

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(10) International Publication Number
WO 02/068409 A1

PCT

(43) International Publication Date
6 September 2002 (06.09.2002)(51) International Patent Classification: C07D 401/12,
471/00, 487/04, A61K 31/444, A61P 29/02

(21) International Application Number: PCT/US02/05226

(22) International Filing Date: 20 February 2002 (20.02.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Date: 23 February 2001 (23.02.2001) US
60271,100(71) Applicant (for all designated States except US): MERCK
& CO., INC. (US); 126 East Lincoln Avenue, Rahway,
NJ 07065-0907 (US).

(72) Inventor: and

(73) Investor/Applicant (for US only): CLAIBORNE,
Christoph F. (US); 126 East Lincoln Avenue,
Rahway, NJ 07065-0907 (US); BUTCHER, John
W. (US); 126 East Lincoln Avenue, Rahway, NJ
07065-0907 (US); CLAREMON, David A. (US);
126 East Lincoln Avenue, Rahway, NJ 07065-0907
(US); LIBBY, Brian E. (US); 126 East Lincoln
Avenue, Rahway, NJ 07065-0907 (US); LIVERSON,
Nigel J. (GB); 126 East Lincoln Avenue, Rahway,
NJ 07065-0907 (US); NUNSON, Peter, M. (US);
126 East Lincoln Avenue, Rahway, NJ 07065-0907
(US); NGUYEN, Kevin T. (US); 126 East Lincoln
Avenue, Rahway, NJ 07065-0907 (US); PHILLIPS,


Published:

with international search report

For two-letter codes and other abbreviations, refer to the "Guide
to the Rules for Coding and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: N-SUBSTITUTED NONARYL-HETEROCYCLIC NMDA/NR2B ANTAGONISTS

(1)

(57) Abstract: Compounds represented by Formula (I) (for
pharmacologically acceptable salts thereof, are effective as
NMDA NR2B antagonists useful for relieving pain.

ccr3 / Merck 2

WO 02/068409

PCT/US02/05226

TITLE OF THE INVENTION

N-SUBSTITUTED NONARYL-HETEROCYCLIC NMDA/NR2B ANTAGONISTS

5 This application claims the benefit of priority of U.S. Patent
Application No. 60/271,100 filed February 23, 2001.

BACKGROUND OF THE INVENTION

Field of the Invention

10 This invention relates to N-substituted nonaryl/heterocyclic compounds.
In particular, this invention relates to N-substituted nonaryl/heterocyclic compounds
that are effective as NMDA NR2B antagonists useful for relieving pain.15 Ions such as glutamate play a key role in processes related to chronic
pain and pain-associated neurotoxicity – primarily by acting through N-methyl-D-
aspartate ("NMDA") receptors. Thus, inhibition of such action – by employing ion
channel antagonists, particularly NMDA antagonists – can be beneficial in the
treatment and control of pain.20 Known NMDA antagonists include ketamine, dextromethan, and 3-(2-
carboxypiperazin-4-yl)-propyl-1-phosphonic acid ("CPP"). Although these
compounds have been reported (J.D. Kristensen, et al., *Pain*, 51:249-253 (1992);
P.K. Eide, et al., *Pain*, 51:221-228 (1995); D.J. Knox, et al., *Anaesthesia Intensive Care*
23:620-622 (1995); and M.B. Max, et al., *Clin. Neuropharmacol.* 18:360-368 (1995))25 to produce symptomatic relief in a number of neuropathies including postherpetic
neuralgia, central pain from spinal cord injury, and phantom limb pain, widespread
use of these compounds is precluded by their undesirable side effects. Such side
effects at analgesic doses include psychotomimetic effects such as dizziness,
headache, hallucinations, dysphoria, and disturbances of cognitive and motor
function. Additionally, more severe hallucinations, sedation, and ataxia are produced
at doses only marginally higher than analgesic doses. Thus, it would be desirable to
provide novel NMDA antagonists that are absent of undesirable side effects or that
produce fewer and/or milder side effects.30 NMDA receptors are heteromeric assemblies of subunits, of which two
major subunit families designated NR1 and NR2 have been cloned. Without being
bound by theory, it is generally believed that the various functional NMDA receptors
in the mammalian central nervous system ("CNS") are only formed by combinations

of NR1 and NR2 subunits, which respectively express glycine and glutamate recognition sites. The NR2 subunit family is in turn divided into four individual subunit types: NR2A, NR2B, NR2C, and NR2D. T. Ishii, et al., *J. Biol. Chem.*, 268:2836-2843 (1993), and D.J. Laurie et al., *Mol. Brain Res.*, 51:23-32 (1997) describe how the various resulting combinations produce a variety of NMDA receptors differing in physiological and pharmacological properties such as ion gating properties, magnesium sensitivity, pharmacological profile, as well as in anatomical distribution.

For example, while NR1 is found throughout the brain, NR2 subunits are differentially distributed. In particular, it is believed that the distribution map for NR2B lowers the probability of side effects while producing pain relief. For example, S.Boyce, et al., *Neuropharmacology*, 38:611-623(1999) describes the effect of selective NMDA NR2B antagonists on pain with reduced side effects. Thus, it would be desirable to provide novel NMDA antagonists that target the NR2B receptor.

U.S. Patent No. 6,020,347 and International Patent Publication WO99/25685 describes 4-substituted 4-piperidine carboxamide derivatives that are antagonists of VLA-4 ("Very Late Antigen-4"). International Patent Publication WO 01/00207 describes substituted pyrimidine compounds that are inhibitors of tyrosine kinases. International Patent Publication WO 00/61551 describes


oxopyrimidinealkanoate compounds that are integrin receptor ligands. International Patent Publication EP 604800 describes Carboxyalkyl-phenyl aminocarbonyl-phenyl-piperidine compounds that are blood platelet aggregation inhibitors. International Patent Publication EP 611660 describes benzimidazoles, xanthines, and analogs as tissue aggregation inhibitors. International Patent Publication EP 771799 and U.S. Patent No 5,861,396 describe purin-6-one derivatives for the treatment of cardiovascular and urogenital diseases. International Patent Publication WO94/21615 describes benzimidazole-piperidine compounds utilized as dopamine D4 antagonists. German Patent No. DE4241632 describes substituted phenyl or cyclohexyl-carboxylic acid derivatives that inhibit cell aggregation.

Phenol compounds described as NMDA antagonists are described in U.S. Patent Nos. 5,306,723 and 5,436,255, and in International Patent Publications WO91/17156, WO92/19502, WO93/02052, WO96/37226, and EP 441506. Benzyl piperidine substituted with phenols or imidazoles are described in Z.-L. Zhou, et al., *J. Medicinal Chemistry*, 42:2993-3000(1999). T.F.Gregory, et al., Poster #94, 218th National Meeting American Chemical Society, New Orleans, Louisiana, August 22-

26, 1999. Other NMDA NR2B selective compounds are described in European Patent Publication EP 787493 and J.N.C. Kew et al., *British J. Pharmacol.*, 123:463(1998). However, there continues to be a need for novel NMDA antagonists that target the NR2B receptor.

SUMMARY OF THE INVENTION

The present invention relates to N-substituted nonary/heterocyclic compounds represented by Formula (I):




(I)

or pharmaceutically acceptable salts thereof. The present invention also forms pharmaceutical compositions utilizing the compounds. Further, this invention includes novel methods to treat pain by utilizing the compounds.

DETAILED DESCRIPTION OF THE INVENTION

The compounds of this invention are represented by Formula (I):



(I)

or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 5-7 membered ring containing 1 or 2 nitrogen ring atoms or an aza bicyclo octane ring;

HeiAr is a 5 or 6 membered heteroaromatic ring containing 1-3 nitrogen ring atoms, or isoxazolyl, thiazolyl, thiadiazolyl, quinolinyl, quinoxalinyl, purinyl, pteridinyl, benzimidazolyl, pyrrolopyrimidinyl, or imidazopyridinyl;

HeiAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

10 A is -C₀₋₄alkyl-;

B is aryl(CH₂)₀₋₃-O-C(O)-, heteroaryl(CH₂)₁₋₃-O-C(O)-, indanyl(CH₂)₀₋₃-O-C(O)-, aryl(CH₂)₁₋₃-C(O)-, aryl-cyclopropyl-C(O)-, heteroaryl-cyclopropyl-C(O)-, heteroaryl(CH₂)₁₋₃-C(O)-, aryl(CH₂)₁₋₃-, heteroaryl(CH₂)₁₋₃-, aryl(CH₂)₁₋₃-NH-C(O)-, aryl(CH₂)₁₋₃-NH-C(NCN)-, aryl(CH₂)₁₋₃-SO₂-, heteroaryl(CH₂)₁₋₃-SO₂-, wherein any of the aryl or heteroaryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

20

In one aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom;

25

HeiAr is a 5 or 6 membered heteroaromatic ring containing 1-3 nitrogen ring atoms, or isoxazolyl, thiazolyl, thiadiazolyl, quinolinyl, quinoxalinyl, purinyl, pteridinyl, benzimidazolyl, pyrrolopyrimidinyl, or imidazopyridinyl;

HeiAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is aryl(CH₂)₀₋₃-O-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

5

In an embodiment of this first aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

10 atom;

HeiAr is a 6 membered heteroaromatic ring containing 1 nitrogen ring atom;

HeiAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

15

A is -C₀₋₄alkyl-;

20

B is aryl(CH₂)₀₋₃-O-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

25

In another embodiment of this first aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

30 atom;

HeiAr is an isoxazolyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

4alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-,
Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is aryl[(CH₂)₀₋₃-O-C(O)-, wherein the aryl is optionally substituted

5 by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

10 In still another embodiment of this first aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

atom;

15 HetAr is a thiadiazolyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-,

20 Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is aryl[(CH₂)₀₋₃-O-C(O)-, wherein the aryl is optionally substituted

by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

25 X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

30 In yet another embodiment of this first aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

atom;

HetAr is a 5 membered heteroaromatic ring containing 2 nitrogen ring atoms;

HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is aryl[(CH₂)₀₋₃-O-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

10 X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

15 In another embodiment of this first aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom;

20 HetAr is a quinolinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is aryl[(CH₂)₀₋₃-O-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

25 X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

30 In still another embodiment of this first aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

atom;
NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

HerAr is a purinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is aryl[(CH₂)₀₋₃-O-C(O)-], wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

15

In yet still another embodiment of this first aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

HerAr is 6 membered heteroaromatic ring containing 2 nitrogen ring

atoms;

HerAr is optionally substituted with 1 or 2 substituents, each

substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is aryl[(CH₂)₀₋₃-O-C(O)-], wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

35

In yet another embodiment of this first aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

5 atom;

HerAr is a thiazolyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano,

methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is aryl[(CH₂)₀₋₃-O-C(O)-], wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

15

In another embodiment of this first aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

atom;

HerAr is a pteridinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano,

methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is aryl[(CH₂)₀₋₃-O-C(O)-], wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

35

In still another embodiment of this first aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

5 NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom;

10 HetAr is a pyrolopyrimidinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -CO-4alkyl-;

15 B is aryl[(CH₂)₀₋₃-O-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

20 In still yet another embodiment of this first aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

atom;

25 HetAr is a imidazopyridinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -CO-4alkyl-;

30 B is aryl[(CH₂)₀₋₃-O-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

5 In yet still another embodiment of this first aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom;

10 HetAr is a benzimidazolyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -CO-4alkyl-;

15 B is aryl[(CH₂)₀₋₃-O-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

20 X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

In a second aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

25 NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom;

HetAr is a 5 or 6 membered heteroaromatic ring containing 1-3 nitrogen ring atoms, or isoxazolyl, thiazolyl, thiadiazolyl, quinolinyl, quinoxalinyl, purinyl, pteridinyl, benzimidazolyl, pyrolopyrimidinyl, or imidazopyridinyl;

30 HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl),

nitro, (C₁-2alkyl)(C₁-2alkyl)NCH₂-, (C₁-2alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀-4alkyl-;

B is aryl[(CH₂)₃-SO₂-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁-4alkyl, C₃-6cycloalkyl, C₁-4alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁-4alkyl, C₁-4alkoxy, NH₂, or X taken with an adjacent bond is =O.

In an embodiment of this second aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

atom;

HeAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring

atoms;

HeAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁-4alkyl, C₁-4alkoxy, C₂-4alkynyl, trifluoromethyl, hydroxy, hydroxyC₁-4alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, -N(C₀-4alkyl)(C₀-4alkyl), nitro, (C₁-2alkyl)(C₁-2alkyl)NCH₂-, (C₁-2alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀-4alkyl-;

B is aryl[(CH₂)₃-SO₂-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁-4alkyl, C₃-6cycloalkyl, C₁-4alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁-4alkyl, C₁-4alkoxy, NH₂, or X taken with an adjacent bond is =O.

In another embodiment of the second aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

atom;

HeAr is a quinazolinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁-4alkyl, C₁-4alkoxy, C₂-4alkynyl, trifluoromethyl, hydroxy, hydroxyC₁-4alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, -N(C₀-4alkyl)(C₀-4alkyl), nitro, (C₁-2alkyl)(C₁-2alkyl)NCH₂-, (C₁-2alkyl)HNCH₂-,

Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀-4alkyl-;

B is aryl[(CH₂)₃-SO₂-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁-4alkyl, C₃-6cycloalkyl, C₁-4alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁-4alkyl, C₁-4alkoxy, NH₂, or X taken with an adjacent bond is =O.

In yet another embodiment of the second aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

atom;

HeAr is purinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁-4alkyl, C₁-4alkoxy, C₂-4alkynyl, trifluoromethyl, hydroxy, hydroxyC₁-4alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, -N(C₀-4alkyl)(C₀-4alkyl), nitro, (C₁-2alkyl)(C₁-2alkyl)NCH₂-, (C₁-2alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀-4alkyl-;

B is aryl[(CH₂)₃-SO₂-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁-4alkyl, C₃-6cycloalkyl, C₁-4alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁-4alkyl, C₁-4alkoxy, NH₂, or X taken with an

adjacent bond is =O.

In still another embodiment of the second aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

atom;

5 HetAr is imidazopyridinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -CO-4alkyl-;

10 B is aryl(CH₂)₃-SO₂-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

15

In yet still another embodiment of the second aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

20 atom;

HetAr is a 6 membered heteroaromatic ring containing 1 nitrogen ring

atom;

25 HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -CO-4alkyl-;

30 B is aryl(CH₂)₃-SO₂-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

35

In a third aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 5 membered ring containing 1 nitrogen ring

5 atom;

HetAr is a 5 or 6 membered heteroaromatic ring containing 1-3 nitrogen ring atoms, or isoxazolyl, thiazolyl, thiadiazolyl, quinolinyl, quinazolinyl, purinyl, pteridinyl, benzimidazolyl, pyrrolopyrimidinyl, or imidazopyridinyl;

10 HetAr is optionally substituted with 1 or 2 substituents; each substituent independently is C₁₋₄alkyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, amino, nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, or NH₂C(O)-;

A is -CO-4alkyl-;

15 B is aryl(CH₂)₃-O-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

20

In an embodiment of the third aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 5 membered ring containing 1 nitrogen ring

atom;

HetAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring

25 atoms;

HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, amino, nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, or NH₂C(O)-;

A is -CO-4alkyl-;

30 B is aryl(CH₂)₃-O-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

35

In another embodiment of the third aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

- 5 NonAr is a nonaromatic 5 membered ring containing 1 nitrogen ring atom;
- HeAr is pteridinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;
- 10 A is -CO-4alkyl-;
- B is aryl(CH₂)₀₋₃-O-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and
- X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.
- 20 In still another embodiment of the third aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein
- NonAr is a nonaromatic 5 membered ring containing 1 nitrogen ring atom;
- 25 HeAr is purinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;
- 30 A is -CO-4alkyl-;
- B is aryl(CH₂)₀₋₃-O-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

5 In still another embodiment of the third aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

- NonAr is a nonaromatic 5 membered ring containing 1 nitrogen ring atom;
- 30 HeAr is benzimidazolyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;
- 15 A is -CO-4alkyl-;
- B is aryl(CH₂)₀₋₃-O-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and
- X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.
- 20 In a fourth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein
- NonAr is an aza bicyclo octane ring;
- 25 HeAr is a 5 or 6 membered heteroaromatic ring containing 1-3 nitrogen ring atoms, or isoxazolyl, thiazolyl, thiadiazolyl, quinolinyl, quinoxalinyl, purinyl, pteridinyl, benzimidazolyl, pyrrolopyrimidinyl, or imidazopyridinyl;
- 30 HeAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;
- A is -CO-4alkyl-;

B is $\text{aryl}(\text{CH}_2)_{0-3}-\text{O}-\text{C}(\text{O})-$, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C_1 -4alkyl, C_3 -6cycloalkyl, C_1 -4alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C_1 -4alkyl, C_1 -4alkoxy, NH_2 , or X taken with an adjacent bond is =O.

In an embodiment of the fourth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

10 NonAr is an aza bicyclo octane ring;

HetAr is a 6 membered heteroaromatic ring containing 1 nitrogen ring atom;

15 HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C_1 -4alkoxy, C_2 -4alkoxy, trifluoromethyl, hydroxy, hydroxy C_1 -4alkyl, fluoro, chloro, bromo, iodo, cyano, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, $-\text{N}(\text{C}_0$ -4alkyl)(C_0 -4alkyl), nitro, (C_1 -2alkyl)(C_1 -2alkyl) NCH_2- , (C_1 -2alkyl) HNCH_2- , $\text{Si}(\text{CH}_3)_3-\text{C}-$, or $\text{NH}_2\text{C}(\text{O})-$;

A is $-\text{C}_0$ -4alkyl-;

20 B is $\text{aryl}(\text{CH}_2)_{0-3}-\text{O}-\text{C}(\text{O})-$, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C_1 -4alkyl, C_3 -6cycloalkyl, C_1 -4alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C_1 -4alkyl, C_1 -4alkoxy, NH_2 , or X taken with an adjacent bond is =O.

25 In another embodiment of the fourth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is an aza bicyclo octane ring;

30 HetAr is purinyl optionally substituted with 1 or 2 substituents, each substituent independently is C_1 -4alkyl, C_1 -4alkoxy, C_2 -4alkoxy, trifluoromethyl, hydroxy, hydroxy C_1 -4alkyl, fluoro, chloro, bromo, iodo, cyano, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, $-\text{N}(\text{C}_0$ -4alkyl)(C_0 -4alkyl),

nitro, (C_1 -2alkyl)(C_1 -2alkyl) NCH_2- , (C_1 -2alkyl) HNCH_2- , $\text{Si}(\text{CH}_3)_3-\text{C}-$, or $\text{NH}_2\text{C}(\text{O})-$;

A is $-\text{C}_0$ -4alkyl-;

5 B is $\text{aryl}(\text{CH}_2)_{0-3}-\text{O}-\text{C}(\text{O})-$, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C_1 -4alkyl, C_3 -6cycloalkyl, C_1 -4alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C_1 -4alkyl, C_1 -4alkoxy, NH_2 , or X taken with an adjacent bond is =O.

10 In still another embodiment of the fourth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is an aza bicyclo octane ring;

HetAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring atom;

15 HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C_1 -4alkyl, C_1 -4alkoxy, C_2 -4alkoxy, trifluoromethyl, hydroxy, hydroxy C_1 -4alkyl, fluoro, chloro, bromo, iodo, cyano, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, $-\text{N}(\text{C}_0$ -4alkyl)(C_0 -4alkyl), nitro, (C_1 -2alkyl)(C_1 -2alkyl) NCH_2- , (C_1 -2alkyl) HNCH_2- , $\text{Si}(\text{CH}_3)_3-\text{C}-$, or $\text{NH}_2\text{C}(\text{O})-$;

A is $-\text{C}_0$ -4alkyl-;

20 B is $\text{aryl}(\text{CH}_2)_{0-3}-\text{O}-\text{C}(\text{O})-$, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C_1 -4alkyl, C_3 -6cycloalkyl, C_1 -4alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

25 X is H, OH, F, C_1 -4alkyl, C_1 -4alkoxy, NH_2 , or X taken with an adjacent bond is =O.

30 In a fifth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is an aza bicyclo octane ring;

HetAr is a 5 or 6 membered heteroaromatic ring containing 1-3 nitrogen ring atoms, or isoxazolyl, thiazolyl, thiadiazolyl, quinolinyl, quinazolinyl, purinyl, pteridinyl, benzimidazolyl, pyrolopyrimidinyl, or imidazopyridinyl;

HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is aryl(CH₂)₁₋₃-SO₂-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

In an embodiment of the fifth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein NonAr is an aza bicyclo octane ring;

HetAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring atom;

HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is aryl(CH₂)₁₋₃-SO₂-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

In a sixth aspect, the compounds of this invention are represented by Formula (II) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

atom;

HetAr is a 5 or 6 membered heteroaromatic ring containing 1-3 nitrogen ring atoms, or isoxazolyl, thiazolyl, thiadiazolyl, quinolinyl, quinoxalinyl, purinyl, pteridinyl, benzimidazolyl, pyrrolopyrimidinyl, or imidazopyridinyl;

HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is heteroaryl(CH₂)₁₋₃-C(O)-, wherein the heteroaryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

In an embodiment of the sixth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

atom;

HetAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring

atom;

HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is heteroaryl(CH₂)₁₋₃-C(O)-, wherein the heteroaryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

In a seventh aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom;

5 HetAr is a 5 or 6 membered heteroaromatic ring containing 1-3 nitrogen ring atoms, or isoxazolyl, thiazolyl, thiadiazolyl, quinolinyl, quinoxalinyl, purinyl, pteridinyl, benzimidazolyl, pyrrolopyrimidinyl, or imidazopyridinyl; HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

15 B is aryl(CH₂)₃-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

20

In an embodiment of the seventh aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

25 atom;

HetAr is 6 membered heteroaromatic ring containing 2 nitrogen ring

atom;

30 HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is aryl(CH₂)₃-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

5 X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

In an eighth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

10 NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom;

HetAr is a 5 or 6 membered heteroaromatic ring containing 1-3 nitrogen ring atoms, or isoxazolyl, thiazolyl, thiadiazolyl, quinolinyl, quinoxalinyl, purinyl, pteridinyl, benzimidazolyl, pyrrolopyrimidinyl, or imidazopyridinyl;

15 HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

20 A is -C₀₋₄alkyl-;

B is aryl-cyclopropyl-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

25 X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

In an embodiment of the eighth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

30 NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom;

HetAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring atoms;

HerAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₂alkyl)(CO₂alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -CO₂alkyl-;

B is aryl-cyclopropyl-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

In an embodiment of the eighth aspect, the compounds of this

invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

atom;

HerAr is a pyrimidinyl ring optionally substituted with 1 or 2

substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₂alkyl)(CO₂alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -CO₂alkyl-;

B is aryl-cyclopropyl-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

In another embodiment of the eighth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

atom;

HerAr is a pyrazinyl ring optionally substituted with 1 or 2

substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₂alkyl)(CO₂alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -CO₂alkyl-;

B is aryl-cyclopropyl-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an

adjacent bond is =O.

In still another embodiment of the eighth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

atom;

HerAr is pyridazinyl ring optionally substituted with 1 or 2

substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₂alkyl)(CO₂alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -CO₂alkyl-;

B is aryl-cyclopropyl-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an

adjacent bond is =O.

In another embodiment of the eighth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring

atom;

HetAr is a pyridyl ring optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl,

trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(Co-4alkyl)(Co-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is aryl-cyclopropyl-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

In a ninth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom;

HetAr is a 5 or 6 membered heteroaromatic ring containing 1-3

nitrogen ring atoms, or isoxazolyl, thiazolyl, thiadiazolyl, quinolinyl, quinazolinyl, purinyl, pteridinyl, benzimidazolyl, pyrrolopyrimidinyl, or imidazopyridinyl;

HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(Co-4alkyl)(Co-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is heteroaryl(CH₂)₃-O-C(O)-, wherein the heteroaryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

In an embodiment of the ninth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom;

HetAr is a 6 membered heteroaromatic ring containing 1 nitrogen ring atom;

HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(Co-4alkyl)(Co-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -C₀₋₄alkyl-;

B is heteroaryl(CH₂)₃-O-C(O)-, wherein the heteroaryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

In a tenth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom;

HetAr is a 5 or 6 membered heteroaromatic ring containing 1-3 nitrogen ring atoms, or isoxazolyl, thiazolyl, thiadiazolyl, quinolinyl, quinazolinyl, purinyl, pteridinyl, benzimidazolyl, pyrrolopyrimidinyl, or imidazopyridinyl;

HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(Co-4alkyl)(Co-4alkyl),

nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -CO₂-alkyl-;

B is aryl(CH₂)₁₋₃-NH-C(NCN)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃-

6cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

10 In an embodiment of the tenth aspect, the compounds of this invention are represented by Formula (I) or pharmaceutically acceptable salts thereof, wherein NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom;

15 HetAr is a 6 membered heteroaromatic ring containing 1 nitrogen ring atom;

HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyl, C₁₋₄alkyl, fluoro, bromo, iodo, cyano, methyl/sulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(C₀₋₄alkyl)(C₀₋₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-;

A is -CO₂-alkyl-;

B is aryl(CH₂)₁₋₃-NH-C(NCN)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃-

6cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro; and

25 X is H, OH, F, C₁₋₄alkyl, C₁₋₄alkoxy, NH₂, or X taken with an adjacent bond is =O.

30 As used herein, "alkyl" as well as other groups having the prefix "alk" such as, for example, alkoxy, alkanoyl, alkenyl, alkynyl and the like, means carbon chains which may be linear or branched or combinations thereof. Examples of alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, ~~sec~~- and ~~tert~~-butyl, pentyl, hexyl, heptyl and the like. "Alkenyl", "alkynyl" and other like terms include carbon chains containing at least one unsaturated C-C bond.

The term "cycloalkyl" means carbocycles containing no heteroatoms, and includes mono-, bi- and tricyclic saturated carbocycles, as well as fused ring systems. Such fused ring systems can include one ring that is partially or fully unsaturated such as a benzene ring to form fused ring systems such as benzofused

5 carbocycles. Cycloalkyl includes such fused ring systems as spirofused ring systems. Examples of cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, decahydronaphthalene, adamantane, indanyl, indenyl, fluorenyl, 1,2,3,4-

tetrahydronaphthalene and the like. Similarly, "cycloalkenyl" means carbocycles containing no heteroatoms and at least one non-aromatic C-C double bond, and include mono-, bi- and tricyclic partially saturated carbocycles, as well as benzofused

10 cycloalkenes. Examples of cycloalkenyl include cyclohexenyl, indenyl, and the like. The term "cycloalkyloxy" unless specifically stated otherwise includes a cycloalkyl group connected to the oxy connecting atom.

The term "alkoxy" unless specifically stated otherwise includes an alkyl group connected to the oxy connecting atom.

15 The term "aryl" unless specifically stated otherwise includes multiple ring systems as well as single ring systems such as, for example, phenyl or naphthyl. The term "aryloxy" unless specifically stated otherwise includes multiple ring systems as well as single ring systems such as, for example, phenyl or

20 naphthyl, connected through the oxy connecting atom to the connecting site. The term "C₀" means that the carbon is not present. Thus, "C₀-C₅" means that there are from none to five carbons present - that is, five, four, three, two, one, or no carbons present. When no carbons are present in a linking alkyl group, the link is a direct bond. When no carbons are present in a terminal alkyl group, the

25 terminus is hydrogen. The term "hetero" unless specifically stated otherwise includes one or more O, S, or N atoms. For example, heterocycloalkyl and heteroaryl include ring systems that contain one or more O, S, or N atoms in the ring, including mixtures of such atoms. The hetero atoms replace ring carbon atoms. Thus, for example, a heterocycloalkyl is a five membered ring containing from 5 to no carbon atoms.

30 Examples of heteroaryl include, for example, pyridinyl, quinolinyl, isoquinolinyl, pyridazinyl, pyrimidinyl, pyrazinyl, quinoxalinyl, furyl, benzofuryl, dibenzofuryl, thienyl, benzthienyl, pyrrolyl, indolyl, pyrazolyl, indazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, benzimidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl.

The term "heteroaryl" unless specifically stated otherwise describes a heteroaryl group connected through an oxygen atom to the connecting site.

Examples of heteroaryl(C₁₋₆alkyl) include, for example, furylmethyl, furyl, thienylmethyl, thienyl, pyrazolylmethyl, oxazolylmethyl, oxazolyl, isoxazolylmethyl, thiazolylmethyl, thiazolyl, imidazolylmethyl, imidazolyl, benzimidazolylmethyl, oxadiazolylmethyl, oxadiazolyl, thiadiazolylmethyl, thiadiazolyl, triazolylmethyl, triazolyl, tetrazolylmethyl, tetrazolyl, pyridinylmethyl, pyridinyl, pyridazinylmethyl, pyrimidinylmethyl, pyrazinylmethyl, quinolinylmethyl, isoquinolinylmethyl and quinoxalylmethyl.

Examples of heterocycloC₃₋₇alkyl include, for example, azetidyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, tetrahydrofuranyl, imidazolyl, pyrrolidin-2-one, piperidin-2-one, and thiomorpholinyl.

Examples of aryl(C₁₋₆alkyl) include, for example, phenyl(C₁₋₆alkyl), and naphthyl(C₁₋₆alkyl).

Examples of heterocycloC₃₋₇alkylcarbonyl(C₁₋₆alkyl) include, for example, azetidyl carbonyl(C₁₋₆alkyl), pyrrolidinyl carbonyl(C₁₋₆alkyl), piperidinyl carbonyl(C₁₋₆alkyl), piperazinyl carbonyl(C₁₋₆alkyl), morpholinyl carbonyl(C₁₋₆alkyl), and thiomorpholinyl carbonyl(C₁₋₆alkyl).

The term "amine" unless specifically stated otherwise includes primary, secondary and tertiary amines.

Unless otherwise stated, the term "carbamoxy" is used to include -NHC(O)OC₁₋₆alkyl, and -OC(O)NHC₁₋₆alkyl.

The term "halogen" includes fluorine, chlorine, bromine and iodine atoms.

The term "optionally substituted" is intended to include both substituted and unsubstituted. Thus, for example, optionally substituted aryl could represent a pentafluorophenyl or a phenyl ring. Further, the substitution can be made at any of the groups. For example, substituted aryl(C₁₋₆alkyl) includes substitution on the aryl group as well as substitution on the alkyl group.

Compounds described herein contain one or more double bonds and may thus give rise to cis/trans isomers as well as other conformational isomers. The present invention includes all such possible isomers as well as mixtures of such isomers.

Compounds described herein may contain one or more asymmetric centers and may thus give rise to diastereomers and optical isomers. The present invention includes all such possible diastereomers as well as their racemic mixtures, their substantially pure resolved enantiomers, all possible geometric isomers, and pharmaceutically acceptable salts thereof. The above Formula I is shown without a definitive stereochemistry at certain positions. The present invention includes all stereoisomers of Formula I and pharmaceutically acceptable salts thereof. Further, mixtures of stereoisomers as well as isolated specific stereoisomers are also included.

During the course of the synthetic procedures used to prepare such compounds, or in using racemization or epimerization procedures known to those skilled in the art, the products of such procedures can be a mixture of stereoisomers.

The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids. When the compound of the present invention is acidic, its corresponding salt can be conveniently prepared from pharmaceutically acceptable non-toxic bases, including inorganic bases and organic bases. Salts derived from such inorganic bases include aluminum, ammonium, calcium, copper (ic and ous), ferric, ferrous, lithium, magnesium, manganese (ic and ous), potassium, sodium, zinc and the like salts. Particularly preferred are the ammonium, calcium, magnesium, potassium and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, as well as cyclic amines and substituted amines such as naturally occurring and synthesized substituted amines. Other pharmaceutically acceptable organic non-toxic bases from which salts can be formed include ion exchange resins such as, for example, arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine and the like.

When the compound of the present invention is basic, its corresponding salt can be conveniently prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Such acids include, for example, acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic,

malic, mandelic, methanesulfonic, mucic, nitric, pantoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid and the like. Particularly preferred are citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, and tartaric acids.

The pharmaceutical compositions of the present invention comprise a compound represented by Formula I (or pharmaceutically acceptable salts thereof) as an active ingredient, a pharmaceutically acceptable carrier and optionally other therapeutic ingredients or adjuvants. The compositions include compositions suitable for oral, rectal, topical, and parenteral (including subcutaneous, intramuscular, and intravenous) administration, although the most suitable route in any given case will depend on the particular host, and nature and severity of the conditions for which the active ingredient is being administered. The pharmaceutical compositions may be conveniently presented in unit dosage form and prepared by any of the methods well known in the art of pharmacy.

In practice, the compounds represented by Formula I, or pharmaceutically acceptable salts thereof, of this invention can be combined as the active ingredient in intimate admixture with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g., oral or parenteral (including intravenous). Thus, the pharmaceutical compositions of the present invention can be presented as discrete units suitable for oral administration such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient. Further, the compositions can be presented as a powder, as granules, as a solution, as a suspension in an aqueous liquid, as a non-aqueous liquid, as an oil-in-water emulsion or as a water-in-oil liquid emulsion. In addition to the common dosage forms set out above, the compound represented by Formula I, or pharmaceutically acceptable salts thereof, may also be administered by controlled release means and/or delivery devices. The compositions may be prepared by any of the methods of pharmacy. In general, such methods include a step of bringing into association the active ingredient with the carrier that constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and intimately admixing the active ingredient with liquid carriers or finely divided solid carriers or both. The product can then be conveniently shaped into the desired presentation.

Thus, the pharmaceutical compositions of this invention may include a pharmaceutically acceptable carrier and a compound or a pharmaceutically acceptable

salt of Formula I. The compounds of Formula I, or pharmaceutically acceptable salts thereof, can also be included in pharmaceutical compositions in combination with one or more other therapeutically active compounds.

The pharmaceutical carrier employed can be, for example, a solid, liquid, or gas. Examples of solid carriers include lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, and stearic acid. Examples of liquid carriers are sugar syrup, peanut oil, olive oil, and water. Examples of gaseous carriers include carbon dioxide and nitrogen.

In preparing the compositions for oral dosage form, any convenient pharmaceutical media may be employed. For example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like may be used to form oral liquid preparations such as suspensions, elixirs and solutions; while carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents, and the like may be used to form oral solid preparations such as powders, capsules and tablets. Because of their ease of administration, tablets and capsules are the preferred oral dosage units whereby solid pharmaceutical carriers are employed. Optionally, tablets may be coated by standard aqueous or nonaqueous techniques.

A tablet containing the composition of this invention may be prepared by compression or molding, optionally with one or more accessory ingredients or adjuvants. Compressed tablets may be prepared by compressing, in a suitable machine, the active ingredient in a free-flowing form such as powder or granules, optionally mixed with a binder, lubricant, inert diluent, surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent. Each tablet preferably contains from about 1mg to about 500mg of the active ingredient and each cachet or capsule preferably containing from about 1 to about 500mg of the active ingredient.

Pharmaceutical compositions of the present invention suitable for parenteral administration may be prepared as solutions or suspensions of the active compounds in water. A suitable surfactant can be included such as, for example, hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof in oils. Further, a preservative can be included to prevent the detrimental growth of microorganisms.

Pharmaceutical compositions of the present invention suitable for injectable use include sterile aqueous solutions or dispersions. Furthermore, the compositions can be in the form of sterile powders for the extemporaneous preparation of such sterile injectable solutions or dispersions. In all cases, the final injectable form must be sterile and must be effectively fluid for easy syringability.

The pharmaceutical compositions must be stable under the conditions of manufacture and storage; thus, preferably should be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol and liquid polyethylene glycol), vegetable oils, and suitable mixtures thereof.

Pharmaceutical compositions of the present invention can be in a form suitable for topical use such as, for example, an aerosol, cream, ointment, lotion, dusting powder, or the like. Further, the compositions can be in a form suitable for use in transdermal devices. These formulations may be prepared, utilizing a compound represented by Formula I of this invention, or pharmaceutically acceptable salts thereof, via conventional processing methods. As an example, a cream or ointment is prepared by mixing hydrophilic material and water, together with about 5 wt% to about 10 wt% of the compound, to produce a cream or ointment having a desired consistency.

Pharmaceutical compositions of this invention can be in a form suitable for rectal administration wherein the carrier is a solid. It is preferable that the mixture forms unit dose suppositories. Suitable carriers include cocoa butter and other materials commonly used in the art. The suppositories may be conveniently formed by first admixing the composition with the softened or melted carrier(s) followed by chilling and shaping in moulds.

In addition to the aforementioned carrier ingredients, the pharmaceutical formulations described above may include, as appropriate, one or more additional carrier ingredients such as diluents, buffers, flavoring agents, binders, surface-active agents, thickeners, lubricants, preservatives (including anti-oxidants) and the like. Furthermore, other adjuvants can be included to render the formulation isotonic with the blood of the intended recipient. Compositions containing a compound described by Formula I, or pharmaceutically acceptable salts thereof, may also be prepared in powder or liquid concentrate form.

Experimental Protocols

Assessing the Activity of Selected Compounds to Inhibit NR1A/2B NMDA Receptor Activation (FLIPR Assay)

5

The activity of selected compounds to inhibit NR1A/2B NMDA receptor activation measured as NR1A/2B receptor-mediated Ca^{2+} influx is assessed by the following procedure:

- 10 NR1A/2B receptor transfected L(tk) cells are plated in 96-well format at 3×10^6 cells per plate and grown for one - two days in normal growth media (Dulbeccos MEM with Na pyruvate, 4500mg glucose, pen/strep, glutamine, 10% FCS and 0.5mg/mL geneticin). NR1A/2B-expression in these cells is induced by the addition of 4nM dexamethasone in the presence of 500 μM ketamine for 16 - 24 hours. After receptor induction cells are washed using a Labsystem Cellwasher two times with assay buffer (Hanks balanced salt solution (HBSS-Mg⁺⁺ free) containing 20mM HEPES, 0.1% BSA, 2mM CaCl₂ and 250 μM probenecid). The cells of each 96 well cell plate are loaded with the Ca^{++} sensitive dye Fluo-3 (Molecular Probes, Inc.) at 4 μM in assay buffer containing 0.5% FBS, and 0.04% pluronic F-127 (Molecular Probes, Inc.) for 1h at 37 °C avoiding light. The cells are then washed with the Cellwasher four times with assay buffer leaving them in 100 μL buffer. Test compounds in solution are pipetted by FLIPR (Fluorometric Imaging Plate Reader) into each test well for a 2min pretreatment. During this time the fluorescence intensity is recorded (excitation at 488nm and emission at 530nm). The glutamate/glycine 50 μL agonist solution (final concentration 1 μM /1 μM) is then added by FLIPR into each well already containing 150 μL of buffer (containing the test compound or vehicle) and the fluorescence is continuously monitored for 10min. The endpoint fluorescence values are used to determine an IC_{50} value comparing the agonist-stimulated signal for the vehicle alone sample and that for the cells incubated with each concentration of test compound.
- 20
- 25
- 30

Determining the Apparent Dissociation Constant (K_i) of Compounds for Human NR1A/NR2B Receptors (Binding Assay):


The radioligand binding assay is performed at room temperature in 96-well microtiter plates with a final assay volume of 1.0mL in 20mM HEPES buffer (pH 7.4) containing 150mM NaCl. Solutions of test compounds were prepared in DMSO and serially diluted with DMSO to yield 20μL of each of 10 solutions differing by 3-fold in concentration. Non-specific binding (NSB) using hot AMD-1 (10μM final concentration) and total binding (TB) by using DMSO (2% final concentration). A solution of NR1A/NR2B receptors (40pM final concentration) and tritiated AMD-2 (1nM final concentration) were added to the test compounds. After 3h of incubation at room temperature, samples are filtered through Packard GF/B filters (presoaked in 0.05% PEI polyethylenimine Sigma P-3143) and washed 10 times with 1mL of cold 20mM HEPES buffer per wash. After vacuum drying of the filter plates, 40μL of Packard Microscint-20 was added and bound radioactivity determined in a Packard TopCount. The apparent dissociation constant (K_i), the maximum percentage inhibition (%I_{max}), the minimum percentage inhibition (%I_{min}) and the hill slope (nH) were determined by a non-linear least squares fitting the bound CPM data to Equation #1 below.

Equation #1:


$$\text{CPM Bound} = \frac{(\text{SB}) (\%I_{\text{max}} - \%I_{\text{min}})}{(1 + ([\text{Drug}]/(K_i(\text{AMD-2})/K_0))^{\text{nH}})} + \text{NSB} + (\text{SB}) (1 - \%I_{\text{max}})$$

where, K₀ is the apparent dissociation constant for the radioligand for the receptor as determined by hot saturation and SB is the specifically bound CPM determined from the difference of TB and NSB.


AMD-1



AMD-2




Compounds AMD-1 and AMD-2 can be synthesized in accordance with the following general reaction schemes.




In accordance with scheme 1, hydrogen chloride is bubbled through a solution of the appropriately substituted benzamide 1 in methanol at room temperature. The volatiles are removed under reduced pressure and the resulting residue is triturated with ether and filtered to yield the desired imidate 2. Imidate 2 is dissolved in methanol at ambient temperature, treated with amine 3 at ambient temperature and stirred under argon. The volatiles are removed under reduced pressure and the residue purified by preparative HPLC or trituration with ether to afford amidine 1a.

15



In accordance with scheme 2, at room temperature under argon, amine 3a is dissolved in ether and was treated with 1-M hydrogen chloride in ether (1 equiv.) in a single portion. The resulting precipitate is stirred vigorously for 10 minutes. The volatiles are removed under reduced pressure. The residue is suspended in toluene, cooled to 0°C under argon, treated with 2.0-M trimethylaluminum (1.05 equiv.) in a dropwise manner, and stirred for 45 minutes at room temperature to afford intermediate 6 (not isolated). Compound 6 is added to a solution of nitrile 1 in toluene. The reaction is heated to 80°C without stirring in a sealed tube for 18h, cooled to ambient temperature, poured onto a silica gel column and eluted with methanol/dichloromethane to give amidine 4.

Preparation of [¹²⁵I]AMD-1



Tritiated AMD-1 was prepared by the following procedure: A mixture of AMD-1, hydrochloride salt, (5mg, 0.012mmol) in dioxane (0.2mL) containing triethylamine (4μL) was treated with hexamethylditin (5μL), a catalytic amount of palladium catalyst and heated at 100°C for 45 minutes. The reaction was cooled to room temperature, filtered through a glass wool plug, rinsed with methanol and concentrated *in vacuo* to give 10.7mg of a brown oil. The oil was dissolved in methylene chloride and passed through a small silica column eluting with methylene chloride followed by 5% methanol/methylene chloride. Fractions containing the trimethylstannane (Rf 0.26 in 10% methanol/methylene chloride) were pooled and

concentrated *in vacuo* to give 4.5mg of the trimethylstannane as a clear colorless oil. This material was further purified by HPLC (C18 Econosil, 10x250mm, 20 minute linear gradient, 30% MeCN:70% H₂O (0.1% TFA) to 90% MeCN, 3mL/min, 254nm, retention time 15 minutes) to give 3mg of the trimethylstannane.

A Na¹²⁵I shipping vial (10mCi, Amersham) was charged with a stir bar, an iodobead, 50μL of methanol and stirred five minutes at room temperature. A solution of the trimethylstannane (0.1mg) in 50μL of methanol containing 5μL of trifluoroacetic acid was added and the reaction was stirred for five minutes. The reaction was quenched with 50μL of ammonium hydroxide and purified by HPLC (C18 Vydac protein and peptide column, 4.6 x 250 mm, 20 minute linear gradient, 30% MeCN:70% H₂O (0.1% TFA) to 90% MeCN, 1 mL/min, retention time 11 minutes). Fractions containing the radioactive product were pooled and concentrated *in vacuo* to give 989μCi of [¹²⁵I]AMD-1 with a specific activity of 898Ci/mmol as measured by UV absorbance at 272nm.

Synthesis of Tritiated AMD-2

Tritiated AMD-2 was prepared by the following procedure: The phenol of AMD-2 (2mg, 0.008mmol) dissolved in dimethylformamide (0.6mL) and potassium carbonate (1.2mg) for 1h. High specific activity tritiated methyl iodide (50mCi, 0.0006mmol, in toluene 1mL, American Radiolabeled Chemicals) was added at room temperature and stirred for 2 hours. The reaction mixture was filtered using a Whatman PTFE 0.45μm syringeless filter device to remove any insoluble potassium carbonate, washed with Abs. ethanol (2mL, Pharmco), and the combined filtrates were concentrated to dryness at room temperature using a rotary evaporator; this also removed any unreacted tritiated methyl iodide. The residue was purified by HPLC chromatography on a Phenomenex Luna C8 semi-prep column (Luna 5 micro C8(2), 250x10.0 mm) using a gradient system of 20/80 acetonitrile/water with 0.1% trifluoroacetic acid to 100% acetonitrile with 0.1% trifluoroacetic acid in 20min. Total activity of the product was 8mCi. Further purification was effected by absorption onto a Waters C-18 Sep-pak column (Waters Sep-Pak PLUS C18) and elution with water followed by absolute ethanol. The product was diluted with absolute ethanol (10mL) before submission for final analysis.

The compounds of this invention exhibit IC₅₀'s of less than 50μM in the FLIPR and binding assays. It is advantageous that the IC₅₀'s be less than 5μM in

the FLIPR and binding assays. It is more advantageous that the IC₅₀'s be less than 1 μM in the FLIPR and binding assays. It is still more advantageous that the IC₅₀'s be less than 0.1 μM in the FLIPR and binding assays. Thus, the compounds and pharmaceutical compositions of this invention have been found to exhibit biological activity as NMDA NR2B antagonists. Accordingly, another aspect of the invention is the treatment of pain, migraine, depression, anxiety, schizophrenia, Parkinson's disease, or stroke - maladies that are amenable to amelioration through inhibition of NMDA NR2B receptors - by the administration of an effective amount of the compounds of this invention.

Thus, pain can be treated by administering once or twice a day, a compound of this invention at 0.1mg, 1mg, 5mg, 10mg, or 25mg per kg of body weight.

Migraine can be treated by administering once or twice a day, a compound of this invention at 0.1mg, 1mg, 5mg, 10mg, or 25mg per kg of body weight.

Depression can be treated by administering once or twice a day, a compound of this invention at 0.1mg, 1mg, 5mg, 10mg, or 25mg per kg of body weight.

Anxiety can be treated by administering once or twice a day, a compound of this invention at 0.1mg, 1mg, 5mg, 10mg, or 25mg per kg of body weight.

Schizophrenia can be treated by administering once or twice a day, a compound of this invention at 0.1mg, 1mg, 5mg, 10mg, or 25mg per kg of body weight.

Parkinson's disease can be treated by administering once or twice a day, a compound of this invention at 0.1mg, 1mg, 5mg, 10mg, or 25mg per kg of body weight.

Stroke can be treated by administering once or twice a day, a compound of this invention at 0.1mg, 1mg, 5mg, 10mg, or 25mg per kg of body weight.

The abbreviations used herein are as follows unless specified otherwise:

BH₃*THF Tetrahydrofuran/borane complex

2,2'-Bis(diphenylphosphino)-1,1'-binaphthyl	BINAP
t-Butoxycarbonyl	BOC
t-Butoxycarbonyl anhydride	BOC ₂ O
Carbobenzyloxy	CBZ
Carbobenzyloxy chloride	CBZ-Cl
Dichloromethane	DCM
Diisopropylethylamine	DIEPA
4-Dimethylaminopyridine	DMAP
N,N-Dimethylformamide	DMF
Dimethylformamide-Dimethylacetal	DMF-DMA
Dimethylsulfoxide	DMSO
3-Ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride	EDC
hours	h
1-Hydroxy-7-azabenzotriazole	HOAt
Hydroxybenzoxazole	HOBt
Isopropanol	IPA
meta Chloroperobenzoic acid	mCPBA
minutes	min
Acetonitrile	MeCN
nuclear magnetic resonance	NMR
room temperature	r.t., RT, or rt
saturated	sat.
Triethylamine	TEA
Trifluoroacetic acid	TFA
Tetrahydrofuran	THF


The following examples are provided to more fully illustrate the present invention, and are not to be construed as limiting the scope of the claims in any manner.

EXAMPLES

The compounds of this invention can be prepared by procedures shown below.

Intermediates:**INTERMEDIATE 1a:**

Carbonic acid 2,5-dioxo-pyrrolidin-1-yl ester 4-methyl-benzyl ester



5

Diisuccinimidy carbonate (5.03g, 19.65mmol) in 30mL MeCN and 30mL DCM was treated with 4-methylbenzyl alcohol (2.4g, 19.6mmol) followed by DMAP (1.20g, 9.82mmol). The resulting cloudy reaction mixture cleared over 2min, stirred overnight at rt, then poured into 100mL water and partitioned. The organic layer was dried over anhydrous sodium sulfate and the solvent evaporated. The solid thus obtained was stirred with approx. 25mL ether, filtered, washed with a small volume of ether and dried to yield carbonic acid 2,5-dioxo-pyrrolidin-1-yl ester 4-methyl-benzyl ester as a white solid. Ref: *Chem. Pharm. Bull.*, 38(1):110-115(1990).

10

15 The following compounds were prepared in the manner similar to that described above for **INTERMEDIATE 1a**:

INTERMEDIATE 1b:

Carbonic acid 2,5-dioxo-pyrrolidin-1-yl ester 4-chloro-benzyl ester

20

INTERMEDIATE 1c:

Carbonic acid 2,5-dioxo-pyrrolidin-1-yl ester 4-fluoro-benzyl ester

INTERMEDIATE 1d:


Carbonic acid 2,5-dioxo-pyrrolidin-1-yl ester 4-ethyl-benzyl ester

INTERMEDIATE 1e:

Carbonic acid 2,5-dioxo-pyrrolidin-1-yl ester 4-isopropyl-benzyl ester

25

Utilizing the carbonic acid derivatives described above for **INTERMEDIATES 1a-1e**, and following the procedure described below in **EXAMPLE 13**, step 1, the following **INTERMEDIATES 2a-2e** were obtained

**INTERMEDIATE 2a:**

4-Methylbenzyl 4-(aminomethyl)piperidine-1-carboxylate

INTERMEDIATE 2b:

4-Chlorobenzyl 4-(aminomethyl)piperidine-1-carboxylate

5

INTERMEDIATE 2c:

4-Fluorobenzyl 4-(aminomethyl)piperidine-1-carboxylate

INTERMEDIATE 2d:

4-Ethylbenzyl 4-(aminomethyl)piperidine-1-carboxylate

10

INTERMEDIATE 2e:


4-Isopropylbenzyl 4-(aminomethyl)piperidine-1-carboxylate

EXAMPLE 1:

Benzyl 4-[(4-pyridinylamino)methyl]-1-piperidinecarboxylate:

15 **Step 1:**

Benzyl 4-[(4-pyridinylamino)carbonyl]-1-piperidinecarboxylate



In DMF (5mL), 1-[(benzyloxy)carbonyl]-4-piperidinecarboxylic acid (P. E. Malignes et al., *Tetrahedron*, 53:10983(1997)) (1.00g, 3.80mmol), 4-aminopyridine (572mg, 6.08mmol), EDC (801mg, 4.18mmol), and HOAt (569mg, 4.18mmol) were combined and aged under N₂ for 4h. The reaction was partitioned


20

between sat. NaHCO₃ and ethyl acetate. The layers were separated and the aqueous layer was extracted with ethyl acetate (2x). The combined organics were washed with water and brine then dried over Na₂SO₄, filtered and concentrated under reduced pressure, affording 1.16g of benzyl 4-[(4-pyridinylamino)carbonyl]-1-piperidinecarboxylate as a yellow oil which was used without further purification.

25

Step 2:

Benzyl 4-[(4-pyridinylamino)methyl]-1-piperidinecarboxylate




The amide prepared as described in Step 1 above (17.82g, 52.50mmol) was dissolved in THF (50mL) and was treated with $\text{BH}_3 \cdot \text{THF}$ (200mmol, 200mL, 1M in THF) over 10min. and was aged at r.t. 3h. The reaction was quenched by slowly adding 2N HCl and stirring vigorously 15h. The reaction was basified with 1M NaOH and extracted with ethyl acetate (3x). The combined organics were washed with brine, dried over Na_2SO_4 , filtered and concentrated *in vacuo*, yielding a white foam which was purified by silica gel chromatography (99:1.0.1 to 90:10:1 CH_2Cl_2 : CH_3OH : NH_4OH) to give 11.53g of benzyl 4-[(4-pyridinylamino)methyl]-1-piperidinecarboxylate

as a viscous pale yellow oil.

^1H NMR (HCl salt 400MHz, CD_3OD): δ 8.09 (brs, 1H, Pyr-H), 7.97 (brs, 1H, Pyr-H), 7.35-7.28 (m, 5H, Ar-H), 6.88 (brs, 2H, Pyr-H), 5.11 (s, 2H, $\text{CH}_2\text{-Ar}$), 4.18 (brd, $J=11.70\text{Hz}$, 2H, CHH), 3.25 (d, $J=6.77\text{Hz}$, 2H, $\text{CH}_2\text{-N}$), 2.86 (brs, 2H, CHH), 1.90-1.77 (m, 3H, CHH, CH), 1.29-1.16 (dq, $J=12.36\text{Hz}$, 4.16Hz, 2H, CHH). M.S. (M+1): 326.47.

EXAMPLE 2:

4-[(3-Methylpyridin-4-ylamino)methyl]piperidine-1-carboxylic acid benzyl ester:




The title compound was prepared as described in EXAMPLE 1, but replacing 4-aminopyridine with 4-amino-3-methylpyridine (Malinowski et al., *J. Prakt. Chem.*, 330:154-158(1988)).

^1H NMR (400MHz, CD_3OD): δ 7.74 (d, $J=5.85\text{Hz}$, 1H, Pyr-H), 7.66 (brs, 1H, Pyr-H), 7.36-7.29 (m, 5H, Ar-H), 6.77 (brs, 1H, Pyr-H), 5.11 (s, 2H, $\text{CH}_2\text{-Ar}$), 4.19 (brd, $J=13.81\text{Hz}$, 3H), 3.31-3.20 (m, 2H, $\text{CH}_2\text{-N}$ + CH_2OH), 2.84 (brs, 2H,

CHH), 2.22 (brs, 2H, CHH), 1.98-1.85 (m, 1H, CH), 1.82 (brd, $J=12.89\text{Hz}$, 2H, CHH), 1.22-1.14 (m, 2H, CHH). M.S. (M+1): 340.27.

5 EXAMPLE 3:

Benzyl 4-[(2-pyridinyl)amino]methyl]-1-piperidinecarboxylate




The title compound was prepared as described in EXAMPLE 1, but replacing 4-aminopyridine with 2-aminopyridine.

^1H NMR (400MHz, CD_3OD): δ 10.00 (brs, 1H, NH), 7.82-7.75 (m, 2H, Pyr-H, Pyr-H), 7.38-7.30 (m, 5H, Ar-H), 6.76-6.70 (m, 2H, Pyr-H, Pyr-H), 5.12 (s, 2H, $\text{CH}_2\text{-Ar}$), 4.24 (brs, 2H, CHH), 3.16 (brs, 2H, $\text{CH}_2\text{-N}$), 2.84 (brs, 2H, CHH), 2.01-1.80 (m, 3H, CH, CHH + H₂O), 1.26-1.18 (m, 2H, CHH). M.S. (M+1): 326.28.

15

EXAMPLE 4:

Benzyl 4-[(3-pyridinyl)amino]methyl]-1-piperidinecarboxylate




The title compound was prepared as described in EXAMPLE 1, but replacing 4-aminopyridine with 3-aminopyridine.

^1H NMR (500MHz, CD_3OD): δ 8.01 (d, $J=2.93\text{Hz}$, 1H, Pyr-H), 7.95 (dd, $J=4.63\text{Hz}$, 1.46Hz, 1H, Pyr-H), 7.37-7.30 (m, 5H, Ar-H), 7.08 (dd, $J=8.30\text{Hz}$, 4.59Hz, 1H, Pyr-H), 6.86-6.84 (m, 1H, Pyr-H), 5.13 (s, 2H, $\text{CH}_2\text{-Ar}$), 4.25 (brs, 2H, CHH), 3.80 (brt, $J=5.86\text{Hz}$, 1H, NH), 3.04 (t, $J=6.33\text{Hz}$, 2H, $\text{CH}_2\text{-N}$), 2.78 (brs, 2H, CHH), 1.78 (brs, 3H, CH, CHH + H₂O), 1.27-1.13 (m, 2H, CHH). M.S. (M+1): 326.31.

25

EXAMPLE 5:

Benzyl 4-[[[(4-methyl-2-pyridinyl)amino]methyl]-1-piperidinecarboxylate




5

The title compound was prepared as described in **EXAMPLE 1**, but replacing 4-aminopyridine with 2-amino-4-methylpyridine (Fluka Co.). M.S. (M+): 340.40.

EXAMPLE 6:

Benzyl 4-[[[(4-ethyl-2-pyridinyl)amino]methyl]-1-piperidinecarboxylate




15

The title compound was prepared as described in **EXAMPLE 1**, but replacing 4-aminopyridine with 2-amino-4-ethylpyridine (Maybridge Chemicals). M.S. (M+): 354.41.

EXAMPLE 7:

Benzyl 4-[(3-isoxazolylamino)methyl]-1-piperidinecarboxylate




20

The title compound was prepared as described in **EXAMPLE 1**, but replacing 4-aminopyridine with 3-aminoisoxazole (Sigma-Aldrich Co.). M.S. (M+): 316.29.

EXAMPLE 8:

Benzyl 4-[(1,3,4-thiadiazol-2-ylamino)methyl]-1-piperidinecarboxylate




The title compound was prepared as described in **EXAMPLE 1**, but replacing 4-aminopyridine with 2-amino-1,3,4-thiadiazole. M.S. (M+): 333.35.

10

EXAMPLE 9:

Benzyl 4-[[[(5-methyl-2-pyridinyl)amino]methyl]-1-piperidinecarboxylate




15

The title compound was prepared as described in **EXAMPLE 1**, but replacing 4-aminopyridine with 2-amino-5-methylpyridine. M.S. (M+): 340.40.

EXAMPLE 10:

Benzyl 4-[[[(1-methyl-1H-imidazol-2-yl)amino]methyl]-1-piperidinecarboxylate




20

The title compound was prepared as described in **EXAMPLE 1**, step 1, but replacing 4-aminopyridine with 2-amino-imidazole hemisulfate and gave the EDC coupling product. This product was refluxed in DMF-DMA for 90min., diluted with ethyl acetate, washed with sat. NaHCO₃, dried over Na₂SO₄, filtered and then concentrated under reduced pressure. The resulting red oil was purified by silica gel chromatography. 50mg (mmol) of the purified product was reacted with borane as described in **EXAMPLE 1**, step 2, to give 26mg of benzyl 4-[[[(1-methyl-1H-imidazol-2-yl)amino]methyl]-1-piperidinecarboxylate.

¹H NMR (400MHz, CDCl₃): 8.73-6.727 (m, 5H, Ar-H), 6.65 (d, J=1.55Hz, 1H, imidazole-H), 6.49 (d, J=1.56Hz, 1H, imidazole-H), 5.12 (s, 2H, CH₂-Ar), 4.19 (brs, 2H, CHH), 3.58 (brs, 1H, NH), 3.34 (s, 3H, CH₃), 3.23 (m, 2H, CH₂-N), 2.79 (brs, 2H, CHH), 1.85-1.70 (m, 3H, CHH, CH), 1.23-1.13 (m, 2H, CHH). M.S. (M+1): 329.27.

EXAMPLE 11: 4-(Quinolin-4-ylaminomethyl)-piperidine-1-carboxylic acid benzyl ester




The title compound was prepared as described in **EXAMPLE 1**, replacing 4-aminopyridine with 4-aminoquinoline. M.S. (M+1): 376.39.

EXAMPLE 12:

Benzyl 4-[[[(1-oxido-4-pyridinyl)amino]methyl]-1-piperidinecarboxylate

Step 1:


Benzyl 4-[[[(1-oxido-4-pyridinyl)amino]carbonyl]-1-piperidinecarboxylate



Benzyl 4-[[[(1-oxido-4-pyridinyl)amino]carbonyl]-1-piperidinecarboxylate (**EXAMPLE 1**, Step 1) (615mg, 1.81mmol) was dissolved in CH₂Cl₂ and treated with mCPBA (3.12g, 18.10mmol) and aged 18h. The reaction was diluted with ethyl acetate and washed with sat. NaHCO₃. The organics were separated, dried over Na₂SO₄, filtered and concentrated under reduced pressure. The resulting oil was purified by silica gel chromatography to afford benzyl 4-[[[(1-oxido-4-pyridinyl)amino]carbonyl]-1-piperidinecarboxylate as a clear oil.

¹H NMR (400MHz, CDCl₃): 8.10-7.72 (s, 1H, NH), 8.03 (d, J=7.50Hz, 2H, Pyr-H), 7.80 (d, J=7.50Hz, 2H, Pyr-H), 7.38-7.28 (m, 5H, Ar-H), 5.12 (s, 2H, CH₂-Ar), 4.18 (brd, J=13.25Hz, 2H, CHH), 2.81 (brs, 2H, CHH), 2.57-2.45 (m, 1H, CH), 1.86-1.68 (m, 4H, CHH, CHH). M.S. (M+1): 356.28.

Step 2:
Benzyl 4-[[[(1-oxido-4-pyridinyl)amino]methyl]-1-piperidinecarboxylate



Benzyl 4-[[[(1-oxido-4-pyridinyl)amino]carbonyl]-1-piperidinecarboxylate (62mg, 0.17mmol) was reduced with borane as described in

EXAMPLE 1, step 2, to afford benzyl 4-[[[(1-oxido-4-pyridinyl)amino]methyl]-1-piperidinecarboxylate as a clear oil.


¹H NMR (400MHz, CDCl₃): 8.799 (d, J=7.31Hz, 2H, Pyr-H), 7.88 (brs, 1H, NH), 7.38-7.30 (m, 5H, Ar-H), 6.66 (brs, 2H, Pyr-H), 5.12 (s, 2H, CH₂-Ar), 4.22 (brs, 2H, CHH), 3.09 (brs, 2H, CH₂-N), 2.77 (brs, 2H, CHH), 1.87-1.71 (m, 3H, CHH, CH), 1.26-1.11 (m, 2H, CHH). M.S. (M+1): 342.33.

EXAMPLE 13:

Benzyl 4-[(9H-purin-6-ylamino)methyl]-1-piperidinecarboxylate

Step 1:

Benzyl 4-(aminomethyl)piperidine-1-carboxylate




4-Aminomethylpiperidine (40g, 350mmol) and benzaldehyde (37.3mL, 368mmol) in toluene (600mL) were heated to reflux under Dean Stark conditions for 2h. The resulting reaction mixture was cooled to room temperature and 500mL dichloromethane was added. The resulting solution was cooled to 5°C and treated with N-(benzyloxycarbonyloxy)succinimide (91.7g, 368mmol). After 10min, the cooling bath was removed and the resulting reaction mixture stirred for 1h. The solvents were evaporated and the residue stirred with 400mL THF and 400mL 2M HCl for 1h. The mixture was concentrated to remove organics and extracted with ether (3x300mL). The aqueous phase was adjusted to pH14 with 50% NaOH and extracted with ethyl acetate. The organic layer was washed with water and brine, dried over anhydrous sodium sulfate, and the solvent evaporated to give benzyl 4-(aminomethyl)piperidine-1-carboxylate as an oil.

¹H NMR (500MHz CDCl₃) δ: 7.4-7.2 (m, 5H); 5.12 (s, 2H); 4.20 (brs, 2H); 2.77 (brs, 2H); 2.58 (d, J=6.6 Hz, 2H) 1.9-1.7 (m, 2H); 1.0-1.5 (m, 5H).

Step 2:

Benzyl 4-[(9H-purin-6-ylamino)methyl]-1-piperidinecarboxylate



In DMF (5mL), benzyl 4-(aminomethyl)-1-piperidinecarboxylate (1.20g, 4.83mmol) and 6-chloropurine (448mg, 2.49mmol) were combined and treated with TEA in a single portion and aged under N₂ at 100°C for 18h. The resulting reaction was diluted with sat. NaHCO₃ and extracted with ethyl acetate (3x). The combined organics were washed with brine, dried over Na₂SO₄, filtered and

concentrated *in vacuo* to give a brown oil which was purified by silica gel chromatography (20g, 32-60μm silica, 99:1:0.1 to 90:10:1 CH₂Cl₂:CH₃OH:NH₄OH) to give benzyl 4-[(9H-purin-6-ylamino)methyl]-1-piperidinecarboxylate as a brown oil.

¹H NMR (400MHz, CDCl₃): δ 8.42 (s, 1H, purine-H), 7.97 (s, 1H, purine-H), 7.36-7.29 (m, 5H, Ar-H), 6.21 (brs, 1H), 5.13 (s, 2H, CH₂-Ar), 4.22 (brs, 2H, CHH), 3.43 (brs, 2H, CH₂-N), 2.80 (brs, 2H, CHH), 1.95-1.79 (m, 3H, CHH, CH), 1.34-1.21 (m, 2H, CHH).


M.S. (M+1): 367.31.

EXAMPLE 14:

4-Methylbenzyl 4-[(4-pyrimidinylamino)methyl]-1-piperidinecarboxylate

Step 1:


4-[(2-Methylsulfonyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid 4-methyl-benzyl ester



The 4-[(2-methylsulfonyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid 4-methyl-benzyl ester was prepared as described in EXAMPLE 13, Step 2, but replacing 6-chloropurine with 4-chloro-2-methylthiopyrimidine and replacing benzyl 4-(aminomethyl)-1-piperidinecarboxylate with 4-methylbenzyl 4-(aminomethyl)-1-piperidinecarboxylate. M.S. (M+1): 387

Step 2:

4-Methylbenzyl 4-[(4-pyrimidinylamino)methyl]-1-piperidinecarboxylate




4-[(2-Methylsulfonyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid 4-methyl-benzyl ester (550mg, 1.42mmol) was dissolved in EtOH (15mL) and treated with Raney Nickel (834mg, 14.20mmol) at room temperature for 3h, filtered, concentrated and purified by silica gel chromatography to give

- 5 **EXAMPLE 14** as a yellow oil.
¹H NMR (400MHz, CDCl₃): δ 8.53 (s, 1H, Pyr-H), 8.13 (brd, J=4.8Hz, 1H, Pyr-H), 7.24 (d, J=7.86Hz, 2H, Ar-H), 7.16 (d, J=7.68Hz, 2H, Ar-H), 6.31 (dd, J=6.00Hz, 1H, Pyr-H), 5.57 (s, 1H, NH), 5.08 (s, 2H, CH₂-Ar), 4.20 (brs, 2H, CHH), 3.23 (brs, 2H, CH₂-N), 2.75 (brs, 2H, CHH), 2.34 (s, 3H, CH₃), 1.82-1.65 (m, 3H, CHH, CH), 1.23-1.09 (m, 2H, CHH).
 M.S. (M+1): 341.35.

EXAMPLE 15:

Benzyl 4-[(4-pyrimidinylamino)methyl]-1-piperidinecarboxylate




The title compound was prepared as described in **EXAMPLE 14**, but replacing 4-methylbenzyl 4-(aminomethyl)-1-piperidinecarboxylate with benzyl 4-(aminomethyl)-1-piperidinecarboxylate.

- 20 ¹H NMR (400MHz, CDCl₃): δ 8.53 (s, 1H, Pyr-H), 8.13 (brd, J=4.85Hz, 1H, Pyr-H), 7.38-7.28 (m, 5H, Ar-H), 6.32 (d, J=6.03Hz, 1H, Pyr-H), 5.51 (brs, 1H, NH), 5.12 (s, 2H, CH₂-Ar), 4.21 (brs, 2H, CHH), 3.24 (brs, 2H, CH₂-N), 2.77 (brs, 2H, CHH), 1.85-1.70 (m, 3H, CHH, CH), 1.27-1.10 (m, 2H, CHH).
 M.S. (M+1): 327.29.

EXAMPLE 16:

Benzyl 4-[(2-pyrimidinylamino)methyl]-1-piperidinecarboxylate




The title compound was prepared as described in **EXAMPLE 13**, except using benzyl 4-(aminomethyl)-1-piperidinecarboxylate (6.50g, 26.19mmol) and 2-chloropyrimidine (990mg, 8.64mmol) as starting materials without a solvent to give the title compound as a yellow oil.

- 5 ¹H NMR (400MHz, CDCl₃): δ 8.26 (d, J=4.85Hz, 1H, Pyr-H), 7.36-7.29 (m, 5H, Ar-H), 6.52 (t, J=4.85Hz, 1H, Pyr-H), 5.12 (s, 2H, CH₂-Ar), 4.21 (brs, 2H, CHH), 3.30 (t, J=6.26Hz, 2H, CH₂-N), 2.78 (brs, 2H, CHH), 1.76-1.62 (m, 3H, CHH, CH), 1.28-1.12 (m, 2H, CHH).
 M.S. (M+1): 327.33.

EXAMPLE 17:

4-Methylbenzyl 4-[(2-pyrimidinylamino)methyl]-1-piperidinecarboxylate




- 15 The title compound was prepared as described in **EXAMPLE 13**, except using 4-methylbenzyl 4-(aminomethyl)-1-piperidinecarboxylate (300mg, 1.14mmol), 2-chloropyrimidine (131mg, 1.14mmol) as starting materials gave the title compound as a yellow oil.

- 20 ¹H NMR (400MHz, CDCl₃): δ 8.26 (d, J=4.76, 2H, Pyr-H), 7.26 (d, J=8.96Hz, 2H, Ar-H), 7.17 (d, J=8.96Hz, 2H, Ar-H), 6.31 (dd, J=4.85Hz, 1H, Pyr-H), 5.28 (s, 1H, NH), 5.08 (s, 2H, CH₂-Ar), 4.19 (brs, 2H, CHH), 3.32 (d, J=6.36Hz, 2H, CH₂-N), 2.76 (brs, 2H, CHH), 2.35 (s, 3H, CH₃), 1.82-1.60 (m, 3H, CHH, CH), 1.25-1.13 (m, 2H, CHH).
 M.S. (M+1): 341.37.

EXAMPLE 18:

Benzyl 4-[(5-methyl-2-pyrimidinyl)amino]methyl]-1-piperidinecarboxylate



The title compound was prepared as described in **EXAMPLE 13**,


except using benzyl 4-(aminomethyl)-1-piperidinecarboxylate (298mg, 1.20mmol), 2-chloro-5-methylpyrimidine (**EXAMPLE 144, Step 1**) (51mg, 0.40mmol) as starting materials and using no solvent and gave the title compound as a yellow oil.

¹H NMR (400MHz, CDCl₃): δ 8.10 (s, 2H, Pyr-H), 7.36-7.28 (m, 5H, Ar-H), 5.47 (br, J=4.98Hz, 1H, NH), 5.12 (s, 2H, CH₂-Ar), 4.19 (brs, 2H, CHH), 3.32 (d, J=6.22Hz, 2H, CH₂-N), 2.76 (brs, 2H, CHH), 2.10 (s, 3H, CH₃), 1.82-1.63 (m, 3H, CHH, CH), 1.25-1.12 (m, 2H, CHH).

M.S. (M+1): 341.40.

EXAMPLE 19:

4-Methylbenzyl 4-((2-(methylsulfany)-4-pyrimidinyl)amino]methyl)-1-piperidinecarboxylate



The title compound was prepared as described in **EXAMPLE 13**,


except using 4-methylbenzyl 4-(aminomethyl)-1-piperidinecarboxylate (600mg, 2.29mmol), and 4-chloro-2-methylthiopyrimidine (386mg, 2.40mmol) as starting materials and gave the title compound as a yellow oil.

¹H NMR (400MHz, CDCl₃): δ 7.99 (bs, 1H, Pyr-H), 7.25 (d, J=8.69Hz, 2H, Ar-H), 7.17 (d, J=8.95Hz, 2H, Ar-H), 6.00 (d, J=5.94Hz, 1H, Pyr-H), 5.08 (s, 2H, CH₂-Ar), 4.97 (bs, 1H, NH), 4.21 (brs, 2H, CHH), 3.24 (brs, 2H, CH₂-N), 2.75 (brs, 2H, CHH), 2.48 (s, 3H, CH₃), 2.35 (s, 3H, CH₃), 1.82-1.65 (m, 3H, CHH, CH), 1.27-1.12 (m, 2H, CHH).

M.S. (M+1): 387.34.

EXAMPLE 20:

Benzyl 4-(((6-chloro-4-pyrimidinyl)amino]methyl)-1-piperidinecarboxylate



The title compound was prepared as described in **EXAMPLE 13**,


except using 4,6-dichloropyrimidine (1.26g, 8.45mmol) in place of 6-chloropurine as starting materials and adding TEA (2.80mL, 20.13mmol) in 10mL DMF. The procedure gave the title compound as a yellow oil.

¹H NMR (400MHz, CDCl₃): δ 8.32 (s, 1H, Pyr-H), 7.37-7.28 (m, 5H, Ar-H), 6.35 (s, 1H, Pyr-H), 5.72 (s, 1H, NH), 5.13 (s, 2H, CH₂-Ar), 4.22 (brs, 2H, CHH), 3.23 (brs, 2H, CH₂-N), 2.78 (brs, 2H, CHH), 1.85-1.66 (m, 3H, CHH, CH), 1.27-1.10 (m, 2H, CHH).

M.S. (M+1): 361.32.

EXAMPLE 21:

Benzyl 4-(((2-amino-9H-purin-6-yl)amino]methyl)-1-piperidinecarboxylate



The title compound was prepared as described in **EXAMPLE 13**,


except using benzyl 4-(aminomethyl)-1-piperidinecarboxylate (300mg, 1.21mmol) and 4-amino-6-chloropurine (68mg, 0.40mmol) as starting material. The procedure gave the title compound as a yellow oil.

¹H NMR (400MHz, CDCl₃): δ 7.60 (s, 1H, purine-H), 7.38-7.28 (m, 5H, Ar-H), 6.01 (vbs, 1H, NH), 5.12 (s, 2H, CH₂-Ar), 4.86 (vbs, 2H, NH₂), 4.19 (brs, 2H, CHH), 3.48 (brs, 2H, CH₂-N), 2.77 (brs, 2H, CHH), 1.88-1.70 (m, 3H, CHH, CH), 1.30-1.13 (m, 2H, CHH).

M.S. (M+1): 382.31.

EXAMPLE 22:

Benzyl 4-[[[(6-chloro-3-pyridazinyl)amino]methyl]-1-piperidinecarboxylate




The title compound was prepared as described in **EXAMPLE 13**, except using benzyl 4-(aminomethyl)-1-piperidinecarboxylate (1.08g, 4.34mmol), 3,6-dichloropyridazine (636mg, 4.34mmol) as starting materials which gave the title compound as a yellow oil.

- 10 ¹H NMR (400MHz, CDCl₃): δ 7.38-7.28 (m, 6H, Pyr-H, Ar-H), 7.15 (d, J=9.24Hz, 1H, Pyr-H), 5.12 (s, 2H, CH₂-Ar), 4.89 (bs, 1H, NH), 4.22 (bs, 2H, CHH), 3.32 (bs, 2H, CH₂-N), 2.78 (bs, 2H, CHH), 1.96-1.82 (m, 1H, CH), 1.77 (brd, J=12.34Hz, 2H, CHH), 1.27-1.12 (m, 2H, CHH).
M.S. (M+1): 361.27.

15

EXAMPLE 23:

Benzyl 4-[(3-pyridazinylamino)methyl]-1-piperidinecarboxylate



- 20 Benzyl 4-[[[(6-chloro-3-pyridazinyl)amino]methyl]-1-piperidinecarboxylate (**EXAMPLE 22**) (400mg, 1.11mmol) was dissolved in absolute ethanol. Raney nickel (65mg, 1.11mmol) was then added and the resulting reaction was stirred under 1atm hydrogen for 18h. The catalyst was filtered and the filtrate was concentrated under reduced pressure. The resulting clear oil was purified by silica gel chromatography to give the title compound as a clear oil.


- 25 ¹H NMR (400MHz, CDCl₃): δ 8.54 (dd, J=4.48Hz, 1H, Pyr-H), 7.38-7.29 (m, 5H, Ar-H), 7.14 (dd, J=9.05Hz, 4.48Hz, 1H, Pyr-H), 6.61 (dd,

J=8.96Hz, 1.28Hz, 1H, Pyr-H), 5.12 (s, 2H, CH₂-Ar), 4.83 (bs, 1H, NH), 4.22 (bs, 2H, CHH), 3.33 (bs, 2H, CH₂-N), 2.78 (bs, 2H, CHH), 1.96-1.71 (m, 3H, CHH, CH), 1.27-1.12 (m, 2H, CHH).
M.S. (M+1): 327.25.

5

EXAMPLE 24:

Benzyl 4-[[[(6-hydroxy-3-pyridazinyl)amino]methyl]-1-piperidinecarboxylate




- 10 Benzyl 4-[[[(6-chloro-3-pyridazinyl)amino]methyl]-1-piperidinecarboxylate (**EXAMPLE 22**) (37mg, 0.10mmol) was dissolved in acetic acid (5mL) with sodium acetate (82mg, 1.00mmol) and was heated to 100°C for 18h. The volatiles were removed under reduced pressure and the residue partitioned between sat. NaHCO₃ and ethyl acetate. The organics were dried over Na₂SO₄, filtered and concentrated under reduced pressure, affording the title compound as a clear oil.

- 15 ¹H NMR (400MHz, CDCl₃): δ 10.78 (bs, 1H, OH), 7.38-7.29 (m, 5H, Ar-H), 6.83 (d, J=10.01Hz, 1H, Pyr-H), 6.78 (d, J=9.77Hz, 1H, Pyr-H), 5.12 (s, 2H, CH₂-Ar), 4.20 (bs, 3H, CHH, NH), 3.11 (bs, 2H, CH₂-N), 2.78 (bs, 2H, CHH), 1.87-1.65 (m, 3H, CHH, CH), 1.23-1.13 (m, 2H, CHH).
M.S. (M+1): 343.34.

20

EXAMPLE 25:

4-(Pyrazin-2-ylaminomethyl)-piperidine-1-carboxylic acid benzyl ester




25

Benzyl 4-formyl-1-piperidinecarboxylate (P. E. Malignes, *Tetrahedron*, 53(22):10983-10992(1997)) (100mg, 0.40mmol) and aminopyrazine (46mg, 0.46mmol) were dissolved in toluene under N₂ and was heated to reflux under Dean Stark conditions for 18h. The volatiles were removed *in vacuo* and the residue taken up in ethanol and treated with solid NaBH₄ (76mg, 2.00mmol) in small portions. The reaction aged at 20°C for 1h then was quenched with 2N HCl. The reaction was basified with 1M NaOH and was extracted with ethyl acetate (2x). The combined organics were dried over Na₂SO₄, filtered and concentrated *in vacuo*. The resulting residue was purified by reverse phase HPLC to give the title compound as a yellow oil.

¹H NMR (400MHz, CD₃OD): δ 8.08 (d, J=1.01Hz, 1H, Pyr-H), 7.95 (dd, J=3.29Hz, 1.37Hz, 1H, Pyr-H), 7.71 (d, J=3.29Hz, 1H, Pyr-H), 7.35-7.28 (m, 5H, Ar-H), 5.10 (s, 2H, CH₂-Ar), 4.18-4.14 (m, 2H, CHH), 3.27 (d, J=2.14Hz, 2H, CH₂-N), 2.83 (brs, 2H, CHH), 1.88-1.65 (m, 3H, CHH, CH), 1.23-1.09 (m, 2H, CHH).
M.S. (M+1): 327.26.

EXAMPLE 26:


Benzyl 4-[(1,3-thiazol-2-ylamino)methyl]-1-piperidinecarboxylate



The title compound was prepared as described in EXAMPLE 25, except using benzyl 4-formyl-1-piperidinecarboxylate (300mg, 1.21mmol) and 2-amino-1,3-thiazole (133mg, 1.33mmol) as starting materials to give the title compound as a yellow oil.


¹H NMR (400MHz, CDCl₃): δ 7.38-7.28 (m, 5H, Ar-H), 7.07 (d, J=3.66Hz, 1H, thiazole-H), 6.43 (d, J=3.66Hz, 1H, thiazole-H), 6.39 (brs, 1H, NH), 5.12 (s, 2H, CH₂-Ar), 4.20 (brs, 2H, CHH), 3.15 (d, J=6.58Hz, 2H, CH₂-N), 2.77 (brs, 2H, CHH), 1.89-1.71 (m, 3H, CHH, CH), 1.26-1.10 (m, 2H, CHH).
M.S. (M+1): 332.34.

EXAMPLE 27:



Step 1:


Benzyl 4-[(3-methyl-2-pyridinyl)amino]methyl-1-piperidinecarboxylate



The benzyl 4-[(3-methyl-2-pyridinyl)amino]methyl-1-piperidinecarboxylate was prepared as described in EXAMPLE 1, except that 1-[(benzyloxy)carbonyl]-4-piperidinecarboxylic acid (5.00g, 18.99mmol), 2-amino-3-methylpyridine (2.16g, 19.94mmol), EDC (4.37g, 22.79mmol), and HOAt (2.71g, 19.94mmol) and DMF (3mL) were used as starting materials. Benzyl 4-[(3-methyl-2-pyridinyl)amino]methyl-1-piperidinecarboxylate was isolated as an off-white solid and used without further purification.

Step 2:


Piperidine-4-carboxylic acid (3-methyl-pyridin-2-yl)-amide



Benzyl 4-[(3-methyl-2-pyridinyl)amino]methyl-1-piperidinecarboxylate from Step 1 above (5.45g, 15.42mmol) was suspended in absolute ethanol (250mL) and was treated with 10% palladium on carbon (1.50g) and stirred vigorously for 18h under 1atm of hydrogen. The catalyst was filtered off and the filtrate was concentrated under reduced pressure giving the piperidine-4-carboxylic acid (3-methyl-pyridin-2-yl)-amide as yellow oil.

Step 3:


4-(3-Methyl-pyridin-2-yl-carbamoyl)-piperidine-1-carboxylic acid 4-methyl-benzyl ester



Piperidine-4-carboxylic acid (3-methyl-pyridin-2-yl)-amide from Step 2 above (100mg, 0.46mmol) and N-[4-(methylbenzyl-oxy)-carbonyloxy]succinimide (127mg, 0.48mmol) were combined in DMF at r.t. and were stirred vigorously for 15min. The resulting reaction mixture was then purified by reverse phase preparatory HPLC to give 4-(3-methyl-pyridin-2-yl-carbamoyl)-piperidine-1-carboxylic acid 4-methyl-benzyl ester as a clear oil.

Step 4:

4-[(3-Methyl-pyridin-2-ylamino)-methyl]-piperidine-1-carboxylic acid 4-methyl-benzyl ester



4-(3-Methyl-pyridin-2-yl-carbamoyl)-piperidine-1-carboxylic acid 4-methyl-benzyl ester from Step 3 above (65mg, 0.18mmol) was treated with 1M $\text{BF}_3 \cdot \text{THF}$ (1.80mmol, 1.80mL, 1M in THF) over 10min. and was aged at r.t. 4h. The reaction was quenched by slowly adding 2N HCl and stirring vigorously for 30 min. The reaction was basified with sat. NaHCO_3 and extracted with ethyl acetate (2x). The combined organics were washed with brine, dried over Na_2SO_4 , filtered and concentrated *in vacuo*, yielding a white foam which was purified by silica gel chromatography (99:10.1 to 95:5.0.5 $\text{CH}_2\text{Cl}_2:\text{CH}_3\text{OH}:\text{NH}_4\text{OH}$) to give **EXAMPLE 27** as a yellow oil.


^1H NMR (400MHz, CD_3OD): δ 8.00 (d, J=2.47Hz, 1H, Pyr-H), 7.26-7.15 (m, 6H, Pyr-H Ar-H), 6.88 (dd, J=7.03Hz, 5.12Hz, 1H, Pyr-H), 5.08 (s, 2H, $\text{CH}_2\text{-Ar}$), 4.18 (brs, 2H, CHH), 3.39 (brs, 2H, $\text{CH}_2\text{-N}$), 2.78 (brs, 2H, CHH), 2.35 (s,

3H, CH_3), 2.07 (s, 3H, CH_3), 1.90-1.60 (m, 3H, CHH, CH), 1.30-1.10 (m, 4.16Hz, 2H, CHH).

M.S. (M+1): 354.41.

EXAMPLE 28:

4-Fluorobenzyl 4-[(3-methyl-2-pyridinyl)amino]methyl]-1-




The piperidine compound (600mg, 2.74mmol) from **EXAMPLE 27**, Step 2, was treated in accordance with Steps 3 and 4 of that **EXAMPLE 27**, except that N-[4-(fluorobenzyl-oxy)-carbonyloxy]succinimide (805mg, 3.01mmol) was used instead of N-[4-(methylbenzyl-oxy)-carbonyloxy]succinimide in Step 3 to give 4-fluorobenzyl 4-[(3-methyl-2-pyridinyl)amino]methyl]-1-piperidinecarboxylate as a clear oil.

^1H NMR (400MHz, CDCl_3): δ 7.99 (d, J=4.29Hz, 1H, Pyr-H), 7.34-7.31 (m, 2H, Ar-H), 7.20-7.18 (m, 1H, Pyr-H), 7.05-7.00 (m, 1H, Pyr-H), 6.50 (dd, J=7.13Hz, 5.12Hz, 2H, Ar-H), 5.08 (s, 2H, $\text{CH}_2\text{-Ar}$), 4.22 (brs, 3H, CHH, NH), 3.38 (brs, 2H, $\text{CH}_2\text{-N}$), 2.77 (brs, 2H, CHH), 2.06 (s, 3H, CH_3), 1.84-1.77 (m, 3H, CHH, CH), 1.26-1.12 (m, 2H CHH).

M.S. (M+1): 358.35.

EXAMPLE 29:

4-Chlorobenzyl 4-[(3-methyl-2-pyridinyl)amino]methyl]-1-



The piperidine compound (600mg, 2.74mmol) from **EXAMPLE 27**, Step 2, was treated in accordance with Steps 3 and 4, except that N-[4-(chlorobenzyl-oxy)carbonyloxy]succinimide (855mg, 3.01mmol) was used instead of N-[4-

(methylbenzoyloxy)succinimide in Step 3 to give 4-chlorobenzyl 4-[(3-methyl-2-pyridinylamino)methyl]-1-piperidinecarboxylate as a clear oil.

¹H NMR (400MHz, CDCl₃): δ 7.99 (dd, J=4.90Hz, 1.23Hz, 1H, Pyr-H), 7.32-7.27 (m, 4H, Ar-H), 7.20-7.18 (m, 1H, Pyr-H), 6.50 (dd, J=7.18Hz, 5.08Hz, 1H, Pyr-H), 5.08 (s, 2H, CH₂-Ar), 4.20 (brs, 3H, CHH, NH), 3.38 (brs, 2H, CH₂-N), 2.78 (brs, 2H, CHH), 2.06 (s, 3H, CH₃), 1.90-1.72 (m, 3H, CHH, CH), 1.26-1.12 (m, 2H, CHH).


M.S. (M+1): 374.31.

10 EXAMPLE 30:

3-Fluorobenzyl 4-[(4-pyridinylamino)methyl]-1-piperidinecarboxylate

Step 1:

N-(4-piperidinylmethyl)-4-pyridinamine




15

Benzyl 4-[(4-pyridinylamino)methyl]-1-piperidinecarboxylate (EXAMPLE 1) (7g, 21mmol) was dissolved in abs. Ethanol (150mL) with 10% palladium on carbon (700mg) and stirred under 1atm of hydrogen for 2h. The catalyst was filtered off and the filtrate was concentrated under reduced pressure to afford the N-(4-piperidinylmethyl)-4-pyridinamine as a clear oil which was used without further purification.

Step 2:

3-Fluorobenzyl 4-[(4-pyridinylamino)methyl]-1-piperidinecarboxylate




3-Fluorobenzyl alcohol (30mg, 0.24mmol) was treated with triphosgene (24mg, 0.08mmol) and N-(4-piperidinylmethyl)-4-pyridinamine (50mg, 0.26mmol), and aged at 40°C for 45min. The resulting reaction solution was partitioned between 0.5M NaOH and ethyl acetate. The organics were separated, dried over Na₂SO₄, filtered and concentrated under reduced pressure. The resulting

oil was purified by preparatory HPLC to give the TFA salt of EXAMPLE 30 as a yellow oil. M.S. (M+1): 344.36.

The following EXAMPLES 32-36 were prepared as described above in EXAMPLE 30, but replacing 3-fluorobenzyl alcohol with the appropriate alcohol:

EXAMPLE 31:

2-Methylbenzyl 4-[(4-pyridinylamino)methyl]-1-piperidinecarboxylate




10

M.S. (M+1): 340.38.

EXAMPLE 32:

3-Methylbenzyl 4-[(4-pyridinylamino)methyl]-1-piperidinecarboxylate




15

M.S. (M+1): 340.39.

EXAMPLE 33:

4-Methylbenzyl 4-[(4-pyridinylamino)methyl]-1-piperidinecarboxylate




20

M.S. (M+1): 340.29.

EXAMPLE 34:

2-Methoxybenzyl 4-[(4-pyridinylamino)methyl]-1-piperidinecarboxylate




M.S. (M+1): 356.37.

5

EXAMPLE 35:

3-Methoxybenzyl 4-[(4-pyridinylamino)methyl]-1-piperidinecarboxylate




10

M.S. (M+1): 356.37.

EXAMPLE 36:

4-Methoxybenzyl 4-[(4-pyridinylamino)methyl]-1-piperidinecarboxylate



15

M.S. (M+1): 356.36.

EXAMPLE 37:

4-Fluorobenzyl 4-[(2-pyrimidinylamino)methyl]-1-piperidinecarboxylate

Benzyl 4-[(2-pyrimidinylamino)methyl]-1-piperidinecarboxylate (EXAMPLE 16) was hydrogenated as described in EXAMPLE 30, Step 1.

Treatment with N-[4-(fluorobenzoyloxy)-carbonyloxy]succinimide as described in


5 EXAMPLE 27, Step 3, afforded the 4-fluorobenzyl 4-[(2-pyrimidinylamino)methyl]-1-piperidinecarboxylate.

¹H NMR (400MHz, CDCl₃): δ 8.26 (d, J=4.89Hz, 2H, Pyr-H), 7.33-7.27 (m, 2H, Ar-H), 7.05-7.01 (m, 2H, Ar-H), 6.53 (t, J=4.76Hz, 1H, Pyr-H), 5.45 (brt, J=5.73Hz, 1H, NH), 5.08 (s, 2H, CH₂-Ar), 4.20 (brd, J=27.6Hz, 2H, CHH), 3.32 (t, J=6.22Hz, 2H, CH₂-N), 2.77 (brs, 2H, CHH), 1.83-1.75 (m, 3H, CHH, CH), 1.26-1.15 (m, 2H CHH).

M.S. (M+1): 345.35.

EXAMPLE 38:

15 4-Chlorobenzyl 4-[(2-pyrimidinylamino)methyl]-1-piperidinecarboxylate



The title compound was prepared as described in EXAMPLE 37, except replacing N-[4-(fluorobenzoyloxy)-carbonyloxy]succinimide with N-[4-(chlorobenzoyloxy)-carbonyloxy] succinimide.

¹H NMR (400MHz, CDCl₃): δ 8.25 (d, J=4.75Hz, 2H, Pyr-H), 7.33-7.27 (m, 4H, Ar-H), 6.51 (t, J=4.84Hz, 1H, Pyr-H), 5.77 (bs, 1H, NH), 5.08 (s, 2H, CH₂-Ar), 4.18 (brs, 2H, CHH), 3.32 (brt, J=6.12Hz, 2H, CH₂-N), 2.77 (brs, 2H, CHH), 1.84-1.75 (m, 3H, CHH, CH), 1.26-1.12 (m, 2H CHH).


M.S. (M+1): 361.32.

25

EXAMPLE 39:

(Cis)-3-hydroxy-4-(pyridin-4-ylaminomethyl)-piperidine-1-carboxylic acid benzyl ester
Step 1:


1-Benzyl-4-hydroxymethyl-piperidin-3-ol



Sodium borohydride (40g) was added in portions to a stirred solution of ethyl N-benzyl-3-oxopiperidine-4-carboxylate hydrochloride (23.6g, 90mmol) in methanol (500mL), over 2h. Water (300mL) was added slowly, the mixture stirred for 15min and then the organics were evaporated. The residue was partitioned between DCM and water (x3), the combined organic layers dried over anhydrous sodium sulfate, and the solvent evaporated to give 1-benzyl-4-hydroxymethyl-piperidin-3-ol product as a cis/trans mixture, which was used in the next step without further purification. M.S (M+1): 222.

Step 2:

3-Hydroxy-4-hydroxymethyl-piperidine-1-carboxylic acid benzyl ester




A solution of the 1-benzyl-4-hydroxymethyl-piperidin-3-ol from Step 1 above (13.5g) in methanol (450mL) was hydrogenated at 50psi over 20% palladium hydroxide on charcoal (10g) for 48h in three batches. The combined reaction mixtures were filtered and the filtrate evaporated to give an oil. This was dissolved in water (100mL) and dioxane (100mL), cooled to 5°C, and benzyl chloroformate (7.8mL) was added slowly, with addition of 1M NaOH to maintain a pH of 10-11. After 30min, the cooling bath was removed and reaction mixture stirred for 30min. The reaction mixture was concentrated to remove dioxane and the residue extracted with EtOAc (x3). The combined extracts were washed with brine, dried over anhydrous sodium sulfate and solvent evaporated to give a mixture of cis and trans 3-hydroxy-4-hydroxymethyl-piperidine-1-carboxylic acid benzyl ester products.

Purification by flash column chromatography (80% EtOAc hexane to 5% MeOH EtOAc) gave the upper Rf cis isomer (major) and the lower Rf trans isomer (minor). M.S (M+1): 266.

Step 3:


(Cis)-3-hydroxy-4-(toluene-4-sulfonyloxymethyl)-piperidine-1-carboxylic acid benzyl ester



A solution of the (cis)-3-hydroxy-4-hydroxymethyl-piperidine-1-carboxylic acid benzyl ester from Step 2 above (7.65g) in chloroform (200mL) was treated with pyridine (2.6mL) and 4-toluenesulfonyl chloride (6.05g) and the reaction mixture heated to 60°C for 18h. Additional pyridine (0.85mL) and 4-toluenesulfonyl chloride (2.0g) were added to the cooled reaction and heating continued for a further 24h. The resulting reaction mixture was cooled to room temperature and washed with 10% aqueous citric acid solution and water, dried over anhydrous sodium sulfate and the solvent evaporated to give, after flash column chromatography, the (cis)-3-hydroxy-4-(toluene-4-sulfonyloxymethyl)-piperidine-1-carboxylic acid benzyl ester.

Step 4:


(Cis)-4-aminomethyl-3-hydroxy-piperidine-1-carboxylic acid benzyl ester



A solution of the tosylate compound (6.80g) from Step 3 above was dissolved in DMF (50mL) and treated with sodium azide (3.16g). The reaction mixture was then heated to 50°C for 48h, cooled to room temperature and partitioned between dilute aqueous sodium bicarbonate and EtOAc. The organic layer was washed with brine, dried over anhydrous sodium sulfate and solvent evaporated to give the azide (5.23g) which was dissolved in THF (50mL) and treated with triphenylphosphine (14.07g) and water (3.25mL). The reaction mixture was stirred

for 18h at room temperature, the volatiles evaporated and the residue purified by flash column chromatography (DCM to 80/20/2 DCM MeOH NH₄OH) to give (cis)-4-aminomethyl-3-hydroxy-piperidine-1-carboxylic acid benzyl ester as an oil.
M.S (M+1): 265.

5 Step 5: (Cis)-3-hydroxy-4-(pyridin-4-ylaminomethyl)-piperidine-1-carboxylic acid benzyl ester



A mixture of the cis 4-aminomethyl-3-hydroxy-piperidine-1-carboxylic acid benzyl ester (245mg) from Step 4 above, 4-chloropyridine (105mg) and isopropanol (0.4mL) was heated to 120°C in a sealed vial for 24h, cooled to room temperature and the solvents evaporated. The resulting crude mixture was purified by flash column chromatography (DCM to 80/20/2 DCM MeOH NH₄OH) to give impure cis 3-hydroxy-4-(pyridin-4-ylaminomethyl)-piperidine-1-carboxylic acid benzyl ester. This was purified by preparative reverse phase HPLC (95% H₂O 5% MeCN to 100% MeCN both containing 0.1% TFA). Evaporation gave an oil which was partitioned between DCM and aqueous sodium bicarbonate solution. The organic layer was dried over anhydrous sodium sulfate and solvent evaporated to give a white solid. M.S (M+1): 342.

EXAMPLE 40:

(-)-(Cis)-3-hydroxy-4-(pyridin-4-ylaminomethyl)-piperidine-1-carboxylic acid benzyl ester and (+)-(cis)-3-hydroxy-4-(pyridin-4-ylaminomethyl)-piperidine-1-carboxylic acid benzyl ester


The enantiomers of (cis)-3-hydroxy-4-(pyridin-4-ylaminomethyl)-piperidine-1-carboxylic acid benzyl ester were separated by preparative HPLC on a Chiralpak® AD column, eluting with 70% (0.1% diethylamine in hexane) 30% isopropanol to give the earlier eluting (-) enantiomer followed by the (+)-enantiomer.

EXAMPLE 41:

(cis)-3-hydroxy-4-(pyridin-4-ylaminomethyl)-piperidine-1-carboxylic acid 4-methyl-benzyl ester

Step 1:


5 3-Hydroxy-4-[(2,3,5,6-tetrachloro-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



2,3,5,6-tetrachloro-4-nitropyridine (S. M. Roberts et al., *J. Chem. Soc. C*, 2844-2848(1968)) (1.7g, 6.5mmol) was added to a solution of (cis)-4-aminomethyl-3-hydroxy-piperidine-1-carboxylic acid benzyl ester (1.71g, 6.49mmol) and N-methylmorpholine (0.785mL, 7.15mmol) in THF (50mL) at room temperature. The resulting reaction mixture was stirred for 18h at room temperature then partitioned between EtOAc and water. The organic layer was washed with saturated sodium bicarbonate solution, dried over anhydrous sodium sulfate and the solvent evaporated to give crude product purified by flash column chromatography (20-80% EtOAc hexane) to give 3-hydroxy-4-[(2,3,5,6-tetrachloro-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester compound. M.S (M+1): 478.

Step 2:


4-(Pyridin-4-ylaminomethyl)-piperidin-3-ol



20 A suspension of 3-hydroxy-4-[(2,3,5,6-tetrachloro-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester compound from Step 1 above (1.64g) and potassium carbonate (6g) in ethanol (200mL) was hydrogenated at 60psi over 1g of 10% palladium on charcoal for 5h. The reaction mixture was filtered and the solids washed well with ethanol. The filtrate was evaporated, taken up in 40% MeOH DCM and refiltered. The filtrate was evaporated to give crude 4-(pyridin-4-ylaminomethyl)-piperidin-3-ol product used without further purification in the next step.

Step 3:

(Cis)-3-hydroxy-4-(pyridin-4-ylaminomethyl)-piperidine-1-carboxylic acid 4-methyl-benzyl ester




5 A suspension of the 4-(pyridin-4-ylaminomethyl)-piperidin-3-ol from Step 2 above (0.076g, 0.367mmol) in DMF (1.5mL) was treated with carbonic acid 2,5-dioxo-pyrrolidin-1-yl ester 4-methyl-benzyl ester (0.097g, 0.37mmol) (INTERMEDIATE 1A) and the resulting reaction mixture stirred at rt for 5min. The mixture was then partitioned between dilute sodium carbonate solution and EtOAc. The organic layer was washed with saturated sodium bicarbonate solution and brine, dried over anhydrous sodium sulfate, and the solvent evaporated to give a crude product. Purification by flash column chromatography (DCM to 80/20/2 DCM MeOH NH₄OH) afforded the cis 3-hydroxy-4-(pyridin-4-ylaminomethyl)-piperidine-1-carboxylic acid 4-methyl-benzyl ester compound. M.S (M+1): 356.

15

EXAMPLE 42:


(Cis)-3-hydroxy-4-(pyridin-4-ylaminomethyl)-piperidine-1-carboxylic acid 4-ethyl-benzyl ester



20 The title compound was prepared as described in EXAMPLE 41, Step 3, but replacing carbonic acid 2,5-dioxo-pyrrolidin-1-yl ester 4-methyl-benzyl ester with carbonic acid 2,5-dioxo-pyrrolidin-1-yl ester 4-ethyl-benzyl ester (INTERMEDIATE 1D). M.S (M+1): 370

25 EXAMPLE 43:

(Cis)-3-hydroxy-4-(pyridin-2-ylaminomethyl)-piperidine-1-carboxylic acid benzyl ester




5 A mixture of (cis)-4-aminomethyl-3-hydroxy-piperidine-1-carboxylic acid benzyl ester (0.1g, 378mmol) and 2-fluoropyridine (0.25mL) was heated to 120°C for 24h. The reaction mixture was partitioned between EtOAc and water. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and the solvent evaporated to give a (cis)-3-hydroxy-4-(pyridin-2-ylaminomethyl)-piperidine-1-carboxylic acid benzyl ester crude product, which was purified by flash column chromatography (50% EtOAc hexane to 5% MeOH EtOAc). M.S (M+1): 342

10

EXAMPLE 44:

4-[(3-Cyano-pyridin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



15


A mixture of benzyl 4-(aminomethyl)piperidine-1-carboxylate (EXAMPLE 13, Step 1) (1g, 4.03mmol) and 3-cyanopyridine (0.25g) was heated to 100°C for 30min. The reaction mixture was partitioned between EtOAc and pH5.2 citrate buffer. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and the solvent evaporated to give a solid which was stirred with 5mL ether

20

and 0.5mL EtOAc for 1h and filtered to give the title compound as a solid. M.S (M+1): 351

EXAMPLE 45:

5 4-[(3-Chloro-pyridin-2-ylamino)-methyl]-piperidine-1-carboxylic acid
benzyl ester




A mixture of benzyl 4-(aminomethyl)piperidine-1-carboxylate (EXAMPLE 13, Step 1) (1g, 4.03mmol) and 2,3-dichloropyridine (0.25g) was heated to 100°C for 12h. The reaction mixture was cooled and partitioned between EtOAc and pH5.2 citrate buffer. The organic layer was washed with brine, dried over anhydrous sodium sulfate and the solvent evaporated to give a crude product. Purification by flash column chromatography (5-50% EtOAc hexane) afforded the title compound. M.S (M+1): 360.

15

EXAMPLE 46:

4-[(3-Trifluoromethyl-pyridin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester




20 A mixture of benzyl 4-(aminomethyl)piperidine-1-carboxylate (EXAMPLE 13, Step 1) (1g, 4.03mmol) and 2-chloro-3-trifluoromethylpyridine (0.25g) was heated to 100°C for 12h. The reaction mixture was cooled and partitioned between EtOAc and pH5.2 citrate buffer. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and the solvent evaporated to give a

crude product. Purification by flash column chromatography (5-50% EtOAc hexane) afforded the title compound. M.S (M+1): 394.

EXAMPLE 47:

5 4-[(3-Chloro-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid
benzyl ester



A mixture of benzyl 4-(aminomethyl)piperidine-1-carboxylate (EXAMPLE 13, Step 1) (1.25g, 5.04mmol) and 2,3-dichloropyrazine (0.25g) was heated to 100°C for 1h. The reaction mixture was cooled and partitioned between EtOAc and pH5.2 citrate buffer. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and the solvent evaporated to give a crude product. Purification by flash column chromatography (5-50% EtOAc hexane) afforded the title compound. M.S (M+1): 361.


15

EXAMPLE 48:

4-[(3-Hydroxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester

Step 1:


20 3-[(Piperidin-4-ylmethyl)-amino]-pyrazin-2-ol



4-[(3-Chloro-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester (EXAMPLE 47) (2.21g, 6.12mmol) and 3M HCl (200mL) was heated to reflux for 18h, cooled to rt and the volatiles evaporated. The residue was azeotroped with ethanol (3x100mL) and then stirred with 50mL ether for 1h, filtered and the solid dried to yield a cream solid.

Step 2:


4-[(3-Hydroxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



- 5 To a solution of 3-[(piperidin-4-ylmethyl)-amino]-pyrazin-2-ol from Step 1 above (0.287g, 1.021mmol) in DMF (5mL) was added triethylamine (0.356mL, 2.55mmol), followed by N-(benzyloxycarbonyloxy)succinimide (0.305g, 1.23mmol). The resulting reaction mixture was stirred at rt for 15min, then partitioned between EtOAc and water. The organic layer was washed with water and brine, dried over anhydrous sodium sulfate and the crude product purified by flash column chromatography (50% EtOAc hexane to 5% MeOH EtOAc) to give an oil which solidified on standing. M.S (M+): 343.24.


EXAMPLE 49:

- 15 4-[(5-Chloro-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester
- Step 1:
- 4-[(2,5,6-Trichloro-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



- 20 To a solution of benzyl 4-(aminomethyl)piperidine-1-carboxylate (EXAMPLE 13, Step 1) and N,N-diisopropylethylamine (2.6g, 20mmol) in THF (40mL) at -78°C was added a solution of tetrachloropyrimidine (4.4g, 20mmol). The cooling bath was removed and the solution was stirred for 45 min. The solution was concentrated and purified by filtering through a pad of silica gel using ether.
- Step 2:


4-[(5-Chloro-2,6-bis-methylsulfonyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



- 5 To 4-[(2,5,6-trichloro-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester (1g, 2.33mmol) in DMF was added sodium thiomethoxide (0.4g, 5.8mmol). The resulting reaction mixture was stirred for 2h and quenched with aqueous ammonium chloride. The product was extracted with ethyl acetate, dried (Na₂SO₄), concentrated, and purified by silica gel chromatography (ether / hexanes).

Step 3:

10 4-[(5-Chloro-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



- 15 4-[(5-Chloro-2,6-bis-methylsulfonyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester (1.0g, 2.2mmol) was suspended in ethanol (15mL) and ethyl acetate added to give a homogeneous solution, and excess Raney Nickel was added. The resulting reaction mixture was stirred overnight. More Raney Nickel was added and the reaction mixture was heated to 80°C for 3h. The mixture was filtered and the solids were washed with hot ethanol/ethyl acetate several times. The organics were concentrated and the resulting residue was purified by silica gel chromatography (isopropanol/methylene chloride). The product was dissolved in ether and treated with ethereal HCl (2.2mmol) to form the HCl salt which was collected by filtration. The resulting 4-[(5-chloro-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester hydrochloride salt was collected by filtration as a colorless solid.
- 25

¹H NMR (400MHz, CD₃OD): δ 8.67 (s, 1h, pyrimidine), 8.45 (s, 1h, pyrimidine), 7.32 (m, 5h, Ar), 5.10 (s, 2h, CHH), 4.15 (d, J = 13.0 Hz, 2h, CHH), 3.58 (d, J = 7.2 Hz, 2h, CHH), 2.83 (m, 2h, CHH), 1.97 (m, 1h, CH), 1.74 (d, J = 12.0 Hz, 2h, CHH).


M.S (M+1): 361.3

EXAMPLE 50:

4-[(2-Hydroxymethyl)-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester

Step 1:


Benzyl 4-(aminomethyl)piperidine-1-carboxylate



4-Aminomethylpiperidine (40g, 350mmol) and benzaldehyde (37.3mL, 368mmol) in toluene (600mL) were heated to reflux under dean stark conditions for 2h. The reaction mixture was cooled to room temperature and 500mL dichloromethane added. The solution was cooled to 5°C and treated with N-(benzyloxy)carbonyloxysuccinimide (91.7g, 368mmol). After 10min., the cooling bath was removed and the reaction mixture stirred for 1h. The solvents were evaporated and the residue stirred with 400mL THF and 400mL 2M HCl for 1h. The mixture was concentrated to remove organics and extracted with ether (3x300mL). The aqueous phase was adjusted to pH14 with 50% NaOH and extracted with ethyl acetate. The organic layer was washed with water and brine, dried over anhydrous sodium sulfate, and the solvent evaporated to give benzyl 4-(aminomethyl)piperidine-1-carboxylate compound.

Step 2:

4-[(1-Benzyloxycarbonyl)-piperidin-4-ylmethyl]-amino)-pyridine-2-carboxylic acid



To a solution of 4-chloropicolinic acid (0.8gm, 0.0051mol) in DMSO (4mL) was added benzyl 4-(aminomethyl)piperidine-1-carboxylate (2.5gm, 0.010mol) and the mixture warmed to 140°C for 18h. The reaction was cooled and diluted with 10 % sodium bicarbonate (100mL) and washed with ether (2x 25mL).


The aqueous extract was washed with dichloromethane (3x 50mL), and the dichloromethane extract dried over sodium sulfate and concentrated to an oil (2.4gm). The oil was chromatographed on silica using dichloromethane/methanol/acetate acid/water-90/10/1/1 to give 4-[(1-benzyloxycarbonyl)-piperidin-4-ylmethyl]-amino)-pyridine-2-carboxylic acid.

¹H NMR 400MHz (δ, DMSO) δ: 8.98 (s, 1H), 8.2-8.0 (m, 1H); 7.6-7.2 (m, 5H); 7.01(brs, 1H); 5.08(s, 2H); 4.02 (brd, 2H); 2.80 (brs, 2H); 1.8-1.6 (m, 3H); 1.3-1.1 (m, 2H).

M.S (M+1): 370.

Step 3:

4-[(2-Hydroxymethyl)-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



To a 0°C solution of 4-[(1-benzyloxycarbonyl)-piperidin-4-ylmethyl]-amino)-pyridine-2-carboxylic acid (0.59gm, 0.0016mol) in THF (2mL) under nitrogen was added a solution of 1.0M borane-tetrahydrofuran (6mL) and the mixture allowed to stir at room temperature for 1h. The reaction was cooled to 0°C, quenched with 1N HCl (10mL), concentrated and diluted with 10% aqueous sodium bicarbonate.

Extraction with dichloromethane (2x 50mL) and concentration of the organic layer gave 540mg of crude material. Column chromatography using

dichloromethane/methanol/ammonium hydroxide-90/10/2 and crystallization from diethyl ether gave 4-[(2-hydroxymethyl)-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester.

¹H NMR (400MHz CDCl₃) δ: 8.13 (d, 1H, J=6.8Hz); 7.5-7.1 (m, 5H); 6.35 (m, 2H); 5.12(s, 2H); 4.61 (s, 2H); 4.20 (brm, 3H); 3.08 (m, 2H); 2.78(m, 2H) 1.8-1.6 (m, 3H); 1.3-1.1 (m, 2H).


M.S (M+1): 356.

EXAMPLE 51:

4-[(2-Dimethylaminomethyl)-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester

5 Step 1:

4-[(2-Dimethylcarbamoyl)-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester




To a mixture of 4-[(1-benzylloxycarbonyl)-piperidin-4-ylmethyl]-1-amino-pyridine-2-carboxylic acid (**EXAMPLE 50**, Step 2) (50mg, 0.000135mol), 1-hydroxybenzotriazole hydrate (31mg, 0.0002mol), 2.0M dimethylamine/THF (0.100mL, 0.0002mol) and triethylamine (0.048mL, 0.0002mol) in DMF (2mL) was added 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (39mg, 0.0002mol) and the mixture allowed to stir at room temperature for 7 days. The mixture was quenched into water (10mL) and extracted with ethyl acetate (20mL). The ethyl acetate extract was washed with 10% aqueous sodium bicarbonate (10mL), brine (5mL), dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue chromatographed (reverse phase C-18 using acetonitrile/0.1% trifluoroacetic acid in water) to give 4-[(2-dimethylcarbamoyl)-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester as its trifluoroacetate salt.

M.S.(M+1): 397.

Step 2:

4-[(2-dimethylaminomethyl)-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



25

To 4-[(2-Dimethylcarbamoyl)-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester (28mg, 0.05mmol) was added a solution of 1.0M

borane-tetrahydrofuran (2mL). The reaction was stirred at room temperature for 24h. The reaction was quenched with 1N HCl (2mL) and concentrated *in vacuo* to an oil. Reverse phase chromatography (C-18 using acetonitrile/0.1% trifluoroacetic acid in water) gave upon concentration *in vacuo* **EXAMPLE 51**.


5

¹H NMR (400MHz CD₃OD) δ: 8.10 (m, 1H); 7.4-7.2 (m, 5H); 7.2-6.8 (m, 2H); 5.12(s, 2H); 4.41 (s, 2H); 4.18 (m, 2H); 3.30(m, 2H); 2.78(m, 2H) 1.8-1.6 (m, 3H); 1.3-1.1 (m, 2H).

M.S.(M+1): 383.

10 **EXAMPLE 52:**

4-[(2-Methylaminomethyl)-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester




The title compound was prepared in a similar manner to **EXAMPLE 51**, except replacing dimethylamine with methylamine in Step 1. M.S.(M+1): 369.

15

EXAMPLE 53:

4-[(3-Chloro-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid 4-fluoro-benzyl ester



20

To 2,3-dichloropyrazine (0.160gm, 0.00107mol) was added 4-fluorobenzyl 4-(aminomethyl)piperidine-1-carboxylate (**INTERMEDIATE 2C**) (0.86gm, 0.00322mol) and the resulting mixture heated under nitrogen at 110°C for 30min. The reaction was cooled, diluted with ethyl acetate (50mL), and washed with 10% aqueous sodium/citric acid pH=5.2 (3X 30mL), and 10% aqueous sodium bicarbonate (30mL). The ethyl acetate extract was dried over sodium sulfate, filtered

25

through a pad of silica and concentrated to an oil. Crystallization from ether/hexane gave 4-(3-chloro-pyrazin-2-ylamino)-methyl-piperidine-1-carboxylic acid 4-fluorobenzyl ester.

¹H NMR (400MHz DMSO d₆) δ: 7.99 (d, 1H, J=2.7 Hz); 7.52 (d, 1H, J=2.7 Hz); 7.41 (d, 1H, J=5.7 Hz); 7.39 (d, 1H, J=5.7 Hz); 7.19 (m, 2H); 7.16 (m, 1H); 5.03 (s, 2H); 3.97 (m, 2H); 3.25 (m, 2H); 2.75 (m, 2H); 1.9 (m, 1H); 1.7 (m, 2H); 1.1-0.9 (m, 2H).


M.S.(M+1): 379.

EXAMPLE 54:

4-Hydroxy-4-(pyridin-4-ylaminomethyl)-piperidine-1-carboxylic acid benzyl ester * TFA salt

Step 1:

4-Aminomethyl-1-benzyl-piperidin-4-ol




15

A mixture of 1-benzyl-4-hydroxy-piperidine-4-carbonitrile (5.00g, 19.78mmol) and BH₃.THF (59.35mmol, 59.35mL, 1M in THF) was heated at 80°C for 1h. Cooled to 0°C and quenched with conc. HCl (20mL), the reaction solution was then stirred at rt in 1h. The reaction solution was basified with 10N NaOH to pH8, and extracted with ethyl acetate (3 x 100mL). The combined extracts were washed with water (50mL), brine (30mL), dried over Na₂SO₄, filtered and concentrated *in vacuo* to give 4-aminomethyl-1-benzyl-piperidin-4-ol.

M.S.(M+1):221.31

Step 2:

4-BOC-aminomethyl-1-benzyl-piperidin-4-ol



25


To a cooled (0°C), stirred solution of 4-aminomethyl-1-benzyl-piperidin-4-ol (4.00g, 18.16mmol) in dry CH₂Cl₂ (40mL), under N₂ was slowly added BOC₂O (4.36g, 19.97mmol) dissolved in dry CH₂Cl₂ (5mL). The ice bath was

removed and the reaction solution allowed to warm to rt over 1h, then concentrated *in vacuo*. The residue was purified by silica gel chromatography, 1 - 10 (10% NH₄OH in MeOH) / 99 - 90 CH₂Cl₂ to give 4-BOC-aminomethyl-1-benzyl-piperidin-4-ol.

M.S.(M+1):321.41

5 Step 3:


4-BOC-aminomethyl-piperidin-4-ol



A mixture of 4-BOC-aminomethyl-1-benzyl-piperidin-4-ol (0.50g, 1.56mmol), Pd(OH)₂ (20% on carbon, 0.05g) in absolute ethanol (15mL) was shaken under 60psi H₂ atmosphere for 3h. Filtered and concentrated, the reaction gave 4-BOC-aminomethyl-piperidin-4-ol. M.S.(M+1):231.28

Step 4:

4-BOC-aminomethyl-1-CBZ-piperidin-4-ol




15

To a cooled (0°C), stirred solution of 4-BOC-aminomethyl-piperidin-4-ol (0.35g, 1.52mmol) in dried CH₂Cl₂ (5mL), under N₂ was slowly added CBZ-Cl (0.24mL, 1.67mmol), followed by triethylamine (0.42mL, 3.04mmol). The ice bath was removed and the reaction solution was stirred to rt in 1h, then concentrated *in vacuo*. The residue was purified by silica gel chromatography (10 CH₂Cl₂ : 1 - 20 IPA : 89 - 10 hexane) to give 4-BOC-aminomethyl-1-CBZ-piperidin-4-ol.

M.S.(M+1):365.39

Step 5:


4-aminomethyl-1-CBZ-piperidin-4-ol



To a stirred solution of 4-BOC-aminomethyl-1-CBZ-piperidin-4-ol (0.50g, 1.37mmol) in dried CH_2Cl_2 (3mL) was slowly added trifluoroacetic acid (3mL). The resulting reaction solution was stirred at rt for 20min., then concentrated *in vacuo*. The residue was dissolved in ethyl acetate (100mL), washed with sat. aq. NaHCO_3 (20mL), water (20mL), brine (10mL), dried over Na_2SO_4 , filtered and concentrated to give 4-aminomethyl-1-CBZ-piperidin-4-ol. M.S.(M+1):265.32

Step 6:

4-Hydroxy-4-(pyridin-4-ylaminomethyl)-piperidine-1-carboxylic acid
benzyl ester * TFA salt




10

A solution of 4-aminomethyl-1-CBZ-piperidin-4-ol (0.10g, 0.38mmol), 4-bromo-pyridine (0.06g, 0.38mmol) in IPA (2mL) was heated at 100°C in a sealed reaction tube for 7h. Cooled to rt, the reaction mixture was diluted with ethyl acetate (100mL), washed with sat. aq. NaHCO_3 (20mL), water (20mL), brine (10mL), dried over Na_2SO_4 , filtered and concentrated. The residue was purified by reverse phase chromatography to give 4-hydroxy-4-(pyridin-4-ylaminomethyl)-piperidine-1-carboxylic acid benzyl ester as a TFA salt. M.S.(M+1):342.35

EXAMPLE 55:

4-[(3-Bromo-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid
benzyl ester




A mixture of benzyl-4-(aminomethyl)piperidine-1-carboxylate

(EXAMPLE 13, Step 1, 0.20g, 0.81mmol), 3,4-dibromo-pyridine (*Chem. Abstracts*, 58:5627) (0.19g, 0.81mmol) in IPA (0.5mL) was heated at 100°C in a sealed reaction tube for 7h, then concentrated *in vacuo*. The residue was purified by silica gel

chromatography (DCM IPA hexane)) to give 4-[(3-Bromo-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester. M.S.(M+1):405.27

EXAMPLE 56:

4-[(3-Fluoro-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid
benzyl ester TFA salt




A mixture of benzyl-4-(aminomethyl)piperidine-1-carboxylate

(EXAMPLE 13, Step 1, 0.20g, 0.81mmol), 3-fluoro-4-iodo-pyridine (*Tetrahedron*, 49:49-64(1993) (0.18g, 0.81mmol) in IPA (0.1mL) was heated at 100°C in a sealed reaction tube for 100h, then concentrated *in vacuo*. The residue was purified by reversed phase chromatography to give 4-[(3-fluoro-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester as a TFA salt. M.S.(M+1):344.36

EXAMPLE 57:

4-[(2-Chloro-6-methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



To a stirred solution of 2,4-dichloro-6-methyl-pyrimidine (3.61g, 22.15mmol), triethylamine (7.02mL, 50.34mmol) in DMF (15mL) was slowly added benzyl-4-(aminomethyl)piperidine-1-carboxylate (EXAMPLE 13, Step 1, 5.00g, 20.13mmol). The resulting reaction solution was stirred at rt for 2h, then diluted with ethyl acetate (400mL), washed with water (3 x 30mL), brine (30mL), dried over Na_2SO_4 , filtered and concentrated. The residue was purified by silica gel chromatography (20 - 80% ethyl acetate in hexane) to give 4-[(2-chloro-6-methyl-

pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester.


M.S. (M+1):375.36

EXAMPLE 58:

4-[(6-Methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester

Step 1:


4-[(6-Methyl-pyrimidin-4-ylamino)-methyl]-piperidine



To a stirred solution of 4-[(2-chloro-6-methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester (EXAMPLE 57, 0.50g, 1.33mmol), Pd/C (10%, 0.05g) in absolute ethanol (15mL) was vigorously stirred under 1 atm H₂ for 6h. Filtered and concentrated, the reaction gave 4-[(6-methyl-pyrimidin-4-ylamino)-methyl]-piperidine. M.S. (M+1):207.30

Step 2:

4-[(6-Methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester




To a stirred solution of 4-[(6-methyl-pyrimidin-4-ylamino)-methyl]-piperidine (0.15g, 0.73mmol), in DMF (1mL) was added carbonic acid benzyl ester 2,5-dioxo-pyrrolidin-1-yl ester (0.18g, 0.73mmol). The resulting reaction solution was stirred at rt for 0.5h, then concentrated *in vacuo*. The residue was purified by silica gel chromatography (90:10:1 DCM MeOH NH₄OH) to give 4-[(6-methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester.

M.S. (M+1):341.37

EXAMPLE 59:

4-[(2-Chloro-5-methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester




To a stirred solution of 2,4-dichloro-5-methyl-pyrimidine (3.61g, 22.15mmol), triethylamine (7.02mL, 50.34mmol) in DMF (15mL) was slowly added benzyl 4-(aminomethyl)piperidine-1-carboxylate (EXAMPLE 13, Step 1, 5.00g, 20.13mmol). The resulting reaction solution was stirred at rt for 2h, then diluted with ethyl acetate (400mL), washed with water (3 x 30mL), brine (30mL), dried over Na₂SO₄, filtered and concentrated. The residue was purified by silica gel chromatography (20 – 80% ethyl acetate in hexane) to give 4-[(2-Chloro-5-methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester. M.S. (M+1):375.36

EXAMPLE 60:

4-[(5-Methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester

Step 1:


4-[(5-Methyl-pyrimidin-4-ylamino)-methyl]-piperidine



A mixture of 4-[(2-chloro-5-methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester (EXAMPLE 59, 2.00g, 5.34mmol), Pd/C (10%, 0.20g) in absolute ethanol (15mL) was vigorously stirred under 1 atm H₂. Filtered and concentrated, the reaction gave 4-[(5-methyl-pyrimidin-4-ylamino)-methyl]-piperidine. M.S. (M+1):207.29

Step 2:

4-[(5-Methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



To a stirred solution of 4-[(5-methyl-pyrimidin-4-ylamino)-methyl]-piperidine (0.20g, 0.97mmol), in DMF (3mL) was added carbonic acid benzyl ester 2,5-dioxo-pyrrolidin-1-yl ester (0.24g, 0.97mmol). The resulting reaction solution was stirred at rt for 0.5h, then concentrated *in vacuo*. The residue was purified by silica gel chromatography (1 – 10 (10% NH_4OH in MeOH) / 99 – 90 CH_2Cl_2) to give 4-[(5-methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester.


^1H NMR (400MHz, CDCl_3) δ 8.50 (s, 1h, Pyr), 7.97 (s, 1h, Pyr), 7.35 (m, 3h, Ar), 5.13 (s, 2h, ArCH_2O), 4.62 (s, 1h, NH), 4.22 (br s, 2h, NCH_2CH_2), 3.43 (s, 2h, NHCH_2CH), 2.79 (br s, 2h, NCH_2CH_2), 2.02 (s, 3h, CH_3), 1.86 (m, 1h, CH), 1.76 (d, $J = 11.7$ Hz, 2h, CHCH_2CH_2), 1.21 (q, $J = 9.7$ Hz, 2h, CHCH_2CH_2);

M.S.(M+):341.39

EXAMPLES 61-63 were prepared as described above in EXAMPLE 60, but replacing the carbonic acid benzyl ester 2,5-dioxo-pyrrolidin-1-yl ester with the appropriately substituted analog:

EXAMPLE 61:

4-[(5-Methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid-4-methyl-benzyl ester




^1H NMR (400MHz, CDCl_3) δ 8.49 (s, 1h, Pyr), 7.97 (s, 1h, Pyr), 7.25 (d, $J = 8.5$ Hz, 2h, Ar), 7.16 (d, $J = 7.9$ Hz, 2h, Ar), 5.08 (s, 2h, ArCH_2O), 4.62 (s, 1h, NH), 4.20 (br s, 2h, NCH_2CH_2), 3.43 (s, 2h, NHCH_2CH), 2.77 (t, $J = 11.0$ Hz, 2h, NCH_2CH_2), 2.35 (s, 3h, PyCH_3), 2.02 (s, 3h, ArCH_3), 1.84 (m, 1h, CH), 1.74 (d, $J = 9.7$ Hz, 2h, CHCH_2CH_2), 1.20 (q, $J = 10.6$ Hz, 2h, CHCH_2CH_2);

M.S.(M+):355.39

EXAMPLE 62:

4-[(5-Methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid-4-chloro-benzyl ester




^1H NMR (400MHz, CDCl_3) δ 8.50 (s, 1h, Pyr), 7.97 (s, 1h, Pyr), 7.34 – 7.26 (m, 4h, Ar), 5.08 (s, 2h, ArCH_2O), 4.62 (s, 1h, NH), 4.20 (br s, 2h, NCH_2CH_2), 3.43 (s, 2h, NHCH_2CH), 2.79 (br s, 2h, NCH_2CH_2), 2.02 (s, 3h, CH_3), 1.85 (m, 1h, CH), 1.76 (d, $J = 12.6$ Hz, 2h, CHCH_2CH_2), 1.20 (q, $J = 10.0$ Hz, 2h, CHCH_2CH_2);

M.S.(M+):375.35

EXAMPLE 63:

4-[(5-Methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid-4-fluoro-benzyl ester



^1H NMR (400MHz, CD_3OD) δ 8.56 (s, 1h, Pyr), 7.96 (s, 1h, Pyr), 7.38 (dd, $J = 5.6$ & 5.4 Hz, 2h, Ar), 7.08 (t, $J = 8.7$ Hz, 2h, Ar), 5.08 (s, 2h, ArCH_2O), 4.14 (d, $J = 13.3$ Hz, 2h, NCH_2CH_2), 6.94 (d, $J = 6.9$ Hz, 2h, NHCH_2CH), 2.81 (br s, 2h, NCH_2CH_2), 2.15 (s, 3h, CH_3), 1.95 (m, 1h, CH), 1.74 (d, $J = 11.4$ Hz, 2h, CHCH_2CH_2), 1.17 (q, $J = 9.2$ Hz, 2h, CHCH_2CH_2);


M.S.(M+):359.36

EXAMPLE 64:

4-[(2-Amino-6-methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester

Step 1:

4-[(2,4-Dimethoxy-benzylamino)-6-methyl-pyrimidin-4-ylamino]-
methyl]-piperidine-1-carboxylic acid benzyl ester




A stirred solution of 4-[(2-chloro-6-methyl-pyrimidin-4-ylamino)-
methyl]-piperidine-1-carboxylic acid benzyl ester (EXAMPLE 57, 0.5g, 1.33mmol)
in 2,4-dimethoxybenzylamine (1.00mL, 6.67mmol) was heated at 100°C for 6h, then
cooled to rt and purified by silica gel chromatography [1 - 10 (10% NH₄OH in
MeOH) / 99 - 90 CH₂Cl₂] to give 4-[(2-(2,4-dimethoxy-benzylamino)-6-methyl-
pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester.

M.S.(M+1):506.46

Step 2:

4-[(2-Amino-6-methyl-pyrimidin-4-ylamino)-methyl]-piperidine-1-
carboxylic acid benzyl ester



To a stirred solution of the 4-[(2,4-dimethoxy-benzylamino)-6-
methyl-pyrimidin-4-ylamino]-methyl]-piperidine-1-carboxylic acid benzyl ester from
Step 1 above (0.4g, 0.79mmol) in CH₂Cl₂ (5mL) was added trifluoroacetic acid
(1mL). The resulting reaction solution was stirred at rt for 1h, then concentrated *in*
vacuo. The residue was purified by silica gel chromatography [1 - 10 (10% NH₄OH
in MeOH) / 99 - 90 CH₂Cl₂] to give 4-[(2-amino-6-methyl-pyrimidin-4-ylamino)-
methyl]-piperidine-1-carboxylic acid benzyl ester. M.S.(M+1):356.36

EXAMPLE 65:

4-[(5,6-Dichloro-pyridazin-4-ylamino)-methyl]-piperidine-1-
carboxylic acid benzyl ester

Step 1:

3,4,5-Trichloropyridazine




A stirred solution of 4,5-dichloro-2,3-dihydro-3-pyridazinone (15.00g,
90.92mmol) in POCl₃ (100mL) was refluxed for 1.5h, then concentrated *in vacuo*.
The residue was dissolved in CH₂Cl₂ (400mL), washed with water (100mL), dried
over Na₂SO₄, filtered and concentrated to give 3,4,5-trichloropyridazine.

M.S.(M+1):185.00

Step 2:

4-[(5,6-Dichloro-pyridazin-4-ylamino)-methyl]-piperidine-1-
carboxylic acid benzyl ester



To a stirred solution of 3,4,5-trichloropyridazine (2.22g, 12.08mmol)
and DIPEA (4.21mL, 24.16mmol) in IPA (25mL) was added benzyl-4-
(aminomethyl)piperidine-1-carboxylate (EXAMPLE 13, Step 1, 3.00g, 12.08mmol).

The resulting reaction solution was stirred at rt for 5h, then concentrated *in vacuo*.
The residue was dissolved in CH₂Cl₂ (200mL), washed with water (50mL), dried over
Na₂SO₄, filtered and concentrated. The crude product was purified by silica gel
chromatography [1 - 7 (10% NH₄OH in MeOH) / 99 - 93 CH₂Cl₂] to give 4-[(5,6-
dichloro-pyridazin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester.

M.S.(M+1):395.28


EXAMPLE 66:

4-[(Pyridazin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester

ester

Step 1:

4-[(Pyridazin-4-ylamino)-methyl]-piperidine



5


A mixture of 4-[(5,6-dichloro-pyridazin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester (EXAMPLE 65, 2.00g, 5.06mmol), Pd/C (10%, 0.20g) in absolute ethanol (15mL) was vigorously stirred under 1atm H₂ (provided by a H₂ balloon for 7h. Filtered and concentrated, the reaction gave 4-[(Pyridazin-4-ylamino)-methyl]-piperidine. M.S. (M+1):193.25

10

Step 2:

4-[(Pyridazin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester

ester



15

To a stirred solution of 4-[(pyridazin-4-ylamino)-methyl]-piperidine (0.20g, 1.04mmol), in DMF (3mL) was added carbonic acid benzyl ester 2,5-dioxo-pyrrolidin-1-yl ester (0.26g, 1.04mmol). The resulting reaction solution was stirred at rt for 0.5h, then concentrated *in vacuo*. The residue was purified by silica gel chromatography (1 - 7 (10% NH₄OH in MeOH) / 99 - 93 CH₂Cl₂) to give 4-[(pyridazin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester.

20

¹H NMR (400MHz, CDCl₃) δ 8.65 (d, J = 6.1 Hz, 1h, Pyr), 8.57 (d, J = 3.1 Hz, 1h, Pyr), 7.36 (m, 5h, Ar), 6.46 (dd, J = 6.1 & 2.9 Hz, 1h, Pyr), 5.13 (s, 2h, ArCH₂O), 4.40 (s, 1h, NH), 4.23 (br s, 2h, NCH₂CH₂), 3.10 (t, J = 6.0 Hz, 2h, NHCH₂CH₂), 2.78 (br s, 2h, NCH₂CH₂), 1.81 (m, 1h, CH), 1.77 (d, J = 12.5 Hz, 2h, CHCH₂CH₂), 1.23 (q, J = 10.3 Hz, 2h, CHCH₂CH₂);


25

M.S. (M+1):327.28

EXAMPLE 67:

- 90 -

4-[(Pyridazin-4-ylamino)-methyl]-piperidine-1-carboxylic acid-4-fluoro-benzyl ester



5


To a stirred solution of 4-[(pyridazin-4-ylamino)-methyl]-piperidine (0.20g, 1.04mmol), from EXAMPLE 66, Step 1) in DMF (3mL) was added carbonic acid-4-fluoro-benzyl ester 2,5-dioxo-pyrrolidin-1-yl ester (0.28g, 1.04mmol). The resulting reaction solution was stirred at rt for 0.5h, then concentrated *in vacuo*. The residue was purified by silica gel chromatography (1-7 (10% NH₄OH in MeOH) / 99-93 CH₂Cl₂) to give 4-[(pyridazin-4-ylamino)-methyl]-piperidine-1-carboxylic acid-4-fluoro-benzyl ester. M.S. (M+1):345.29

10


EXAMPLES 68A and 68B:

EXAMPLE 68A: 4-[(6-Chloro-pyridazin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester

15



EXAMPLE 68B: 4-[(5-Chloro-pyridazin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



20

A mixture of 4-[(5,6-dichloro-pyridazin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester (EXAMPLE 65, 0.15g, 0.38mmol), washed Raney Nickel (0.15g), NH₄OH (1mL) in absolute ethanol (10mL) was vigorously stirred under 1atm H₂ for 7h. The reaction mixture was filtered and concentrated and

- 91 -

the residue was purified by silica gel chromatography (1 - 7 (10% NH_4OH in MeOH) / 99 - 93 CH_2Cl_2) to give 4-[(6-chloro-pyridazin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester. M.S. (M+1): 361.25 and 4-[(5-chloro-pyridazin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester. M.S. (M+1): 361.25

EXAMPLE 69:

4-[(2-Chloro-5-fluoro-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester

Step 1:


2,4-Dichloro-5-fluoro-pyrimidine



A solution of 5-fluoro-uracil (5.00g, 38.44mmol) and N,N-dimethylaniline (5mL) in POCl_3 (20mL) was refluxed for 1h. The solution was then concentrated *in vacuo*. The resulting residue was quenched with water (20mL) at 0°C , and extracted with ether (3 x 150mL). The combined ether layers were washed with water (2 x 50mL), sat. aq. NaHCO_3 , water (50mL), dried over Na_2SO_4 , filtered and concentrated to give 2,4-dichloro-5-fluoro-pyrimidine compound.

Step 2:


4-[(2-Chloro-5-fluoro-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



To a stirred solution of 2,4-dichloro-5-fluoro-pyrimidine (0.67g, 4.03mmol) and triethylamine (0.84mL, 6.04mmol) in DMF (5mL) was added benzyl 4-(aminomethyl)piperidine-1-carboxylate (EXAMPLE 13, Step 1) (1.00g, 4.03mmol). The resulting reaction solution was stirred at rt for 1h, and concentrated *in vacuo*. The residue was purified by silica gel chromatography (CH_2Cl_2 / IPA / hexanes) to give 4-[(2-chloro-5-fluoro-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester. M.S. (M+1): 379.25

EXAMPLE 70:

4-[(5-Fluoro-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



A mixture of 4-[(2-chloro-5-fluoro-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester (EXAMPLE 69, 0.15g, 0.40mmol), washed Raney-Nickel® (0.15g), NH_4OH (1mL) in absolute ethanol (10mL) was vigorously stirred under 1atm H_2 for 2h. The reaction mixture was filtered and concentrated and the residue was purified by silica gel chromatography (1 - 10 (10% NH_4OH in MeOH) / 99 - 90 CH_2Cl_2) to give 4-[(5-fluoro-pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester. M.S. (M+1): 345.28

EXAMPLE 71:

4-[(5-Fluoro-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester

Step 1:


2-Chloro-5-fluoro-pyrimidine



To a refluxing mixture of 2,4-dichloro-5-fluoro-pyrimidine (EXAMPLE 69, Step 1, 3.25g, 19.47mmol) and zinc (8 - 30 mesh, 3.82g, 58.39mmol) in THF (30mL) was slowly added acetic acid (1.11mL, 19.47mmol). This reaction mixture was refluxed for 7h, then cooled to rt, filtered and concentrated to give 2-chloro-5-fluoro-pyrimidine compound.

Step 2:

4-[(5-Fluoro-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester




A solution of benzyl-4-(aminomethyl)piperidine-1-carboxylate

(EXAMPLE 13, Step 1, 0.10g, 0.40mmol), 2-chloro-5-fluoro-pyrimidine (0.053g, 0.40mmol) and triethylamine (0.11mL, 0.81mmol) in DMF (0.5mL) was heated at 100°C for 6h, then concentrated *in vacuo*. The residue was purified by silica gel chromatography (10 CH₂Cl₂ : 1 – 20 IPA : 89 – 70 hexane) to give 4-[(5-fluoro-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester.

M.S.(M+):345.29

10 EXAMPLE 72:

4-[(5-Fluoro-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid-4-methyl-benzyl ester



The solution of (4-methyl-benzyl)-4-(aminomethyl)piperidine-1-


15 carboxylate (INTERMEDIATE 2a) (0.20g, 0.76mmol), 2-chloro-5-fluoro-pyrimidine (EXAMPLE 71, Step 1) (0.10g, 0.76mmol) and triethylamine (0.21mL, 1.53mmol) in DMF (1mL) was heated at 100°C for 6h, then concentrated *in vacuo*. The residue was purified by silica gel chromatography (10 CH₂Cl₂ : 1 – 10 IPA : 89 – 80 hexane) to give 4-[(5-fluoro-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid-4-methyl-benzyl ester. M.S.(M+):359.33

EXAMPLE 73:

4-[(5-Fluoro-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid-4-cyclopropyl-benzyl ester

25 Step 1:

4-Cyclopropyl-benzoic acid ethyl ester




Indium trichloride (2.2g, 10mmol) and THF (50mL) were combined under nitrogen and cooled to -70°C. Cyclopropylmagnesium bromide solution (33mL, 30mmol, 0.92 M) was added dropwise while maintaining the reaction

5 temperature ≤60°C. After the addition was complete, the reaction was stirred 0.5h with cooling then 0.5h with the cooling bath removed. The resulting solution was added via cannula to a refluxing solution of ethyl-4-iodobenzoate (5.5g, 20mmol), *trans*-dichlorobis(triphenylphosphine)palladium(II) (421mg, 0.60mmol) and THF (100mL) under nitrogen. After 24h, the contents of the reaction flask were cooled and the solvent was removed *in vacuo*. Water (100mL) and 5% KHSO₄ were added and the mixture was extracted with CH₂Cl₂ (3x100mL). The combined organic extracts were washed with brine, dried with Na₂SO₄, and filtered. The filtrate was removed *in vacuo* and the remaining residue was purified by flash column chromatography (hexane:EtOAc 95:5) to give 4-cyclopropyl-benzoic acid ethyl ester as an orange oil.

15 Step 2:

(4-Cyclopropyl-phenyl)-methanol




4-Cyclopropyl-benzoic acid ethyl ester (2.46g, 13mmol), and THF (250mL) were combined under nitrogen and cooled in an IPA/dry ice bath to -70°C. Lithium aluminum hydride solution (20mL, 20mmol, 1.0M) was added dropwise.

20 After 2h excess lithium aluminum hydride was quenched by adding EtOAc dropwise. The reaction was warmed to 25°C, then the solvent was removed *in vacuo*. Water (200mL) and a few drops of HCl(aq, 6N) were added. The mixture was extracted with EtOAc (3x100mL). The combined organic extracts were washed with brine, dried with Na₂SO₄, and filtered. The filtrate was removed *in vacuo* and the remaining residue was purified by flash column chromatography (hexane:EtOAc 40:60) to give (4-cyclopropyl-phenyl)-methanol as a colorless oil.


Step 3:

Carbonic acid 4-cyclopropyl-benzyl ester 2,5-dioxo-pyrrolidin-1-yl ester




The title compound was prepared from (4-cyclopropyl-phenyl)-38(L):110-115(1990) and INTERMEDIATE 1A).

Step 4:
4-Aminomethyl-piperidine-1-carboxylic acid 4-cyclopropyl-benzyl ester



The title compound was prepared from carbonic acid 4-cyclopropyl-benzyl ester 2,5-dioxo-pyrrolidin-1-yl ester as described in EXAMPLE 13, Step 1.


Step 5:
4-[(5-Fluoro-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid-4-cyclopropyl-benzyl ester



A solution of (4-cyclopropyl-benzyl)-4-(aminomethyl)piperidine-1-carboxylate (0.10g, 0.35mmol), 2-chloro-5-fluoro-pyrimidine (EXAMPLE 71, Step 1, 0.046g, 0.35mmol) and triethylamine (0.097mL, 0.69mmol) in DMF (1mL) was heated at 100°C for 6h, then concentrated *in vacuo*. The residue was purified by gel chromatography (CH₂Cl₂ / IPA / hexanes) to give 4-[(5-fluoro-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid-4-cyclopropyl-benzyl ester. M.S.(M+1):385.31

EXAMPLE 74:


4-[(5-Fluoro-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid-4-chloro-benzyl ester



A solution of (4-chloro-benzyl)-4-(aminomethyl)piperidine-1-carboxylate (INTERMEDIATE 2B) (0.10g, 0.35mmol), 2-chloro-5-fluoro-pyrimidine (0.047g, 0.35mmol) and triethylamine (0.099mL, 0.71mmol) in DMF (1mL) was heated at 100°C for 6h, then concentrated *in vacuo*. The residue was purified by silica gel chromatography (CH₂Cl₂ / IPA / hexanes) to give 4-[(5-fluoro-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid-4-chloro-benzyl ester. M.S.(M+1):379.26

EXAMPLE 75:


4-[(5-Fluoro-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid-4-fluoro-benzyl ester



A solution of (4-fluoro-benzyl)-4-(aminomethyl)piperidine-1-carboxylate (INTERMEDIATE 2C) (0.10g, 0.38mmol), 2-chloro-5-fluoro-pyrimidine (0.05g, 0.38mmol) and triethylamine (0.11mL, 0.75mmol) in DMF (1mL) was heated at 100°C for 6h, then concentrated *in vacuo*. The residue was purified by silica gel chromatography (CH₂Cl₂ / IPA / hexanes) to give 4-[(5-fluoro-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid-4-fluoro-benzyl ester. M.S.(M+1):363.31

EXAMPLE 76:

4-Methylbenzyl 4-[(2-pyrimidinylamino)methyl]-1-piperidinecarboxylate



A stirred solution of 4-methylbenzyl 4-(aminomethyl)-1-

- 5 piperidinecarboxylate (INTERMEDIATE 2A) (20.00g, 76.23mmol), 2-chloropyrimidine (8.73g, 76.23mmol) and triethylamine (21.25mL, 152.46mmol) in DMF (40mL) was heated at 100°C for 6h. The reaction solution was cooled to rt, then diluted with ethyl acetate (800mL), washed with sat. aq. NaHCO₃ (100mL), water (3 x 100mL), brine (100mL), dried over Na₂SO₄, filtered and concentrated. The residue was purified by silica gel chromatography (CH₂Cl₂ / IPA / hexanes) to give 4-methylbenzyl 4-[(2-pyrimidinylamino)methyl]-1-piperidinecarboxylate. M.S. (M+): 341.30


EXAMPLE 77:

- 15 [1-(2-Phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-pyrimidin-2-yl-

amine

Step 1:

4-Aminomethyl-piperidine-1-carboxylic acid tert-butyl ester




- 20 To a mixture of 4-aminomethylpiperidine (15g) in 250mL of anhydrous tetrahydrofuran cooled to -78°C was added, dropwise over 45min., a solution of di-tert-butyl di-carbonate (24g) in 100mL of anhydrous tetrahydrofuran. After stirring for 1h at -78°C, the mixture was allowed to warm to room temperature and stirred overnight. The mixture was concentrated to near dryness and diluted with 200mL of 10% aqueous citric acid. The mixture was extracted with 3 x 100mL of ether, then made basic with sodium hydroxide pellets and extracted with 3 x 200mL of chloroform. The combined chloroform extracts were dried over magnesium sulfate and concentrated to dryness under reduced pressure. The resulting oil was

homogeneous by TLC (development with 90:10 chloroform saturated with ammonia; methanol).

¹H NMR (400MHz, CDCl₃): δ 4.1 (br s, 2 H), 2.7 (br m, 2H), 2.6 (d, 2H), 1.7 (m, 3H), 1.42 (s, 9H), 1.1 (m, 2H).

Step 2:

5 4-(Benzyloxycarbonylamino-methyl)-piperidine-1-carboxylic acid tert-butyl ester




To a solution of 4-aminomethyl-piperidine-1-carboxylic acid tert-butyl ester (21g) in 100mL of ethyl acetate cooled to 0°C was added 100mL of saturated sodium carbonate and benzyl chloroformate (17g). The solution was stirred for 3h, then separated. The organic layer was dried over magnesium sulfate and concentrated under reduced pressure. Drying under vacuum gave the product as an oil:

¹H NMR (400MHz, CDCl₃): δ 7.35 (m, 5H), 5.3 (d, 1H), 5.1 (s, 2H), 4.1 (br s, 2 H), 3.0 (br m, 2H), 2.6 (br m, 2H), 1.7 (m, 3H), 1.42 (s, 9H), 1.1 (m, 2H).

Step 3:

Piperidin-4-ylmethyl-carbamic acid benzyl ester



A mixture of 4-(benzyloxycarbonylamino-methyl)-piperidine-1-

20 carboxylic acid tert-butyl ester (35g) and 50mL of 4N HCl in dioxane was stirred at room temperature for 3h, then diluted with 200mL of ether and filtered. There was obtained piperidin-4-ylmethyl-carbamic acid benzyl ester hydrochloride salt as a white fluffy solid. The free base was obtained by partitioning the hydrochloride between chloroform (50mL) and saturated aqueous Na₂CO₃ (50mL).


¹H NMR (400MHz, CDCl₃): δ 7.35 (m, 5H), 5.15 (s, 2H), 4.9 (br s, 1 H), 3.1 (m, 2H), 2.6 (m, 3H), 1.7 (m, 2H), 1.6 (m, 2H), 1.1 (m, 2H).

MS (m+1) = 249.

Step 4:

[1-(2-Phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-carbamic acid

30 benzyl ester




A mixture of piperidin-4-ylmethyl-carbamate benzyl ester hydrochloride (2g), 25mL of dichloromethane, trans-2-styrenesulfonyl chloride (1.5g), and 3mL of N,N-diisopropylethylamine was stirred at room temperature overnight, then diluted with 200mL of chloroform and washed with 100mL of saturated sodium carbonate. The chloroform extracts were dried over magnesium sulfate and concentrated. There was obtained 1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl-carbamate benzyl ester as a white solid.

¹H NMR (400MHz, CDCl₃): δ 7.5-7.2 (m, 10H), 6.65 (m, 1H), 5.15 (s, 2H), 4.8 (br s, 1H), 3.8 (d, 2H), 3.1 (dd, 2H), 2.6 (dd, 2H), 1.8 (d, 2H), 1.6 (m, 2H), 1.35 (m, 2H)

MS (m+1) = 415.

Step 5:

C-[1-(2-Phenyl-ethanesulfonyl)-piperidin-4-yl]-methylamine



A mixture of 1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl-carbamate benzyl ester (2.5g), 20% palladium hydroxide (1g) on carbon, 200mL of methanol and 50mL of tetrahydrofuran were shaken under 50psi of hydrogen for 2 days at room temperature. The catalyst was filtered off and washed with 250mL of methanol. Concentration under reduced pressure gave C-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-yl]-methylamine as white solid.


¹H NMR (400MHz, CDCl₃): δ 7.4-7.2 (m, 5H), 5.1 (s, 2H), 3.8 (d, 2H), 3.1 (m, 4H), 2.7 (dd, 2H), 1.8 (d, 2H), 1.6 (m, 5H), 1.3 (m, 2H)

MS (m+1) = 283.

Step 6:

[1-(2-Phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-pyrimidin-2-yl-

amine



A mixture of 0.5g of 1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl-pyrimidin-2-yl-amine, 0.56g of 2-bromopyrimidine, 25mL of 2-propanol and 0.5mL of N,N-diisopropylethylamine was heated to reflux overnight. Purification of the residue obtained after concentration under reduced pressure by chromatography on silica, eluting with ethyl acetate gave 1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl-pyrimidin-2-yl-amine as a white solid.

¹H NMR (400MHz, CDCl₃): δ 8.15 (d, 2H), 7.3-7.18 (m, 5H), 6.5 (dd, 1H), 5.5 (dd, 1H), 3.8 (d, 2H), 3.35 (d, 2H), 3.15 (dd, 4H), 2.7 (m, 2H), 1.9 (d, 2H), 1.8 (m, 1H), 1.3 (m, 2H)

MS (m+1) = 361.


EXAMPLE 78:

[1-[2-(4-Fluoro-phenyl)-ethanesulfonyl]-piperidin-4-yl]methyl]-

pyrimidin-2-yl-amine

Step 1:

1-(2-Chloro-ethyl)-4-fluoro-benzene




A mixture of 7g 2-(4-fluoro-phenyl)-ethanol, 25mL of chlorobenzene, 42mL of 37% HCl, and 0.9g of Aliquat® 336 (tricaprylmethyl ammonium chloride) was heated to reflux for 3 days, cooled and extracted into 3 x 100mL of hexane. The combined extracts were dried over magnesium sulfate and concentrated under reduced pressure. The resulting oil was mainly 1-(2-chloro-ethyl)-4-fluoro-benzene:

¹H NMR (400MHz, CDCl₃): δ 7.3 (dd, 2H), 7.0 (dd, 2H), 3.7 (t, 2H),

3.05 (t, 2H).

Step 2:

Thioacetic acid S-[2-(4-fluoro-phenyl)-ethyl] ester




A mixture of 2.4g of 1-(2-chloro-ethyl)-4-fluoro-benzene, 30mL of DMF and 25mL of potassium thioacetate was stirred under nitrogen for 24h. The mixture was diluted with 200mL of water and extracted with 3 X 50mL of dichloromethane. The combined organic layers were dried over magnesium sulfate and concentrated under reduced pressure. Drying under vacuum gave an oil:

¹H NMR (400MHz, CDCl₃): δ 7.18 (dd, 2H), 6.98 (dd, 2H), 3.08 (t, 2H), 2.81 (t, 2H), 2.32 (s, 3H).

Step 3:

2-(4-Fluoro-phenyl)-ethanesulfonyl chloride




A stream of chlorine gas was dispersed into a stirred, ice cold mixture of 2.5g of thioacetic acid S-[2-(4-fluoro-phenyl)-ethyl] ester, 30mL of dichloromethane and 30mL of water over 1h. The mixture was diluted with 200mL of dichloromethane, shaken and separated. The combined organic layers were dried over magnesium sulfate and concentrated under reduced pressure. Trituration with hexane gave a white solid:

¹H NMR (400MHz, CDCl₃): δ 7.2 (dd, 2H), 7.0 (dd, 2H), 3.1 (dd, 2H), 3.3 (dd, 2H), 2.32 (s, 3H).

Step 4:

4-(tert-Butoxycarbonylamino-methyl)-piperidine-1-carboxylic acid benzyl ester




To an ice cold, stirred solution of 21g of benzyl 4-(aminomethyl)piperidine-1-carboxylate (EXAMPLE 13, Step 1) in 250mL of dichloromethane was added 18g of di-tert-butylidicarbonate in 100mL of

dichloromethane over 30 min. After stirring overnight, the mixture was concentrated to dryness. Trituration with hexane gave a white solid:

¹H NMR (400MHz, CDCl₃): δ 7.4 (m, 5H), 5.15 (s, 2H), 4.6 (br s, 1H), 4.2 (br s, 2H), 3.0 (br s, 2H), 2.8 (m, 2H), 1.7 (m, 3H), 1.42 (s, 9H), 1.15 (m, 2H).

Step 5:

Piperidin-4-ylmethyl-carbamic acid tert-butyl ester




A mixture of 28g of 4-(tert-butoxycarbonylamino-methyl)-piperidine-1-carboxylic acid benzyl ester, 1g of 10% palladium on carbon, 100mL of THF and 200mL of methanol was stirred under an atmosphere of hydrogen for 2 days. The mixture was filtered concentrated under reduced pressure. Drying under reduced pressure gave a white solid:

¹H NMR (400MHz, CDCl₃): δ 4.8 (br s, 1H), 3.05 (d, 2H), 2.9 (dd, 2H), 2.6 (m, 3H), 1.6 (d, 2H), 1.5 (m, 1H), 1.4 (s, 9H), 1.05 (m, 2H).

Step 6:


{1-[2-(4-Fluoro-phenyl)-ethanesulfonyl]-piperidin-4-ylmethyl}-carbamic acid tert-butyl ester



To an ice cold, stirred solution of 0.2g of piperidin-4-ylmethyl-carbamic acid tert-butyl ester and 0.2mL of N,N-diisopropylethylamine in 20mL of dichloromethane was added 0.3g of 2-(4-fluoro-phenyl)-ethanesulfonyl chloride. After stirring overnight the mixture was diluted with 50mL of chloroform, washed with 50mL of saturated sodium carbonate, dried over magnesium sulfate and concentrated to dryness under reduced pressure. Trituration with hexane gave a white solid:


¹H NMR (400MHz, CDCl₃): δ 7.2 (m, 2H), 7.0 (dd, 2H), 4.6 (br m, 1H), 3.8 (d, 2H), 3.1 (m, 3H), 3.0 (m, 2H), 2.7 (dd, 2H), 1.8 (d, 2H), 1.6 (br m, 2H), 1.42 (s, 9H), 1.3 (m, 2H).

Step 7:
C-{1-[2-(4-Fluoro-phenyl)-ethanesulfonyl]-piperidin-4-yl}-
methylamine



5 A mixture of 0.4g of {1-[2-(4-fluoro-phenyl)-ethanesulfonyl]-
piperidin-4-ylmethyl}-carbamate in 5mL of 4N HCl in dioxane
was stirred at room temperature for 3h, then diluted with 50mL of chloroform, washed
with 50mL of saturated sodium carbonate, dried over magnesium sulfate and
concentrated to dryness under reduced pressure. The product was a white solid:
10 ¹H NMR (400MHz, CDCl₃): δ 7.2 (m, 2H), 7.0 (dd, 2H), 3.92 (d, 2H),
3.1 (s, 4H), 2.7 (dd, 2H), 2.6 (d, 2H), 1.8 (d, 2H), 1.5 (br m, 3H), 1.3 (m, 2H)
MS (m+1) = 301.

Step 8:
15 {1-[2-(4-Fluoro-phenyl)-ethanesulfonyl]-piperidin-4-ylmethyl}-
pyrimidin-2-yl-amine




A mixture of 0.3g of C-{1-[2-(4-Fluoro-phenyl)-ethanesulfonyl]-
piperidin-4-yl}-methylamine, 0.3g of 2-bromopyrimidine, 25mL of 2-propanol and
0.3mL of N,N-diisopropylethylamine was heated to reflux overnight. Purification of
the residue obtained after concentration under reduced pressure by preparative
chromatography, eluting with ethyl acetate gave a white solid.
20 ¹H NMR (400MHz, CDCl₃): δ 8.25 (d, 2H), 7.2 (m, 2H), 7.0 (dd,
2H), 6.58 (dd, 1H), 5.25 (br m, 1H), 3.82 (d, 2H), 3.4 (dd, 2H), 3.15 (s, 4H), 2.75 (dd,
2H), 1.9 (d, 2H), 1.8 (m, 1H), 1.3 (m, 2H)
MS (m+1) = 379.

EXAMPLE 79:

ester
3-(Pyrimidin-2-ylaminomethyl)-pyrrolidine-1-carboxylic acid benzyl

Step 1:

5 1-Benzyl-pyrrolidine-3-carboxylic acid amide




To a mixture of 4.4g 1-benzyl-pyrrolidine-3-carboxylic acid methyl
ester (M. J. Kornet, P. A. Thio, S. E. Tan, *J. Organic Chemistry*, 23:3637-3639(1968)
and 3g formamide in 10mL of anhydrous DMF heated to 100°C, a solution of sodium
methoxide, from 0.33g of sodium dissolved in methanol, was added dropwise over 20
10 minutes. After stirring for 1h at 100°C, the mixture was allowed to cool to room
temperature and added to 100mL of isopropanol. The mixture was concentrated to
dryness. The residue was triturated with 200mL of chloroform, filtered and
concentrated to dryness under reduced pressure. The resulting oil was fairly
homogeneous by TLC (development with 90:10 chloroform saturated with ammonia:
15 methanol):

¹H NMR (400MHz, CDCl₃): δ 7.1 (5H), 4.3 (br s, 2 H), 3.5 (d, 2H),
3.4 (m, 1H), 2.6 (m, 2H), 2.5 (m, 1H), 2.25 (m, 1H), 1.9 (m, 1H).

Step 2:

20 3-Carbamoyl-pyrrolidine-1-carboxylic acid benzyl ester




A mixture of 4.5g 1-benzyl-pyrrolidine-3-carboxylic acid amide,
200mL THF, 20mL methanol and 1g 20% palladium hydroxide on carbon was shaken
under 50psi of hydrogen for 12h. The catalyst was filtered off and the filtrate
concentrated under reduced pressure. Drying under vacuum gave 3g of an oil. To a
25 stirred solution of the crude residue in 500mL of chloroform was added 5.5g of N-
(benzyloxycarbonyloxy)succinimide and 2.2mL of triethylamine. The mixture was
allowed to stir overnight then washed with 50mL of saturated sodium carbonate, dried

over magnesium sulfate, and concentrated to dryness. Purification by chromatography on silica gel, eluting with 90:10 ethyl acetate: methanol, gave 3-carbamoyl-pyrrolidine-1-carboxylic acid benzyl ester:

¹H NMR (400MHz, CDCl₃): δ 7.35 (m, 5H), 5.6 (br m, 2H), 3.6 (m, 3H), 3.4 (m, 1H), 2.9 (br m, 1H), 2.1 (m, 2H).

Step 3:

3-Aminomethyl-pyrrolidine-1-carboxylic acid benzyl ester




A mixture of 1g 3-carbamoyl-pyrrolidine-1-carboxylic acid benzyl ester and 24mL 1M borane-THF was stirred at room temperature for 24h, then carefully quenched with 50mL of 3N HCl. The mixture was concentrated under reduced pressure, then partitioned between 50mL chloroform and 25mL saturated aqueous sodium carbonate. Concentration of the combined extracts after drying over magnesium sulfate gave 3-aminomethyl-pyrrolidine-1-carboxylic acid benzyl ester:

¹H NMR (400MHz, CDCl₃): δ 7.35 (m, 5H), 5.15 (s, 2H), 3.7-4 (complex, 4H), 2.7 (m, 1H), 2.4-2.0 (complex, 2H), 1.6 (m, 4H).

Step 4:

3-(Pyrimidin-2-ylaminomethyl)-pyrrolidine-1-carboxylic acid benzyl ester



A mixture of 3-aminomethyl-pyrrolidine-1-carboxylic acid benzyl ester (0.15g), 2-bromopyrimidine (0.25g), 2-propanol (10mL), and of N,N-diisopropylethylamine (0.1mL) was heated to reflux overnight. Purification of the residue obtained after concentration under reduced pressure by preparative chromatography, and eluting with ethyl acetate, gave 3-(pyrimidin-2-ylaminomethyl)-pyrrolidine-1-carboxylic acid benzyl ester as a solid:

¹H NMR (400MHz, CDCl₃): δ 8.15 (d, 2H), 7.3 (m, 5H), 6.5 (dd, 1H), 5.8 (m, 1H), 5.1 (s, 2H), 3.5 (m, 2H), 3.4 (m, 3H), 3.2 (m, 1H), 2.55 (m, 1H), 2.0 (m, 1H), 1.7 (m, 1H)

MS (m+1) = 313.


EXAMPLE 80:

(R,S) 4-[1-(Pyridin-4-ylamino)-ethyl]-piperidine-1-carboxylic acid

benzyl ester

Step 1:

4-Acetyl-piperidine-1-carboxylic acid benzyl ester




To a solution of 5g of 4-(N-methoxy-N-methyl-carbamoyl)-piperidine-1-carboxylic acid benzyl ester (S. Nahm and S. W. Weinreb, *Tetrahedron Letters*, 22:3815-3818(1981)) in 50mL of anhydrous THF cooled to 0°C, was added dropwise 6mL of 3M methylmagnesium bromide in ether over 10 minutes. After stirring for 1h at 0°C, the resulting mixture was quenched with 50mL of 1N HCl and extracted with 3 x 50mL of ether. The combined extracts were dried over magnesium sulfate and concentrated to dryness under reduced pressure. Drying under vacuum gave 4-Acetyl-piperidine-1-carboxylic acid benzyl ester as a white solid:

¹H NMR (400MHz, CDCl₃): δ 7.35 (m, 5H), 5.15 (s, 2H), 4.2 (br s, 2H), 2.9 (br t, 2H), 2.5 (m, 1H), 2.2 (s, 3H), 1.9 (m, 2H), 1.6 (m, 2H).

Step 2:

4-(1-Hydroxyimino-ethyl)-piperidine-1-carboxylic acid benzyl ester




A mixture of 4.0g of 4-acetyl-piperidine-1-carboxylic acid benzyl ester, 25mL of pyridine, and 6g of hydroxylamine hydrochloride were heated to 100°C for 12h. The mixture was concentrated under reduced pressure and partitioned between 200mL of ethyl acetate and 50mL of 1N HCl. The organic extract was dried over magnesium sulfate and concentrated to dryness under reduced pressure. Drying

under vacuum gave 4-(1-(1-hydroxyimino-ethyl)-piperidine-1-carboxylic acid benzyl ester as a solid:

¹H NMR (400MHz, CDCl₃): δ 7.35 (m, 5H), 5.15 (s, 2H), 4.3 (br s, 2H), 2.8 (br t, 2H), 2.3 (m, 1H), 2.05 and 1.85 (2s, 3H), 1.8 (m, 2H), 1.5 (m, 2H).

Step 3:

4-(1-(1-Hydroxyimino-ethyl)-piperidine-1-carboxylic acid tert-butyl ester




A mixture of 3.2g of 4-(1-(1-hydroxyimino-ethyl)-piperidine-1-carboxylic acid benzyl ester, 0.4g of di-tert-butyl dicarbonate, 0.15g of 10% palladium on carbon and 20mL of THF was stirred under atmosphere of hydrogen for 2h. The mixture was filtered and concentrated under reduced pressure. Drying under vacuum gave 4-(1-(1-hydroxyimino-ethyl)-piperidine-1-carboxylic acid tert-butyl ester:

¹H NMR (400MHz, CDCl₃): δ 4.15 (br s, 2H), 2.7 (br t, 2H), 2.25 (m, 1H), 1.8 (s, 3H), 1.7 (m, 2H), 1.42 (m, 2H), 1.4 (s, 9H).

Step 4:

(R,S) 4-[1-(Amino-ethyl)-piperidine-1-carboxylic acid tert-butyl ester




A mixture of 3g of 4-(1-(1-hydroxyimino-ethyl)-piperidine-1-carboxylic acid tert-butyl ester, 5g of wet Raney-nickel and 100mL of 5% ammonia in ethanol was shaken under 55psi of hydrogen for 12h. The mixture was filtered and concentrated under reduced pressure. The resulting crude product was taken up in 250mL of chloroform, dried over magnesium sulfate, and concentrated under reduced pressure. Drying under vacuum gave (R,S) 4-(1-(1-amino-ethyl)-piperidine-1-carboxylic acid tert-butyl ester:

¹H NMR (400MHz, CDCl₃): δ 4.05 (br s, 2H), 2.6 (br m, 3H), 2.25 (m, 1H), 1.6 (dd, 2H), 1.4 (s, 9H), 1.2 (m, 2H), 1.1 (m, 2H), 1.0 (d, 3H).

Step 5:

(R,S) 4-[1-(2-Pyridin-4-ylamino-ethyl)-piperidine-1-carboxylic acid

tert-butyl ester




A mixture of 3g of 4-(1-(1-amino-ethyl)-piperidine-1-carboxylic acid tert-butyl ester, 2.5g of 4-bromopyridine hydrochloride, 3.6g of sodium tert-butoxide, 0.14g of palladium acetate, 0.38g of racemic BINAP and 50mL of THF was heated to reflux for 12h. The mixture was cooled, diluted with 50mL of water and concentrated under reduced pressure. The resulting residue was partitioned between 500mL of chloroform and 200mL of water. The extracts were dried over magnesium sulfate and concentrated under reduced pressure. Purification by chromatography, eluting with 90:10 chloroform saturated with ammonia: methanol gave (R,S) 4-[1-(pyridin-4-ylamino)-ethyl]-piperidine-1-carboxylic acid tert-butyl ester resin:

¹H NMR (400MHz, CDCl₃): δ 8.15 (d, 2H), 6.4 (d, 2H), 4.3 (d, 1H), 4.15 (br s, 2H), 3.2 (m, 1H), 2.65 (m, 2H), 2.5 (m, 1H), 1.7 (dd, 2H), 1.6 (m, 1H), 1.42 (s, 9H), 1.25 (m, 2H), 1.15 (m, 2H), 1.1 (d, 3H).

Step 6:

(R,S) 4-[1-(2-Pyridin-4-ylamino-ethyl)-piperidine-1-carboxylic acid benzyl ester



A mixture of 0.1g of 4-[1-(pyridin-4-ylamino)-ethyl]-piperidine-1-carboxylic acid tert-butyl ester and 10mL of 4N HCl in dioxane was stirred at room temperature for 2h, then concentrated to dryness. The residue was diluted with 50mL of chloroform and 1mL of saturated sodium carbonate, cooled to 0°C and treated with 0.05mL of benzyl chloroformate. The resulting solution was allowed to stir for 3h then separated. The organic layer was dried over magnesium sulfate and concentrated under reduced pressure. Purification by preparative chromatography eluting with 90:10 chloroform saturated with ammonia: methanol gave (R,S) 4-[1-(pyridin-4-ylamino)-ethyl]-piperidine-1-carboxylic acid benzyl ester:


¹H NMR (400MHz, CDCl₃): δ 8.15 (d, 2H), 7.3 (m, 5H), 6.4 (d, 2H), 4.38 (d, 1H), 4.15 (br s, 2H), 3.4 (m, 1H), 2.9 (m, 1H), 2.75 (m, 2H), 1.65 (dd, 2H), 1.6 (m, 1H), 1.32 (m, 4H), 1.1 (d, 3H)

MS (m+1) = 340.

The following **EXAMPLES 81-103** were prepared from a primary amine described herein and a chloro-substituted heterocycle using conditions and procedures similar to those described in **EXAMPLE 77**, Step 6 unless otherwise stated:

EXAMPLE 81:


N2-[1-(2-Phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-quinazoline-2,4-diamine



EXAMPLE 81 was prepared from C-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-yl]-methylamine and 2-chloro-quinazolin-4-ylamine (2-chloro-quinazolin-4-ylamine was prepared from 2,4-dichloroquinazoline and ammonia in THF at room temperature; N.B. Chapman, G. M. Gibson, F.G. Mann, *J. Chem. Soc.*, 1947, 890-899); MS (m+1) = 426.

EXAMPLE 82:


[1-(2-Phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-(9H-purin-2-yl)-amine



EXAMPLE 82 was prepared from C-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-yl]-methylamine and 2-chloro-9H-purine (2-chloro-9H-purine was prepared according to S. R. Brashears, S. S. Wang, S. G. Bechtolt, B. E. Christensen, *J. Am. Chem. Soc.*, 81:3789-3792(1959)); MS (m+1) = 401.

EXAMPLE 83:


2-[1-(2-Phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-amino)-pyrimidine-4-carboxylic acid amide



EXAMPLE 83 was prepared from C-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-yl]-methylamine and 2-chloro-pyrimidine-4-carboxylic acid amide (2-chloro-pyrimidine-4-carboxylic acid amide was prepared according to G. D. Davies, D. E. O'Brien, L. R. Lewis, C. C. Cheng, *J. Heterocyclic Chem.*, 1:130-131(1964)); MS (m+1) = 404.

EXAMPLE 84:


(9-Methyl-9H-purin-6-yl)-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-amine



EXAMPLE 84 was prepared from C-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-yl]-methylamine and 6-chloro-9-methyl-9H-purine (6-chloro-9-methyl-9H-purine prepared according to G. B. Eilon, *J. Org. Chem.*, 27:2478-2491(1962)); MS (m+1) = 415.

EXAMPLE 85:


(7-Methyl-7H-purin-6-yl)-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-amine



EXAMPLE 85 was prepared from C-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-yl]-methylamine and 6-chloro-7-methyl-7H-purine (6-chloro-7-methyl-7H-purine was prepared according to G. B. Eilon, *J. Org. Chem.*, 27:2478-2491(1962)); MS (m+1) = 415.

EXAMPLE 86:


4-(Pteridin-4-ylaminomethyl)-piperidine-1-carboxylic acid benzyl ester



EXAMPLE 86 was prepared from 4-aminomethyl-piperidine-1-carboxylic acid benzyl ester and 4-methylthio-pteridine (4-methylthio-pteridine was prepared according to A. A. Brown, D. J. Brown, h. C. S. Wood, *J. Chem. Soc.*, 1954, 3832-3839). MS (m+1) = 379.

EXAMPLE 87:


4-[(7H-Pyrrolo[2,3-d]pyrimidin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



EXAMPLE 87 was prepared from 4-aminomethyl-piperidine-1-carboxylic acid benzyl ester and 4-chloro-7H-pyrrolo[2,3-d]pyrimidine (4-chloro-7H-pyrrolo[2,3-d]pyrimidine was prepared according to U. Lupke, F. Seela, *Chem. Ber.*, 112:3832-3839(1979). MS (m+1) = 366.

EXAMPLE 88:

4-[(1H-Imidazo[4,5-c]pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester




EXAMPLE 88 was prepared from 4-aminomethyl-piperidine-1-carboxylic acid benzyl ester and 7-chloro-3H-imidazo[4,5-b]pyridine (7-chloro-3H-

imidazo[4,5-b]pyridine was prepared according to Y. Mizuno, T. Itoh, K. Saio, *Chem. Pharm. Bull.*, 12:866-872(1964). MS (m+1) = 366.

EXAMPLE 89:


5 (3-Chloro-pyrazin-2-yl)-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-amine



EXAMPLE 89 was prepared from C-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-yl]-methylamine and 2,3-dichloropyrazine (refluxing 2-butanol). MS (m+1) = 396.

EXAMPLE 90:


[1-(2-Phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-pyrazin-2-yl-amine



EXAMPLE 90 was prepared from (3-chloro-pyrazin-2-yl)-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-amine by hydrogenation in ethanol-triethylamine over 5% palladium on carbon, 1 atm of hydrogen. MS (m+1) = 361.

EXAMPLE 91:


20 (2-Chloro-5-methyl-pyrimidin-4-yl)-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-amine



EXAMPLE 91 was prepared from C-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-yl]-methylamine and 2,4-dichloro-5-methyl-pyrimidine: MS (m+1) = 410.

5 EXAMPLE 92:


(5-Methyl-pyrimidin-4-yl)-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-amine



EXAMPLE 92 was prepared from (2-chloro-5-methyl-pyrimidin-4-yl)-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-amine by hydrogenation in ethanol-triethylamine over 5% palladium on carbon, 1atm of hydrogen: MS (m+1) = 375.5.

15 EXAMPLE 93:

[1-(2-Phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-pyrimidin-4-yl-amine



EXAMPLE 93 was prepared from C-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-yl]-methylamine and 2,4-dichloro-pyrimidine followed by hydrogenation in ethanol-triethylamine over 5% palladium on carbon, 1atm of hydrogen: MS (m+1) = 361.5.


25 EXAMPLE 94:

(4-Methyl-pyrimidin-2-yl)-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-amine

EXAMPLE 94 was prepared from C-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-yl]-methylamine and 2-chloro-4-methyl-pyrimidine: MS (m+1) = 375.5.

5 EXAMPLE 95:


5-Fluoro-N2-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-pyrimidine-2,4-diamine



EXAMPLE 95 was prepared from C-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-yl]-methylamine and 2-chloro-5-fluoro-pyrimidin-4-ylamine: MS (m+1) = 394.5.

15 EXAMPLE 96:


N2-[1-(2-Phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-pyrimidine-2,4-diamine



EXAMPLE 96 was prepared from C-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-yl]-methylamine and 2-chloro-pyrimidin-4-ylamine (prepared from 2,4-chloro-pyrimidin-4-ylamine by hydrogenation in ethanol over 5% palladium on carbon, 1atm of hydrogen): MS (m+1) = 376.5.

25 EXAMPLE 97:


(3-Methyl-pyrazin-2-yl)-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-amine



EXAMPLE 97 was prepared from C-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-yl]-methylamine and 3-bromo-pyrazine-2-carboxylic acid methyl ester followed by reduction with lithium tri-sec-butylborohydride at 0°C in THF: MS (m+1) = 375.5.

EXAMPLE 98:


[1-[2-(2-Fluoro-phenyl)-ethanesulfonyl]-piperidin-4-ylmethyl]-pyrimidin-2-yl-amine



EXAMPLE 98 was prepared from 2-(2-fluoro-phenyl)-ethanol as described in **EXAMPLE 78**, Steps 1-7 above: MS (m+1) = 378.5.

EXAMPLE 99:


[1-[2-(4-Chloro-phenyl)-ethanesulfonyl]-piperidin-4-ylmethyl]-pyrimidin-2-yl-amine



EXAMPLE 99 was prepared from 2-(4-chloro-phenyl)-ethanol as described in **EXAMPLE 78**, Steps 1-7 above: MS (m+1) = 396.

EXAMPLE 100:


Pyrimidin-2-yl-[1-(2-p-tolyl-ethanesulfonyl)-piperidin-4-ylmethyl]-amine



EXAMPLE 100 was prepared from 2-(4-methyl-phenyl)-ethanol as described in **EXAMPLE 78**, Steps 1-7 above: MS (m+1) = 375.5.

EXAMPLE 101:


3-(Pteridin-4-ylaminomethyl)-pyrrolidine-1-carboxylic acid benzyl ester



EXAMPLE 101 was prepared from 3-aminomethyl-pyrrolidine-1-carboxylic acid benzyl ester (**EXAMPLE 79**, Step 3) and 4-methylthio-pteridine (A. Brown, D. J. Brown, J. C. S. Wood, *J. Chem. Soc.*, 1954, 3832-3839): MS (m+1) = 365.4.

EXAMPLE 102:


3-[(9H-Purin-6-ylamino)-methyl]-pyrrolidine-1-carboxylic acid benzyl ester



EXAMPLE 102 was prepared from 3-aminomethyl-pyrrolidine-1-carboxylic acid benzyl ester (**EXAMPLE 79**, Step 3) and 6-chloro-9H-purine: MS (m+1) = 353.4.


EXAMPLE 103:

3-Nitro-N⁶-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-pyridine-2,6-diamine




EXAMPLE 103 was prepared from C-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-yl]-methylamine and 6-chloro-3-nitro-pyridin-2-ylamine: MS (m+1) = 420.5.

EXAMPLE 104:
(1H-Imidazo[4,5-b]pyridin-5-yl)-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-amine



EXAMPLE 104 was prepared from 3-nitro-N⁶-[1-(2-phenyl-ethanesulfonyl)-piperidin-4-ylmethyl]-pyridine-2,6-diamine (**EXAMPLE 103**) (1mmol scale) by hydrogenation in 15mL of THF/methanol over 0.5 of Raney-nickel under 1atm of hydrogen for 1h, followed by immediate conversion of the crude, air sensitive triaminopyridine into the imidazo[4,5b]pyridine by heating with 5mL of 96% formic acid and 2mL of 37% hydrochloric acid at reflux overnight. The free base was liberated with sodium hydroxide and purified by preparative chromatography, eluting with 90:10 chloroform: methanol: MS (m+1) = 400.5.


EXAMPLE 105:
4-[(1H-Benzimidazol-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



EXAMPLE 105 was prepared from 1H-benzimidazol-4-ylamine (The 1H-benzimidazol-4-ylamine was prepared by heating 1.5g of 3-nitro-benzene-1,2-diamine in 50mL of triethyl orthoformate with 10mg of p-toluenesulfonic acid monohydrate at reflux overnight, concentration to dryness under reduced pressure, hydrolysis with refluxing 3N HCl for 1h and neutralization with NaOH. Then, cooling and collection yielded the 4-nitro-benzimidazole product by filtration.

Catalytic reduction using Raney Nickel® in ethanol under 1atm of hydrogen for 1h gave 1H-benzimidazol-4-ylamine as an air sensitive solid) and 4-formyl-piperidine-1-carboxylic acid benzyl ester (prepared from 4-(N-methoxy-N-methyl-carbamoyl)-piperidine-1-carboxylic acid benzyl ester, using the procedures described by S. Nahm and S. W. Weinreb, *Tetrahedron Letters*, 22:3815-3818(1981)) on a 1mmol scale by reductive amination in 5mL of 1,2-dichloromethane using sodium triacetoxyborohydride over 0.5 of Raney Nickel® under 1atm of hydrogen for 1h, followed by immediate conversion of the crude, air sensitive triaminopyridine into the imidazo[4,5b]pyridine by heating with 5mL of 96% formic acid and 2mL of 37% hydrochloric acid at reflux overnight. The free base was liberated with sodium hydroxide and purified by preparative chromatography, eluting with 90:10 chloroform: methanol: MS (m+1) = 363.5.

EXAMPLE 106:
4-[(3-Hydroxy-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



EXAMPLE 106 was prepared from 4-(3-hydroxy-pyridin-4-yl-carbamoyl)-piperidine-1-carboxylic acid benzyl ester (which was prepared by EDC coupling of 4-amino-pyridin-3-ol and N-benzyloxycarbonyl piperidine-4-carboxylic acid) by borane-THF reduction overnight at room temperature. The reaction was quenched by slow addition of 1N HCl until pH = 2, then basified to pH = 10 with 10 N NaOH. Extraction with chloroform yielded a crude product which was purified by preparative chromatography, eluting with 90:10 chloroform saturated with ammonia: methanol to give 4-[(3-hydroxy-pyridin-4-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester: MS (m+1) = 342.4.

EXAMPLE 107:

3-*exo*-(Pyridin-4-ylaminomethyl)-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester hydrochloride

Step 1:

(8-Benzyl-8-azabicyclo[3.2.1]oct-3-*exo*-yl)methylamine




In a three-neck flask equipped with an addition funnel, a nitrogen inlet, and a rubber septum was placed a 1M solution of lithium aluminum hydride in tetrahydrofuran (5.5mL, 5.5mmol). To that solution, a solution of 8-benzyl-8-azabicyclo[3.2.1]octane-3-*exo*-carboximide (EP 31219 A1 19810701) (1.13g, 5.0mmol) in dry tetrahydrofuran was added dropwise via syringe. The resulting mixture was stirred 3 hours at 60°C. The mixture was cooled in an ice-bath and 3N sodium hydroxide solution (25mL) was added dropwise. The mixture was extracted with ethyl acetate (2x100mL). The combined extract was washed with water (50mL) and brine (50mL), dried (sodium sulfate), filtered, and the solvent was evaporated under reduced pressure to give crude (8-benzyl-8-azabicyclo[3.2.1]oct-3-*exo*-yl)methylamine product as an oil.

¹H NMR (CDCl₃) δ 7.38 (2H, d, J 7 Hz), 7.34–7.23 (3H, m), 3.54 (2H, s), 3.21 (2H, m), 2.55 (2H, d, J 6.5 Hz), 2.01 (2H, m), 1.67 (1H, m), 1.60 (2H, d, J 8 Hz), 1.56–1.34 (6H, m).

Mass spec.: 231.50 (M+1).

Step 2:

(8-Benzyl-8-aza-bicyclo[3.2.1]oct-3-*exo*-ylmethyl)pyridin-4-yl-amine



To a mixture of (8-benzyl-8-azabicyclo[3.2.1]oct-3-*exo*-yl)methylamine (0.999g, 4.3mmol), 4-bromopyridine hydrochloride (0.719g,


3.7mmol), palladium acetate (0.033g, 0.15mmol), and (±)-BINAP (0.092g, 0.15mmol) in tetrahydrofuran (34mL) under nitrogen, was added sodium t-butoxide (0.86g, 8.9mmol). The mixture was stirred at 70°C under nitrogen for 18h. The mixture was diluted with ether (35mL), washed with brine (2x35mL), dried (sodium sulfate), filtered, and the solvent was evaporated under reduced pressure to give crude product (1.42g) as a brown gum. The crude product was flash chromatographed on silica gel, eluting first with methanol: methylene chloride (10:90) to remove impurities, then with methanol: methylene chloride: ammonium hydroxide (10:90:1 increasing to 20:80:2) to give a yellow foam (1.08g). The foam was triturated with ether to give a crystalline solid. The solid was filtered off and dried in vacuo to give (8-benzyl-8-aza-bicyclo[3.2.1]oct-3-*exo*-ylmethyl)pyridin-4-yl-amine product as a yellow solid.

¹H NMR (CDCl₃) δ 8.16 (2H, m), 7.39 (2H, d, J 1.5 Hz), 7.32 (2H, m), 7.26 (1H, m), 6.41 (2H, m), 4.25 (1H, br s), 3.55 (2H, s), 3.25 (2H, m), 3.02 (2H, t, J 6 Hz), 2.05 (2H, m), 1.97 (1H, m), 1.55 (6H, m).

Mass spec.: 308.36 (M+1).

Step 3:

(8-Benzyl-8-aza-bicyclo[3.2.1]oct-3-*exo*-ylmethyl)pyridin-4-yl-carbamate acid *tert*-butyl ester




A mixture of (8-benzyl-8-aza-bicyclo[3.2.1]oct-3-*exo*-ylmethyl)pyridin-4-yl-amine (0.707g, 2.3mmol), 4-dimethylaminopyridine (0.037g, 0.30mmol, 0.13 equiv.), and di-*tert*-butyl dicarbonate (0.79g, 3.6mmol) in acetonitrile was stirred under nitrogen at ambient temperature for 18h. The mixture was concentrated under reduced pressure and the residue was taken up in methylene chloride (60mL). The mixture was washed with saturated sodium bicarbonate solution (30mL), water (30mL), and brine (30mL), dried (sodium sulfate), filtered, and the solvent was evaporated under reduced pressure to give a crude product (0.96g) as an orange gum. The crude product was flash chromatographed on silica gel eluting first with methanol: methylene chloride (10:90), then with methanol: methylene

chloride: ammonium hydroxide (10:90:1) to give (8-benzyl-8-aza-bicyclo[3.2.1]oct-3-*exo*-ylmethyl)pyridin-4-yl-carbamic acid *tert*-butyl ester product as a yellow oil.

¹H NMR (CDCl₃) δ 8.52 (2H, m), 7.40-7.23 (5H, m), 7.19 (2H, m), 3.60 (2H, d, J 7 Hz), 3.51 (2H, m), 3.18 (2H, br s), 1.99 (3H, m), 1.48 (9H, s), 1.42 (6H, m).

5 Step 4:

(8-Aza-bicyclo[3.2.1]oct-3-*exo*-ylmethyl)pyridin-4-yl-carbamic acid *tert*-butyl ester




10 A mixture of (8-benzyl-8-aza-bicyclo[3.2.1]oct-3-*exo*-ylmethyl)pyridin-4-yl-carbamic acid *tert*-butyl ester (0.917g, 2.25mmol) and 10% palladium on carbon (0.60g) in methanol (25mL) was hydrogenated (53psi hydrogen) for 18 h. The catalyst was removed by filtration through Celite. The filter cake was washed with methanol (3x25mL) and the filtrate was concentrated under reduced pressure to give crude product (0.592g) as a gum. The crude product was flash chromatographed on silica gel eluting with methanol: methylene chloride: ammonium hydroxide (10:90:1 increasing to 20:80:2) to give product as a solid white foam.

15 ¹H NMR (CDCl₃) δ 8.53 (2H, m), 7.19 (2H, m), 3.80 (2H, s), 3.64 (2H, d, J 7 Hz), 2.6-2.0 (1H, br s), 2.10 (1H, m), 2.07 (2H, m), 1.63 (6H, m), 1.48 (9H, s).

20 Step 5:

3-*exo*-[(*tert*-Butoxycarbonyl-pyridin-4-yl-amino)methyl]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester




To a rapidly stirred mixture of (8-aza-bicyclo[3.2.1]oct-3-*exo*-ylmethyl)pyridin-4-yl-carbamic acid *tert*-butyl ester (95mg, 0.30mmol), sodium bicarbonate (76mg, 0.90mmol), methylene chloride (0.8mL), and water (0.8mL) cooled in an ice-bath, was added benzyl chloroformate (57μL, 68mg, 0.40mmol). The mixture was stirred 18h while warming from ice-bath to ambient temperature. The mixture was diluted with dichloromethane (5mL) and the layers were separated. The organic layer was washed with water (2mL), and brine (2mL), dried (sodium sulfate), filtered, and the solvent was evaporated under reduced pressure to give a crude product (112mg) as a pale yellow oil. The crude product was chromatographed on a 2mm silica gel prep plate eluting with ethyl acetate : hexane (3:2) to give 3-*exo*-[(*tert*-butoxycarbonyl-pyridin-4-yl-amino)methyl]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester product as a colorless gum.

¹H NMR (CDCl₃) δ 8.53 (2H, d, J 6 Hz), 7.34 (5H, m), 7.17 (2H, d, J 6 Hz), 5.12 (2H, s), 4.29 (2H, br s), 3.56 (2H, d, J 7 Hz), 2.17 (1H, m), 1.92 (2H, m), 1.55-1.31 (15H, m).

15 Step 6:

3-*exo*-(Pyridin-4-ylaminomethyl)-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester hydrochloride



20

Into a solution of 3-*exo*-[(*tert*-butoxycarbonyl-pyridin-4-yl-amino)methyl]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester (54mg,

0.12mmol) in ethyl acetate (1mL), cooled in an ice-bath, was bubbled hydrogen chloride for 2 minutes. The solution was stirred one hour with ice-bath cooling, degassed with nitrogen, then concentrated under reduced pressure. The residual gum was dissolved in methylene chloride (0.5mL) and the solution was diluted with ether (5mL) to deposit a gum. The supernatant was decanted, the gum was triturated with ether, and the resulting solid was filtered off and dried in vacuo to give 3-*exo*-(pyridin-4-ylaminomethyl)-8-*aza*-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester hydrochloride as an off-white solid.

¹H NMR (DMSO-*d*₆) δ 13.34 (1H, br s), 8.68 (1H, m), 8.19 (1H, br s), 8.06 (1H, br s), 7.36 (5H, m), 6.90 (2H, d, 7.7 Hz), 5.08 (2H, s), 4.20 (2H, br s), 3.11 (2H, t, 7.6 Hz), 2.17 (1H, m), 1.88 (2H, m), 1.65 (4H, m), 1.31 (2H, m).

Mass spec.: 352.41 (M+1).

EXAMPLE 108:

3-*exo*-(9H-Purin-6-ylamino-methyl)-8-*aza*-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester

Step 1:

(8-Benzyl-8-*aza*-bicyclo[3.2.1]oct-3-*exo*-ylmethyl)carbamic acid *tert*-butyl ester



To a solution of (8-benzyl-8-*aza*-bicyclo[3.2.1]oct-3-*exo*-yl)methylamine (EXAMPLE 107, Step 1) (0.65g, 2.8mmol) in dichloromethane

(30mL) was added di-*tert*-butyl dicarbonate (0.65mL, 0.69g, 3.0mmol). The solution was stirred 18h under nitrogen. The solution was diluted with dichloromethane (50mL), washed with saturated sodium bicarbonate solution (25mL), water (25mL), and brine (25mL), dried (sodium sulfate), filtered, and the solvent was evaporated under reduced pressure to give a crude product (0.993g) as a pale yellow solid. A solution of the crude product in ethyl acetate (5mL) was filtered through a pad of silica gel, eluting with ethyl acetate: hexane (2:1). The filtrate was evaporated under reduced pressure to give (8-benzyl-8-*aza*-bicyclo[3.2.1]oct-3-*exo*-ylmethyl)carbamic acid *tert*-butyl ester product as a white solid.

¹H NMR (CDCl₃) δ 7.37 (2H, d, 7.7 Hz), 7.30 (2H, t, 7.7 Hz), 7.24 (1H, m), 4.55 (1H, br s), 3.53 (2H, s), 3.19 (2H, s), 2.99 (2H, m), 2.00 (2H, m), 1.80 (1H, m), 1.55 (4H, m), 1.44 (11H, m).

Step 2:

(8-Aza-bicyclo[3.2.1]oct-3-*exo*-ylmethyl)carbamic acid *tert*-butyl ester




A mixture of *tert*-butyl (8-benzyl-8-*aza*-bicyclo[3.2.1]oct-3-*exo*-yl)methylcarbamate (0.892g, 2.7mmol) and 10% palladium on carbon (0.55g) in methanol (50mL) was hydrogenated under a hydrogen balloon for 18h. The catalyst was removed by filtration through Celite. The filter cake was washed with methanol (3x25mL) and the filtrate was concentrated under reduced pressure to give crude (8-*aza*-bicyclo[3.2.1]oct-3-*exo*-ylmethyl)carbamic acid *tert*-butyl ester product as a white solid.

¹H NMR (CDCl₃) δ 4.57 (1H, br s), 3.53 (2H, s), 2.96 (2H, m), 1.95-1.77 (4H, m), 1.72-1.50 (4H, m), 1.44 (9H, m), 1.24 (2H, m).

Mass spec.: 241.32 (M+1).

Step 3:

3-*exo*-(*tert*-Butoxycarbonylamino-methyl)-8-*aza*-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester



To a mixture of *tert*-butyl 8-*aza*-bicyclo[3.2.1]oct-3-*exo*-ylmethylcarbamate (0.84g, 3.5mmol) in acetonitrile (35mL) was added 1-[[[(benzyloxy)carbonyloxy]pyrrolidine-2,5-dione (0.87g, 3.5mmol)]. The mixture


was stirred 18h under nitrogen. The resulting solution was concentrated under reduced pressure. The residue was partitioned between ethyl acetate (150mL) and water (75mL) and the layers were separated. The organic layer was washed with

water (2x75mL) and brine (50mL), dried (sodium sulfate), filtered, and the solvent was evaporated under reduced pressure to give a crude product (1.31g) as a white solid. The crude product was purified by flash column chromatography on silica gel, eluting with ethyl acetate: hexane (30:70 increasing to 50:50) to give 3-*exo*-(*tert*-butoxycarbonylamino-methyl)-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester product as a white solid.

¹H NMR (CDCl₃) δ 7.36 (5H, m), 5.13 (2H, s), 4.56 (1H, br s), 4.32 (2H, br s), 2.94 (2H, m), 2.00 (3H, m), 1.62 (4H, m), 1.48-1.25 (11H, m).

Mass spec.: 375.39 (M+1).

10 Step 4: 3-*exo*-Aminomethyl-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester




Benzyl 3-*exo*-[[(*tert*-butoxycarbonyl)amino]methyl]-8-

15 azabicyclo[3.2.1]octane-8-carboxylate (0.94g, 2.5mmol) was placed in a round-bottom flask under nitrogen and cooled in an ice-bath. Trifluoroacetic acid (6mL) was added dropwise and the mixture was stirred one hour with ice-bath cooling. The mixture was poured into ice-cold 5N sodium hydroxide solution (16mL) and the aqueous mixture was extracted with methylene chloride (4x50mL). The extract was washed with brine (50mL), dried (sodium sulfate), filtered, and the solvent was evaporated under reduced pressure to give product as a colorless oil.

¹H NMR (CDCl₃) δ 7.36 (5H, m), 5.14 (2H, s), 4.33 (2H, br s), 2.52 (2H, d, J 6 Hz), 1.96 (2H, m), 1.88 (1H, m), 1.67 (2H, d, J 7 Hz), 1.61 (2H, m), 1.42-1.25 (4H, m).

25 Mass spec.: 275.34 (M+1).

Step 5: 3-*exo*-[(9H-Purin-6-ylamino)-methyl]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester




A solution of 3-*exo*-aminomethyl-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester (27mg, 0.10mmol), 6-chloropurine (31mg, 0.20mmol), and diisopropylethylamine (35μL, 0.20mmol) in isopropanol (2mL) was heated at reflux for 18h. The resulting mixture was concentrated under reduced pressure and the residue was taken up in ethyl acetate (3mL). The resulting mixture was washed with saturated sodium bicarbonate solution (1mL), water (2x1mL), and brine (1mL), dried (sodium sulfate), filtered, and the solvent was evaporated under reduced pressure to give a crude product (39mg) as a yellow solid. The solid was triturated in hot ethyl acetate (1mL), the mixture cooled to ambient temperature, and the solid precipitate filtered off and dried *in vacuo* to give 3-*exo*-[(9H-purin-6-ylamino)-methyl]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester product as a white solid.

¹H NMR (DMSO-d₆) δ 12.86 (1H, br s), 8.16 (1H, s), 8.07 (1H, s), 7.61 (1H, br s), 7.35 (5H, m), 5.08 (2H, d, J 2 Hz), 4.17 (2H, br s), 3.32 (2H, m), 2.26 (1H, m), 1.86 (2H, br s), 1.61 (4H, m), 1.34 (2H, m).

Mass spec.: 393.36 (M+1).

EXAMPLE 109:

20 3-*exo*-[(3-Chloropurin-2-ylamino)methyl]-8-azabicyclo[3.2.1]octane-8-carboxylic acid benzyl ester



Employing the procedure substantially as described for 3-*exo*-(9H-purin-6-ylamino)-methyl]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester (EXAMPLE 108), but substituting 2,3-dichloropyrazine for 6-chloropurine, the crude

product (51mg) was obtained as an oil. The crude product was filtered through a pad of silica gel eluting with ethyl acetate: hexane (2:1), and the filtrate was concentrated under reduced pressure. The residual oil was dissolved in ether, the solvent evaporated under reduced pressure, and the residue dried in vacuo to give 3-*exo*-[(3-chloropropyl-2-ylamino)methyl]-8-aza-bicyclo[3.2.1]octane-8-carboxylic acid benzyl ester as a yellow gum.

¹H NMR (CDCl₃) δ 7.93 (1H, d, J 3 Hz), 7.56 (1H, d, J 3 Hz), 7.36 (5H, m), 5.20 (1H, m), 5.15 (2H, s), 4.34 (2H, br s), 3.32 (2H, m), 2.21 (1H, m), 1.97 (2H, m), 1.66 (4H, m), 1.60-1.40 (2H, m).
Mass spec.: 387.27 (M+1).


EXAMPLE 110:

[8-(2-Phenylethanesulfonyl)-8-aza-bicyclo[3.2.1]oct-3-*exo*-

ylmethyl]pyrimidin-2-yl-amine

Step 1:

[8-(2-*trans*-Phenylethanesulfonyl)-8-aza-bicyclo[3.2.1]oct-3-*exo*-ylmethyl]carbamate acid *tert*-butyl ester



To a solution of *tert*-butyl 8-azabicyclo[3.2.1]oct-3-*exo*-


ylmethylcarbamate (EXAMPLE 107, Step 1) (0.60g, 2.5mmol) and diisopropylethylamine (0.52mL, 0.39g, 3.0mmol) in methylene chloride (15mL), under nitrogen cooled in an ice-bath, was added dropwise over 10 minutes a solution of *trans*-2-phenylethanesulfonyl chloride (0.57g, 2.8mmol) in methylene chloride (10mL). The resulting mixture was stirred 18h under nitrogen while warming from ice-bath to ambient temperature. The solution was diluted with dichloromethane (125mL), washed with 1N sodium hydroxide solution (50mL), water (50mL), and brine (50mL), dried (sodium sulfate), filtered, and the solvent was evaporated under reduced pressure to give a crude product (0.95g) as yellow gum. The crude product was purified by flash column chromatography on silica gel, eluting with ethyl acetate: hexane (33:67 increasing to 50:50) to give [8-(2-*trans*-phenylethanesulfonyl)-8-aza-

bicyclo[3.2.1]oct-3-*exo*-ylmethyl]carbamate acid *tert*-butyl ester product as a colorless gum.

¹H NMR (CDCl₃) δ 7.50-7.40 (6H, m), 6.65 (1H, d, J 15 Hz), 4.58 (1H, br s), 4.24 (2H, br s), 3.00 (2H, m), 1.96 (3H, m), 1.69 (3H, m), 1.54 (3H, m), 1.44 (9H, m).

Step 2:

[8-(2-Phenylethanesulfonyl)-8-aza-bicyclo[3.2.1]oct-3-*exo*-ylmethyl]carbamate acid *tert*-butyl ester




A mixture of [8-(2-*trans*-phenylethanesulfonyl)-8-aza-

bicyclo[3.2.1]oct-3-*exo*-ylmethyl]carbamate acid *tert*-butyl ester (0.61g, 1.5mmol) and 20% palladium hydroxide on carbon (0.30g) in ethanol (50mL) was hydrogenated (52psi hydrogen) for 18h. The catalyst was removed by filtration through Celite. The filter cake was washed with ethanol (3x25mL) and the filtrate was concentrated under reduced pressure to give crude [8-(2-phenylethanesulfonyl)-8-aza-bicyclo[3.2.1]oct-3-*exo*-ylmethyl]carbamate acid *tert*-butyl ester product as a gum.

¹H NMR (CDCl₃) δ 7.35-7.20 (5H, m), 4.56 (1H, br s), 4.24 (2H, br s), 3.24 (2H, m), 3.11 (2H, m), 2.98 (2H, t, J 6 Hz), 2.02 (2H, m), 1.92 (1H, m), 1.74-1.51 (4H, m), 1.44 (9H, s), 1.37 (2H, m).

Step 3:

C-[8-(2-Phenylethanesulfonyl)-8-aza-bicyclo[3.2.1]oct-3-*exo*-yl]methylamine



A solution of crude [8-(2-phenylethanesulfonyl)-8-aza-

bicyclo[3.2.1]oct-3-*exo*-ylmethyl]carbamate acid *tert*-butyl ester (0.64g, 1.5mmol) in


dioxane (2mL) and 3N hydrochloric acid (2mL) was heated at reflux for 3h. The solvent was removed under reduced pressure. The aqueous residue was cooled in an ice-bath and made basic with 3N sodium hydroxide solution. The aqueous mixture was extracted with methylene chloride (4x20mL). The organic layer was washed with brine (20mL), dried (sodium sulfate), filtered, and the solvent was evaporated under reduced pressure to give a crude product (0.404g) as a pale yellow oil. A solution of the crude product in methylene chloride was filtered through a pad of silica gel eluting with methanol: methylene chloride: ammonium hydroxide (20:80:2) to give C-[8-(2-phenylethanesulfonyl)-8-aza-bicyclo[3.2.1]oct-3-*exo*-yl]methylamine product as a yellow oil.

¹H NMR (CDCl₃) δ 7.32 (2H, m), 7.26 (1H, m), 7.21 (2H, d, J 7 Hz), 4.24 (2H, m), 3.24 (2H, m), 3.11 (2H, m), 2.56 (2H, d, J 6 Hz), 2.03 (2H, m), 1.82-1.65 (5H, m), 1.37 (4H, m).

Mass spec.: 309.33 (M+1).

Step 4:

[8-(2-Phenylethanesulfonyl)-8-aza-bicyclo[3.2.1]oct-3-*exo*-ylmethyl]pyrimidin-2-yl-amine




A solution of C-[8-(2-phenylethanesulfonyl)-8-aza-bicyclo[3.2.1]oct-3-*exo*-yl]methylamine (31mg, 0.10mmol), 2-bromopyrimidine (32mg, 0.20mmol), and diisopropylethylamine (35μL, 0.20mmol) in isopropanol (2mL) was heated at reflux for 18h. The mixture was concentrated under reduced pressure and the residue was taken up in ethyl acetate (3mL). The resulting mixture was washed with saturated sodium bicarbonate solution (1mL), water (2x1mL), and brine (1mL), dried (sodium sulfate), filtered, and the solvent was evaporated under reduced pressure to give a crude product (39mg) as a yellow solid. The crude product was chromatographed on a 1mm silica gel prep plate eluting with ethyl acetate: hexane (2:1) to give a colorless gum (27mg). The gum was crystallized from ethyl acetate, the precipitate filtered off, and dried *in vacuo* to give [8-(2-phenylethanesulfonyl)-8-aza-bicyclo[3.2.1]oct-3-*exo*-ylmethyl]pyrimidin-2-yl-amine product as a white solid.

¹H NMR (CDCl₃) δ 8.26 (2H, d, J 5 Hz), 7.32 (2H, m), 7.26 (1H, m), 7.21 (2H, d, J 7 Hz), 6.53 (1H, t, J 5 Hz), 5.11 (1H, m), 4.25 (2H, m), 3.31 (2H, t, J 6.5 Hz), 3.24 (2H, m), 3.12 (2H, m), 2.03 (3H, m), 1.74 (4H, m), 1.46 (2H, m).

Mass spec.: 387.31 (M+1).

EXAMPLE 111:


1-[4-(Pyrimidin-2-ylaminomethyl)-piperidin-1-yl]-4-thiophen-2-yl-butan-1-one



Benzyl 4-[(2-pyrimidinylamino)methyl]-1-piperidinecarboxylate (EXAMPLE 16) was hydrogenated as described in EXAMPLE 30, Step 1. The resulting piperidine was combined with EDC (1.3equiv.), HOBT (1.0equiv.), and 4-thiophen-2-yl-butyric acid (1.0equiv.) in DMF and stirred for 2h. The resulting reaction solution was partitioned into ethyl acetate and aqueous sodium bicarbonate. The organic layer was separated and washed with pH 4.5 citric acid buffer (10% citric acid and sodium hydroxide), dried (sodium sulfate), and concentrated to yield the desired 1-[4-(pyrimidin-2-ylaminomethyl)-piperidin-1-yl]-4-thiophen-2-yl-butan-1-one. M.S. (M+1): 345.25

EXAMPLE 112:


3-Phenyl-1-[4-(pyrimidin-2-ylaminomethyl)-piperidin-1-yl]-propan-1-one



The title compound was prepared as described in EXAMPLE 111, except substituting 4-thiophen-2-yl-butyric acid with 3-phenylpropionic acid. M.S. (M+1): 325.28

EXAMPLE 113:


(2-Phenyl-cyclopropyl)-[4-(pyrimidin-2-ylaminomethyl)-piperidin-1-yl]-methanone



The title compound was prepared as described in **EXAMPLE 111**, except substituting 4-thiophen-2-yl-butyric acid with 2-phenyl-cyclopropanecarboxylic acid. M.S. (M+1): 337.27

EXAMPLE 114:


2-Phenoxy-1-[4-(pyrimidin-2-ylaminomethyl)-piperidin-1-yl]-ethanone



The title compound was prepared as described in **EXAMPLE 111**, except substituting 4-thiophen-2-yl-butyric acid with phenoxycetic acid. M.S. (M+1): 341.27

EXAMPLE 115:


4-(Pyridin-4-ylaminomethyl)-piperidine-1-carboxylic acid thiophen-3-ylmethyl ester



The title compound was prepared as described in **EXAMPLE 30**, except substituting 3-fluorobenzyl alcohol with thiophen-3-yl-methanol. M.S. (M+1): 332.31

EXAMPLE 116:


N-benzyl-N'-cyano-N''-[4-(pyridin-4-ylaminomethyl)piperidinyl]guanidine



To a solution of diphenyl cyanocarbonimidate (0.44mmol) in THF (3mL) at -78°C was added benzylamine (0.44mmol, in 2mL THF) dropwise. The cooling bath was removed, and after reaching 20°C, piperidin-4-ylmethyl-pyridin-4-yl-amine (0.44mmol, in 2 mL DMF, **EXAMPLE 30**) was added. The resulting reaction mixture was heated to 90°C for 14h, cooled, the volatiles were removed under vacuum, and the resulting residue purified by silica gel chromatography. M.S. (M+1): 349.38

EXAMPLE 117:


4-[(3-Chloro-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid 4-chloro-benzyl ester



The title compound was prepared as described in **EXAMPLE 47**, reacting 2,3-dichloropyrazine with **INTERMEDIATE 2b** to give the title compound. M.S. (M+1): 395.

EXAMPLE 118:

4-[(3-Chloro-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid 4-methyl-benzyl ester




The title compound was prepared as described in **EXAMPLE 47**, reacting 2,3-dichloropyrazine with **INTERMEDIATE 2a** to give the title compound. M.S.(M+1): 375.

5


EXAMPLE 119:

4-[(3-Chloro-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid indan-2-yl ester



10 Step 1:

4-Aminomethyl-piperidine-1-carboxylic acid indan-2-yl ester



The title compound was prepared in the same way as described for the preparation of **INTERMEDIATES 2A-E**.

15 Step 2:


4-[(3-Chloro-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid indan-2-yl ester

The title compound was prepared as described in **EXAMPLE 47**, reacting 2,3-dichloropyrazine with the amine described in **STEP 1**. M.S.(M+1): 387.

20


EXAMPLE 120:

4-[(3-Chloro-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzylamide



5 Step 1:

4-Aminomethyl-piperidine-1-carboxylic acid benzylamide



The title compound was prepared in the same way as described for the preparation of **INTERMEDIATES 2A-E**, replacing the **INTERMEDIATE 1A-E** with benzyl isocyanate

10

Step 2:

4-[(3-Chloro-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzylamide


The title compound was prepared as described in **EXAMPLE 47**, reacting 2,3-dichloropyrazine with the amine described in **STEP 1**, to give the title compound. M.S.(M+1): 360

15

EXAMPLE 121:

4-[(3-Cyano-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester


20



The title compound was prepared in a manner similar to that described for the preparation of **EXAMPLE 47**, utilizing 2-chloro-3-cyanopyrazine (Maybridge Chemicals) in place of 2,3-dichloropyrazine. M.S.(M+1): 352.

EXAMPLE 122:


4-[(3-Aminomethyl-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester trifluoroacetic acid salt



To a solution of 4-[(3-cyano-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester (130mg) (**EXAMPLE 121**) in ethanol (10mL) under nitrogen, was added Raney Nickel (20mg) and the mixture stirred under hydrogen (1atm) for 8h. The reaction was filtered, concentrated in vacuo, and then purified using reverse phase chromatography C-18 (gradient elution 0.1% aqueous trifluoroacetic acid/acetonitrile) to give the title compound as the trifluoroacetic acid salt. M.S.(M+1): 356.

EXAMPLE 123:


4-[(6-Aminomethyl-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester trifluoroacetic acid salt



This was prepared in a manner similar to that described for the preparation of **EXAMPLE 122**, from 2-chloro-6-cyanopyrazine (L. Bernadi et al *Gazz. Chim. Ital.*, 91, 1431 (1961) and benzyl 4-(aminomethyl)piperidine-1-carboxylate (**EXAMPLE 13**, Step 1). M.S.(M+1): 356

EXAMPLE 124:


4-[(3-Methoxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



4-[(3-Chloro-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester (2.72g, 7.54mmol) and 0.5 M sodium methoxide in methanol (40mL) were heated under nitrogen at 60°C for 2 days, cooled, evaporated and the residue partitioned between EtOAc and water. The organic layer was washed with brine, dried and solvent evaporated to afford crude material, purified by flash chromatography on silica (gradient 25 to 100%EtOAc hexane) to give the desired compound as a solid. The solid was stirred with approx. (10mL) 2:1 isopropyl acetate: hexane and filtered to give the title compound as white solid. M.S.(M+1): 357

EXAMPLE 125:

4-[(3-Ethoxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid
benzyl ester




The title compound was prepared as described for **EXAMPLE 124**,
using sodium ethoxide in ethanol in place of sodium methoxide in methanol.

M.S.(M+1): 371

EXAMPLE 126:

4-[(3-isopropoxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic
acid benzyl ester



The title compound was prepared as described for **EXAMPLE 124**,
using sodium isopropoxide in isopropanol in place of sodium methoxide in methanol.


M.S.(M+1): 385

EXAMPLE 127:

(4-[(3-Methoxy-pyrazin-2-ylamino)-methyl]-piperidin-1-yl)-(1R,2R)-
2-phenyl-cyclopropyl)-methanone

Step 1:

4-[(3-Chloro-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid
tert-butyl ester




5

2,3-Dichloropyrazine (1.0g, 0.0067mol), tert-butyl-4-(aminomethyl)piperidine-1-carboxylate (1.6g, 0.0075mol) (Astatech) and cesium carbonate (2.4g, 0.0075mol) in acetonitrile (10 mL) were heated to 90°C under nitrogen for 18h. The reaction was concentrated in vacuo, diluted with ethyl acetate (50mL) and washed with water (50mL). The organic extract was dried over sodium sulfate, filtered and chromatographed on silica using a gradient of 10 to 30% ethyl acetate/hexane to give the title compound as a foam. M.S.(M+1): 327

Step 2:

4-[(3-Methoxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic
acid tert-butyl ester




4-[(3-Chloro-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid
tert-butyl ester (3.0g, 0.0092mol) and 0.5M sodium methoxide in methanol (40mL)
were heated under nitrogen at 75°C for 18h. The reaction was concentrated in vacuo,
diluted with methylene chloride (100mL) and washed with water (pH=9, adjusted

20

with NaOH). The organic extract was dried over sodium sulfate filtered and concentrated to give the title compound. M.S. (M+1): 323.

Step 3:


4-[(3-Methoxy-pyrazin-2-ylamino)-methyl]-piperidine



4-[(3-Methoxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester (0.5g, 0.0015mol) and trifluoroacetic acid (5mL) were allowed to stir under nitrogen for 0.5h. The reaction was concentrated in vacuo, and chromatographed on silica using methylene chloride/methanol/ammonium hydroxide (90/10/2) to give the title compound. M.S. (M+1): 223.

Step 4:


4-[(3-Methoxy-pyrazin-2-ylamino)-methyl]-piperidin-1-yl)-(1R,2R)-2-phenyl-cyclopropyl)-methanone



A mixture of 4-[(3-methoxy-pyrazin-2-ylamino)-methyl]-piperidine (0.093g, 0.00042mol), 1-hydroxybenzotriazole (0.078g, 0.0005mol), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (0.097g, 0.0005mole) and (1R,2R)-2-phenylcyclopropanecarboxylic acid (T. Riley et al., *J. Med. Chem.*, 15, 1187, 1972) (0.072g, 0.00044mol) in DMF (2mL) were stirred at rt for 18h. The reaction was diluted with ethyl acetate (30mL), washed with 10% aqueous sodium bicarbonate (20mL) followed by brine (10mL), concentrated in vacuo and chromatographed on silica using 50-100% ethyl acetate/hexane. Crystallization from ether/hexane gave the title compound. M.S. (M+1): 367.

EXAMPLE 128:

[2-((1R,2R)-(2-Fluoro-phenyl))-cyclopropyl]-(4-[(3-methoxy-pyrazin-2-ylamino)-methyl]-piperidin-1-yl)-methanone




The title compound was prepared in a manner similar to that described for the preparation of EXAMPLE 127, Step 4 using (1R,2R)-2-(2-fluorophenyl)cyclopropanecarboxylic acid, prepared as described below:

5 M.S. (M+1): 385.

Step 1:

(R,R)-2-(2-Fluoro-phenyl)-cyclopropanecarboxylic acid tert-butyl ester




To a solution of copper triflate (2:1 benzene complex) (21mg,

0.041mmol) in chloroform (20mL) under nitrogen was added 2,2'-isopropylidenebis-(4S)-4-tert-butyl-2-oxazoline (12.5mg, 0.042mmol) and the mixture allowed to stir at rt for 1h. The reaction was filtered under nitrogen into a flask and 2-fluorostyrene (1.0gm, 8.19mmole) added. A solution of t-butyl diazoacetate (0.63mL, 4.09mmole) in chloroform (10mL) was added dropwise over 1.5h and the mixture allowed to stir overnight at rt. The reaction was concentrated in vacuo and chromatographed on silica using 3-10% ethyl acetate/hexane to give (R,R)-trans of the title compound as an oil.

¹H NMR 400 MHz (δ, CDCl₃) δ: 1.22(m, 1H), 1.48(s, 9H), 1.54(m, 1H), 1.84(m, 1H), 2.58(m, 1H), 6.9-7.1(m, 3H), 7.17(m, 1H).

Step 2:

2-(2-Fluoro-phenyl)-cyclopropanecarboxylic acid




To the t-butyl ester from Step 1 (0.52g, 0.0022mole) in dichloromethane at 0°C was added trifluoroacetic acid and the mixture stirred at rt for 30min. The reaction was concentrated in vacuo to give the title compound as an oil.

Analysis of the acid by chiral HPLC (Chirepak AD, 250X4.6 mm) using 95/5(A/B), 0.2% trifluoroacetic acid in hexane(A) and ethanol(B), 1mL/min, showed the material to have a purity of $\geq 94\%$ EE. M.S.(M+1): 181.

5 EXAMPLE 129:

[2-((1R,2R)-2,6-Difluoro-phenyl)-cyclopropyl]-4-[(3-methoxy-pyrazin-2-ylamino)-methyl]-piperidin-1-yl)-methanone




The title compound was prepared in a manner similar to that described for the preparation of EXAMPLE 127, Step 4 using (1R,2R)-2-(2,6-

difluorophenyl)cyclopropanecarboxylic acid (prepared in a similar manner to that described for 2-(2-fluoro-phenyl)-cyclopropanecarboxylic acid (EXAMPLE 128). M.S.(M+1): 403.

15 EXAMPLE 130:


4-[(3-Methoxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid 4-methyl-benzyl ester



A mixture of 4-[(3-methoxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid (EXAMPLE 127, STEP 3) (0.093g, 0.00042mol) and N-(4-methylbenzyl)oxycarbonyloxysuccinimide (INTERMEDIATE 1A) (118mg) in DMF (2mL) was stirred at rt for 18h. The reaction was diluted with ethyl acetate (30mL), washed with 10 % aqueous sodium bicarbonate (20mL) followed by brine (10mL), concentrated in vacuo and chromatographed on silica using a gradient elution of 5-15% acetone/dichloromethane. Concentration in vacuo followed by crystallization from ether/hexane gave the title compound. M.S.(M+1): 371.

EXAMPLE 131:


4-[(5-Cyano-3-methoxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



5

Step 1:

4-[(5-Bromo-3-methoxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester




10

To 4-[(3-methoxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester (EXAMPLE 127, STEP 2) (2.0g, 0.0062mol) in chloroform (160mL) and under nitrogen was added pyridine (0.528mL, 0.0064mol), followed by a slow addition (~1h) of a solution of bromine (1.044g, 0.0064mol) in chloroform (16mL). The reaction was diluted with water (100mL) and the organic layer removed, dried over sodium sulfate, filtered and concentrated to an oil. The oil was chromatographed on silica using a gradient of 0 to 4% acetone/dichloromethane to give the title compound as a foam. M.S.(M+1): 401.

Step 2:

4-[(5-Cyano-3-methoxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester


20



To 4-[(5-bromo-3-methoxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester (0.5g, 0.00125mol) in DMSO (10mL), under nitrogen, was added copper cyanide (0.56g, 0.00625mol) and the mixture heated to 150°C for 1.5h. The reaction mixture was cooled to rt, diluted with a mixture of 20% ammonium hydroxide in water (50mL) and dichloromethane (50mL) and allowed to stir for 1h. The organic layer was removed, dried over sodium sulfate, filtered and concentrated to an oil. The oil was chromatographed on silica using a gradient of 20-40 % ethyl acetate/hexane to give the title compound as a foam. M.S.(M+1): 348.

10 Step 3:

4-[(5-Cyano-3-methoxy-pyrazin-2-ylamino)-methyl]-piperidine



The above compound was prepared in a similar manner as described in EXAMPLE 127, STEP 3 from 4-[(5-cyano-3-methoxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester. M.S.(M+1): 248.


15 Step 4:

4-[(5-Cyano-3-methoxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester

The above compound was prepared in a similar manner as described in EXAMPLE 130 from 4-[(5-cyano-3-methoxy-pyrazin-2-ylamino)-methyl]-piperidine using N-(benzyloxycarbonyloxy)succinimide (Sigma-Aldrich). M.S.(M+1): 382.

EXAMPLE 132:


6-Methoxy-5-[(1-(2-(1R,2R)-phenyl-cyclopropanecarbonyl)-piperidin-4-yl)methyl]-amino)-pyrazine-2-carbonitrile



The above compound was prepared in a similar manner as described in EXAMPLE 127, STEP 4 from 4-[(5-cyano-3-methoxy-pyrazin-2-ylamino)-methyl]-piperidine (EXAMPLE 131, STEP 3). M.S.(M+1): 392.


EXAMPLE 133:

4-[(3-Methoxy-5-methyl-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



Step 1:


4-[(3-Methoxy-5-methyl-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester



To 4-[(5-bromo-3-methoxy-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester (EXAMPLE 131, STEP 1) (0.20g, 0.0005mmol) in tetrahydrofuran (1mL), under nitrogen, was added 1,3-bis(diphenylphosphino)propane nickel(II) chloride (0.034g, 0.00625mmol) followed by a dropwise addition of 2.0M dimethylzinc in toluene (0.313mL, 0.000625mmol). The reaction mixture was stirred for 1.5h, diluted with water (5mL) and ethyl acetate (30mL). The organic layer was removed, dried over sodium sulfate, filtered and concentrated to an oil. The oil was chromatographed on silica using a gradient of 20-50 % ethyl acetate/hexane to give the title compound as a foam. M.S.(M+1): 337.

Step 2


4-[(3-Methoxy-5-methyl-pyrazin-2-ylamino)-methyl]-piperidine



The above compound was prepared in a similar manner as described in EXAMPLE 127, STEP 3 from 4-[(3-methoxy-5-methyl-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid tert-butyl ester. M.S.(M+1): 237

Step 3:


4-[(3-Methoxy-5-methyl-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



The above compound was prepared in a similar manner as described in EXAMPLE 130 from 4-[(3-methoxy-5-methyl-pyrazin-2-ylamino)-methyl]-piperidine using N-(benzyloxycarbonyloxy)succinimide (Sigma-Aldrich). M.S.(M+1): 371.

EXAMPLE 134:

4-[(3-Methoxy-5-methyl-pyrazin-2-ylamino)-methyl]-piperidine-1-carboxylic acid 4-methyl-benzyl ester




The above compound was prepared in a similar manner as described in EXAMPLE 130 from 4-[(3-methoxy-5-methyl-pyrazin-2-ylamino)-methyl]-piperidine (EXAMPLE 130, STEP 2) using N-(4-methylbenzyloxycarbonyloxy)succinimide (INTERMEDIATE 1A). M.S.(M+1):

385.

EXAMPLE 135:

4-[(3-Methoxy-5-methyl-pyrazin-2-ylamino)-methyl]-piperidine-1-yl)-(2-((1R,2R)-phenyl)-cyclopropyl)-methanone



15


The above compound was prepared in a similar manner as described in EXAMPLE 127, STEP 4, from 4-[(3-methoxy-5-methyl-pyrazin-2-ylamino)-methyl]-piperidine (EXAMPLE 133, STEP 2). M.S.(M+1): 381.

EXAMPLE 136:

trans N-[(1-[(2-(2-Fluorophenyl)cyclopropyl]carbonyl]piperidin-4-yl)methyl]-5-fluoropyrimidin-2-amine

Step 1:


Methyl (2E)-3-(2-fluorophenyl)prop-2-enoate



5 HCl gas was bubbled through a stirring solution of 2-fluorocinnamic acid in anhydrous methanol. The reaction mixture was allowed to cool to room temperature, then concentrated to yield the title compound. M.S. (M+1): 181.

Step 2:


Methyl 2-(2-fluorophenyl)cyclopropanecarboxylate



10 Diazomethane was prepared as follows: To a stirring solution of ether (290mL) and 40% KOH (aq, 90mL) at 0°C, was added 1-methyl-3-nitro-1-nitrosoguanidine (24.42g, 166.53mmol), portionwise. After stirring for 1h, the mixture was cooled to -78°C and allowed to stir for an additional ten minutes. The ether layer and palladium acetate (approx 200mg) were then both added in approx. 10 portions to a stirred solution of methyl (2E)-3-(2-fluorophenyl)prop-2-enoate (3.0g, 16.65mmol) in ether (20mL) at 0°C. After stirring at rt for approximately thirty minutes, the reaction mixture was then filtered through silica gel and concentrated. M.S. (M+1): 195.


Step 3:

Preparation of 2-(2-fluorophenyl)cyclopropanecarboxylic acid



To a stirred solution of methyl 2-(2-fluorophenyl)cyclopropanecarboxylate (4.8g, 24.72mmol) in tetrahydrofuran (25mL), was added 10M sodium hydroxide solution (approximately 2mL), a small amount of water, and sufficient methanol to achieve a homogeneous reaction mixture. The reaction mixture was then allowed to stir at rt for approximately 2h. After concentrating the reaction mixture, 1N HCl was added until the mixture was acidic. The organic layer was extracted twice with ethyl acetate, then washed with brine, dried over anhydrous Na₂SO₄, and concentrated to afford the title compound. M.S. (M+1): 181.

Step 4:




N-((1-((2-(2-fluorophenyl)cyclopropyl)carbonyl)piperidin-4-yl)methyl)-5-fluoropyrimidin-2-amine

A solution of 4-[(5-fluoro-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester (EXAMPLE 71, STEP 2) was hydrogenated at 1atm. of hydrogen over 10% Pd/C in ethanol until debenzoylation was complete. The reaction mixture was then filtered, the catalyst washed with ethanol and solvent evaporated to give the deprotected amine which was coupled with 2-(2-fluorophenyl)cyclopropanecarboxylic acid using the conditions described in EXAMPLE 127, STEP 4 to give the title compound after chromatography on silica. ¹H NMR (400 MHz): δ 8.15 (m, 2H); 7.17 (brs, 1H); 7.04 (m, 2H); 5.47 (brs, 1H); 4.65 (brs, 1H); 4.15 (d, 1H); 3.31 (m, 2H); 3.08 (t, 1H); 2.56 (m, 2H); 2.02 (brs, 1H); 1.90 (m, 3H); 1.67 (m, 1H); 1.22 (m, 4H). M.S. (M+1): 373.

EXAMPLE 137:


(S,S) and (R,R) N-((1-((2-(2-fluorophenyl)cyclopropyl)carbonyl)piperidin-4-yl)methyl)-5-fluoropyrimidin-2-amine



Separation of the two enantiomers of N-((1-([2-(2,6-difluorophenyl)cyclopropyl]carbonyl)piperidin-4-yl)methyl)-5-fluoropyrimidin-2-amine was accomplished on a Chiralpak AD column, eluting with 0.1% diethylamine in hexane/2-propanol.

EXAMPLE 138:

N-((1-([2-(2,6-difluorophenyl)cyclopropyl]carbonyl)piperidin-4-yl)methyl)-5-fluoropyrimidin-2-amine




The title compound was prepared in a manner similar to that described for EXAMPLE 136, starting with 2,6-difluorocinnamic acid.

¹H NMR (400 MHz, CDCl₃): δ 8.16 (s, 2H); 7.13 (brs, 1H); 6.82 (dd, 2H); 5.20 (s, 1H); 4.67 (m, 1H); 4.23 (m, 1H); 3.32 (dd, 2H); 3.11 (m, 1H); 2.64 (m, 1H); 2.38 (m, 1H); 2.30 (m, 1H); 1.94-1.80 (m, 2H); 1.66 (m, 2H); 1.40-1.39 (m, 1H); 1.27-1.22 (m, 2H).

M.S. (M+1): 391.

EXAMPLE 139:


(S,S) and (R,R) N-((1-([2-(2,6-difluorophenyl)cyclopropyl]carbonyl)piperidin-4-yl)methyl)-5-fluoropyrimidin-2-amine



Separation of the two enantiomers of N-((1-([2-(2-fluorophenyl)cyclopropyl]carbonyl)piperidin-4-yl)methyl)-5-fluoropyrimidin-2-amine was accomplished on a Chiralpak AD column, eluting with 0.1% diethylamine in hexane/2-propanol.

EXAMPLE 140:

N-((1-([2-(2,3-difluorophenyl)cyclopropyl]carbonyl)piperidin-4-yl)methyl)-5-fluoropyrimidin-2-amine




The title compound was prepared in a manner similar to that described for EXAMPLE 136 using 2,3-difluorocinnamic acid.

¹H NMR (400 MHz): δ 8.15 (s, 2H); 7.00 (m, 2H); 6.77 (brs, 1H); 5.44 (brs, 1H); 4.65 (brs, 1H); 4.14 (d, 1H); 3.09 (t, 1H); 2.59 (m, 2H); 2.05 (brs, 1H); 1.89 (m, 3H); 1.69 (brs, 1H); 1.26 (m, 3H).

M.S. (M+1): 391.

EXAMPLE 141:


(S,S) and (R,R) N-((1-([2-(2,3-difluorophenyl)cyclopropyl]carbonyl)piperidin-4-yl)methyl)-5-fluoropyrimidin-2-amine



Separation of the two enantiomers of N-((1-([2-(2,3-difluorophenyl)cyclopropyl]carbonyl)piperidin-4-yl)methyl)-5-fluoropyrimidin-2-amine was accomplished on a Chiralpak AD column, eluting with 0.1% diethylamine in hexane/2-propanol.


EXAMPLE 142:

Benzyl 4-[[[(5-fluoro-pyrimidin-2-yl)amino]methyl]piperidin-1-carboxylate



Step 1:

2,4-dichloro-5-fluoropyrimidine



A stirred solution of 5-fluorouracil (15.0g, 0.115mol), N,N-dimethylaniline (7.31mL, 0.058mol) in POCl_3 (107mL) was heated to reflux for 1h. The reaction mixture was concentrated in vacuo and the residue quenched with ice (100g) at 0°C. The solution was then extracted with ethyl ether (3 x 200mL). The combined ether layer was washed with aqueous saturated NaHCO_3 (100mL), water (100mL), brine (50mL), dried over Na_2SO_4 , filtered and concentrated to give the title compound.

^1H NMR (400 MHz, CDCl_3): δ 8.49 (s, 1 H, Ar).

Step 2:


2-Chloro-5-fluoropyrimidine (7)



To a stirred, refluxing mixture of 2,4-dichloro-5-fluoropyrimidine (17.0g, 0.102mol) and zinc (100mesh, 20.0g, 0.305mol) in THF (100mL) was slowly added acetic acid (5.8mL, 0.102mol). The resulting reaction mixture was refluxed for 3h, then cooled to RT. Solids were removed by filtration and the filtrate concentrated in vacuo. The residue was chromatographed on silica gel 60 (200g), eluting with 10-50% ethyl acetate in hexane to give the title compound. ^1H NMR (400 MHz, CDCl_3): δ 8.53 (s, 2 H, Ar).

Step 3:

Benzyl 4-[[[(5-fluoro-pyrimidin-2-yl)amino]methyl]piperidin-1-carboxylate




A stirred mixture of benzyl 4-(aminomethyl)piperidin-1-carboxylate (EXAMPLE 13, STEP 1) (10.0g, 0.040mol), 2-chloro-5-fluoropyrimidine (5.3g, 0.040mol) and cesium carbonate (26.2g, 0.081mol) in DMF (100mL) was heated at 100°C for 2h. The reaction mixture was cooled to rt, diluted with ethyl acetate (400mL), washed with aqueous saturated NaHCO_3 (100mL), water (5 x 100mL), and brine (50mL), dried over anhydrous Na_2SO_4 , filtered and concentrated. The residue was chromatographed on silica gel 60 (0.5kg), eluting with 50-100% ethyl acetate in hexane to give the title compound. M.S. ($M+1$): 345.

^1H NMR (400 MHz, CDCl_3): δ 8.15 (s, 2 H, Pyr), 7.35 (m, 5 H, Ar), 5.12 (s, 2 H, ArCH_2), 4.21 (brs, 2 H, NCH_2), 3.29 (t, $J = 6.4$ Hz, 2 H, NHCH_2), 2.78 (brs, 2 H, NCH_2), 1.80 (m, 2 H, CH), 1.77 (m, 2 H, CHCH_2CH_3), 1.20 (m, 2 H, CHCH_2CH_3).


EXAMPLE 143:

5-Fluoro-2-[[[(1-[[[(1*R*,2*R*)-2-phenylcyclopropyl]carbonyl]piperidin-4-yl)methyl]amino]pyrimidine



Step 1:

5-Fluoro-*N*-(piperidin-4-yl)methylpyrimidin-2-amine




A mixture of benzyl 4-(((5-fluoropyrimidin-2-

yl)amino)methyl]piperidine-1-carboxylate (EXAMPLE 142, STEP 3) (9.0g, 0.026mol) and Pd/C (10%, 0.9g) in anhydrous methanol (250mL) was vigorously stirred under hydrogen atmosphere provided by a hydrogen balloon for 2h. The reaction mixture was filtered and the filtrate was concentrated to give the title compound. M.S. (M+1): 211.

¹H NMR (400 MHz, CDCl₃): δ 8.15 (s, 2 H, Pyr), 5.19 (s, 1 H, NH), 3.27 (t, J = 6.3 Hz, 2 H, NHC₂H₅), 3.11 (d, J = 9.1 Hz, 2 H, NHC₂H₅), 2.61 (t, J = 12.1 Hz, 2 H, NHC₂H₅), 1.77 (d, J = 12.7 Hz, 2 H, CH₂CH₂CH₃), 1.73 (m, 1 H, CH), 1.24 (m, 2 H, CH₂CH₂CH₃).

Step 2:

5-Fluoro-2-(((1-(((1*R*,2*R*)-2-phenylcyclopropyl)carbonyl)piperidin-4-yl)methyl)amino)pyrimidine




A solution of 5-fluoro-*N*-(piperidin-4-yl)methyl)pyrimidin-2-amine (1.00g, 4.76mmol), (1*R*,2*R*)-2-phenylcyclopropanecarboxylic acid (T. Riley et al., *J. Med. Chem.*, 15, 1187, 1972) (0.77g, 4.76mmol), EDC (1.37g, 7.13mmol) and HOBT (0.96g, 7.13mmol) in DMF (10mL) was stirred at rt for 2h. The reaction mixture was diluted with ethyl acetate (200mL), washed with aqueous saturated NaHCO₃ (50mL), water (5 x 50mL), brine (20mL), dried over Na₂SO₄, filtered and concentrated. The residue was chromatographed on silica gel 60 (90g), eluting with 10:1-15:89-75 CH₂Cl₂:2-propanol:hexane to give the title compound. M.S. (M+1): 355.

¹H NMR (400 MHz, CDCl₃): δ 8.15 (s, 2 H, Pyr), 7.28 (t, J = 7.6 Hz, 2 H, Ar), 7.19 (t, J = 6.6 Hz, 1 H, Ar), 7.10 (d, J = 7.4 Hz, 2 H, Ar), 5.15 (s, 1 H, NH), 4.64 (d, J = 13.5 Hz, 1 H, NCH₂), 4.14 (d, J = 12.7 Hz, 1 H, NCH₂), 3.30 (s, 2 H, CH₂NH), 3.06 (q, J = 12.8 Hz, 1 H, NCH₂), 2.62 (t, J = 12.1 Hz, 1 H, NCH₂), 2.46 (brs, 1 H, ArCH), 1.98 (m, 1 H, CHCO), 1.87 (m, 1 H, CH₂CHCH₂), 1.82 (m, 2 H, CH₂CH₂CH₃), 1.65 (m, 1 H, CHCH₂CH), 1.26 (m, 1 H, CHCH₂CH), 1.21 (m, 2 H, CH₂CH₂CH₃).


EXAMPLE 144:

Benzyl 4-(((5-methylpyrimidin-2-yl)amino)methyl)piperidine-1-carboxylate



Step 1:

2-chloro-5-methylpyrimidine




To a stirred, refluxing mixture of 2,4-dichloro-5-methylpyrimidine ([1780-31-0](Sigma-Aldrich) (40.0g, 0.245mol) and zinc (100mesh, 48.1g, 0.736mol) in THF (250mL) was slowly added acetic acid (14.0mL, 0.245mol). The resulting reaction mixture was refluxed for 3h, then cooled to RT. The solids were removed by filtration and the filtrate concentrated in vacuo. The residue was chromatographed on silica gel 60 (0.5kg), eluting with a gradient of 10-50% ethyl acetate in hexane to give the title compound. M.S. (M+1): 129.

¹H NMR (400 MHz, CDCl₃): δ 8.47 (s, 2 H, Ar), 2.32 (s, 3 H, CH₃).

Step 2:

Benzyl 4-(((5-methylpyrimidin-2-yl)amino)methyl)piperidine-1-carboxylate




A stirred mixture of benzyl 4-(aminomethyl)piperidine-1-carboxylate (EXAMPLE 13, STEP 1)(20.0g, 0.081mol), 2-chloro-5-methylpyrimidine (10.4g, 0.081mol) and cesium carbonate (52.5g, 0.161mol) in DMF (200mL) was heated at 150°C for 6h. The reaction mixture was cooled to rt, diluted with ethyl acetate (700mL), washed with aqueous saturated NaHCO₃ (200mL), water (5 x 200mL), and brine (100mL), dried over anhydrous sodium sulfate, filtered and concentrated. The

residue was chromatographed on silica gel 60 (1kg), eluting with 50-100% ethyl acetate in hexane to give the title compound. M.S. (M+1): 341.

¹H NMR (400 MHz, CDCl₃): δ 8.11 (s, 2 H, Pyr), 7.35 (m, 5 H, Ar), 5.12 (s, 2 H, ArCH₂), 5.00 (s, 1 H, NH), 4.20 (brs, 2 H, NCH₂), 3.31 (t, J = 6.3 Hz, 2 H, NCH₂), 2.77 (brs, 2 H, NCH₂), 2.12 (s, 3 H, CH₃), 1.78 (m, 1 H, CH), 1.77 (m, 2 H, CHCH₂CH₃), 1.20 (m, 2 H, CHCH₂CH₃).


EXAMPLE 145:

5-Methyl-2-(((1-((1R,2R)-2-phenylcyclopropyl)carbonyl)piperidin-4-yl)methyl)amino]pyrimidine



Step 1:

5-Methyl-N-(piperidin-4-ylmethyl)pyrimidin-2-amine




15 A mixture of benzyl 4-(((5-methylpyrimidin-2-

yl)amino)methyl)piperidine-1-carboxylate (EXAMPLE 144) (13.0g, 0.038mol) and Pd/C (10%, 1.3g) in anhydrous methanol (500mL) was vigorously stirred under a hydrogen atmosphere provided by a hydrogen balloon for 6h. The reaction mixture was filtered and the filtrate was concentrated to give 5-methyl-N-(piperidin-4-yl)methyl)pyrimidin-2-amine. M.S. (M+1): 207.

¹H NMR (400 MHz, CDCl₃): δ 8.11 (s, 2 H), 5.10 (s, 1 H), 3.30 (t, J = 6.4 Hz, 2 H), 3.20 (d, J = 12.3 Hz, 2 H), 2.65 (dt, J = 12.3 & 2.6 Hz, 2 H), 2.12 (s, 3 H), 1.82 (d, J = 13.5 Hz, 2 H), 1.77 (m, 1 H), 1.31 (q, J = 12.1 & 3.7 Hz, 2 H).

Step 2:

25 5-Methyl-2-(((1-((1R,2R)-2-phenylcyclopropyl)carbonyl)piperidin-4-yl)methyl)amino]pyrimidine




A solution of 5-methyl-N-(piperidin-4-ylmethyl)pyrimidin-2-amine (5.00g, 0.024mol), ((1R,2R)-2-phenylcyclopropanecarboxylic acid (T. Riley et al., *J. Med. Chem.*, (1972), 15, 1187) (3.93g, 0.024mol), EDC (6.97g, 0.036mol) and HOBT (4.91g, 0.036mol) in DMF (50mL) was stirred at RT for 2h. The reaction mixture was diluted with ethyl acetate (400mL), washed with aqueous saturated NaHCO₃ (100mL), water (5 x 100mL), brine (50mL), dried over anhydrous sodium sulfate, filtered and concentrated. The residue was chromatographed on silica gel 60 (400g), eluting with 50-100% ethyl acetate in hexane to give 5-methyl-2-(((1-((1R,2R)-2-phenylcyclopropyl)carbonyl)piperidin-4-yl)methyl)amino]pyrimidine. M.S. (M+1): 351.

¹H NMR (400 MHz, CDCl₃): δ 8.11 (s, 2 H), 7.28 (t, J = 7.0 Hz, 2 H), 7.19 (t, J = 7.2 Hz, 1 H), 7.11 (d, J = 7.6 Hz, 2 H), 5.01 (s, 1 H), 4.63 (d, J = 12.1 Hz, 1 H), 4.13 (d, J = 13.2 Hz, 1 H), 3.31 (s, 2 H), 3.05 (q, J = 12.2 Hz, 1 H), 2.62 (t, J = 12.5 Hz, 1 H), 2.46 (brs, 1 H), 2.12 (s, 3 H), 1.97 (s, 1 H), 1.86 (m, 1 H), 1.81 (d, J = 13.3 Hz, 2 H), 1.64 (s, 1 H), 1.26 (s, 1 H), 1.22 (m, 2 H).

EXAMPLE 145A:

5-Methyl-2-(((1-((1R,2R)-2-phenylcyclopropyl)carbonyl)piperidin-4-yl)methyl)amino]pyrimidinium chloride




5-Methyl-2-(((1-((1R,2R)-2-phenylcyclopropyl)carbonyl)piperidin-4-yl)methyl)amino]pyrimidine (6.81g, 19.4mmol) (EXAMPLE 145) was dissolved in EtOH (400mL) and 1M HCl in ether (19.4mL, 19.4 mmol) added. The solution was then concentrated and the residue was crystallized from 30% 2-propanol in ether (100mL) to give the title compound. Melting Point 157.5 °C. M.S. (M+1): 351.


¹H NMR (500 MHz, CD₃OD): δ 8.42 (s, 2 H), 7.25 (m, 2 H), 7.17 (m, 1 H), 7.14 (m, 2 H), 4.55 (d, *J* = 12.9 Hz, 1 H), 4.26 (m, 1 H), 3.38 (m, 2 H), 3.14 (q, *J* = 12.9 Hz, 1 H), 2.69 (t, *J* = 12.1 Hz, 1 H), 2.33 (m, 1 H), 2.24 (s, 3 H), 2.19 (brs, 1 H), 1.97 (m, 1 H), 1.82 (d, *J* = 12.9 Hz, 2 H), 1.53 (m, 1 H), 1.29 (m, 1 H), 1.17 (m, 2 H).

EXAMPLE 146:

5-Methyl-N-((1-((2-(5-methylthien-2-yl)cyclopropyl)carbonyl)piperidin-4-yl)methyl)pyrimidin-2-amine

**Step 1:**


tert-Butyl (2E)-3-(5-methylthien-2-yl)prop-2-enoate



To a solution of *tert*-butyl diethylphosphonoacetate (1.12mL, 4.76mmol) in THF (5mL) at -78 °C was added LHMDS (1.0M in THF, 4.76mL, 4.76mmol). After 5min at -78 °C 5-methyl-2-thiophene-carboxaldehyde (0.43mL, 3.96mmol) was added. The reaction mixture was warmed to RT, stirred for 10min and poured onto EtOAc/H₂O. The layers were separated and the organic layer was washed with H₂O, dried over Na₂SO₄, filtered and concentrated. The residue was chromatographed on silica (gradient elution; hexanes to 4:1 hexanes:EtOAc) to give *tert*-butyl (2E)-3-(5-methylthien-2-yl)prop-2-enoate as a clear oil.

Step 2:

tert-Butyl 2-(5-methylthien-2-yl)cyclopropanecarboxylate



tert-Butyl (2E)-3-(5-methylthien-2-yl)prop-2-enoate was cyclopropanated according to the procedure described for **EXAMPLE 136, STEP 2**, providing, after chromatography, *tert*-butyl 2-(5-methylthien-2-yl)cyclopropanecarboxylate.

Step 3:

2-(5-Methylthien-2-yl)cyclopropanecarboxylic acid




To a solution of *tert*-butyl 2-(5-methylthien-2-yl)cyclopropanecarboxylate (100 mg, 0.42 mmol) in CH₂Cl₂ (3 mL) at RT was added

trifluoroacetic acid (1 mL). The reaction mixture was stirred at RT for 10 min and concentrated in vacuo. The crude 2-(5-methylthien-2-yl)cyclopropanecarboxylic acid was used without further purification. M.S. (M+1): 182

Step 4:

5-Methyl-N-((1-((2-(5-methylthien-2-yl)cyclopropyl)carbonyl)piperidin-4-yl)methyl)pyrimidin-2-amine




2-(5-Methylthien-2-yl)cyclopropanecarboxylic acid was coupled to amine 5-methyl-N-(piperidin-4-ylmethyl)pyrimidin-2-amine (**EXAMPLE 144, STEP 3**) according to the procedure described for **EXAMPLE 144, STEP 4**, providing, after chromatography, 5-methyl-N-((1-((2-(5-methylthien-2-yl)cyclopropyl)carbonyl)piperidin-4-yl)methyl)pyrimidin-2-amine. M.S. (M+1): 371.


EXAMPLE 147:

N-((4-fluoro-1-((1R,2R)-2-phenylcyclopropyl)carbonyl)piperidin-4-

yl)methyl]-5-methylpyrimidin-2-amine

**Step 1:**


tert-Butyl 1-oxa-6-azaspiro[2.5]octane-6-carboxylate



To a solution of tert-butyl 4-oxopiperidine-1-carboxylate (0.50g, 2.51mmol) in THF/DMF (2:1, 6mL) at 60 °C was added trimethylsulfonium iodide (0.58g, 2.63mmol) and sodium t-butoxide (0.25g, 2.63mmol). The reaction mixture was stirred at 60 °C for 30min, cooled to RT and concentrated. Water was added and the mixture was extracted with EtOAc twice. The combined organics were dried over Na₂SO₄, filtered and concentrated. Purification on silica gel (3:1, hexanes:EtOAc) gave tert-butyl 1-oxa-6-azaspiro[2.5]octane-6-carboxylate as a clear oil that solidified upon standing.

Step 2:

Benzyl 4-fluoro-4-(hydroxymethyl)piperidine-1-carboxylate




To a solution of tert-butyl 4-oxopiperidine-1-carboxylate (7.0g, 32.8mmol) in CH₂Cl₂ (14mL) at -10 °C was added HF-pyridine (11.6mL, 82.1mmol) portionwise. The reaction mixture was stirred for 10min at -10 °C, warmed to RT. After stirring for 16h, the reaction was quenched with aqueous NaHCO₃ and extracted with CH₂Cl₂. The aqueous layer was concentrated to a white paste that was suspended in CH₂Cl₂ (100mL). N-benzylloxycarbonyloxysuccinimide (8.2g, 32.8mmol) was added and the mixture was stirred at RT for 3h. The reaction mixture was partitioned between EtOAc and H₂O, the organic layer was dried over Na₂SO₄, filtered and concentrated. Purification on silica gel (10:1 to 1:1 hexanes:EtOAc) gave benzyl 4-fluoro-4-(hydroxymethyl)piperidine-1-carboxylate as a clear oil. M.S. (M+1): 268

Step 3:

Benzyl 4-fluoro-4-(((methylsulfonyl)oxy)methyl)piperidine-1-carboxylate


To a solution of benzyl 4-(azidomethyl)-4-fluoropiperidine-1-carboxylate (1.5g, 5.1mmol) in THF (10mL) at RT was added water (0.92mL, 0.92mmol) and triphenylphosphine (4.3g, 15.4mmol). The reaction mixture was stirred for 60h, concentrated, dissolved in HCl (1M) and extracted with Et₂O four times. The aqueous layer was basified to pH 11 and extracted with EtOAc twice. The



To a solution of benzyl 4-fluoro-4-(hydroxymethyl)piperidine-1-carboxylate (1.0g, 3.7mmol) in CH₂Cl₂ (10mL) at RT was added methanesulfonyl chloride (0.29mL, 3.7mmol) and triethylamine (1.04mL, 7.5mmol). The reaction mixture was stirred at rt for 5min, and partitioned between ethyl acetate and water. The organic layer was dried over Na₂SO₄, filtered, concentrated and purified on silica gel (10:1 to 1:2 hexanes:EtOAc) to give benzyl 4-fluoro-4-(((methylsulfonyl)oxy)methyl)piperidine-1-carboxylate. M.S. (M+1): 346

Step 4:

Benzyl 4-(azidomethyl)-4-fluoropiperidine-1-carboxylate




To a solution of benzyl 4-fluoro-4-

(((methylsulfonyl)oxy)methyl)piperidine-1-carboxylate (1.3g, 3.7mmol) in DMF (10mL) at RT was added NaN₃ (2.4g, 37.0mmol). The reaction mixture was heated to 110 °C and stirred for 60h, cooled and partitioned between EtOAc and H₂O. The organic layer was dried over Na₂SO₄, filtered, concentrated and purified on silica gel (10:1 to 1:2 hexanes:EtOAc) to give benzyl 4-(azidomethyl)-4-fluoropiperidine-1-carboxylate. (0.86 g, 80% yield). M.S. (M+1): 293

Step 5:

Benzyl 4-(aminomethyl)-4-fluoropiperidine-1-carboxylate




To a solution of benzyl 4-(azidomethyl)-4-fluoropiperidine-1-carboxylate (1.5g, 5.1mmol) in THF (10mL) at RT was added water (0.92mL, 0.92mmol) and triphenylphosphine (4.3g, 15.4mmol). The reaction mixture was stirred for 60h, concentrated, dissolved in HCl (1M) and extracted with Et₂O four times. The aqueous layer was basified to pH 11 and extracted with EtOAc twice. The

organic layer was dried over Na_2SO_4 , filtered and concentrated. The crude mixture was chromatographed on silica gel (CH_2Cl_2 to 80:20:2 CH_2Cl_2 :MeOH/NH₄OH) to give benzyl 4-(aminomethyl)-4-fluoropiperidine-1-carboxylate. M.S. (M+1): 267

Step 6:

5 N-[(4-fluoro-1-[(1R,2R)-2-phenylcyclopropyl]carbonyl]piperidin-4-yl)methyl]-5-methylpyrimidin-2-amine




Benzyl 4-(aminomethyl)-4-fluoropiperidine-1-carboxylate was coupled to 2-chloro-5-methylpyrimidine, deprotected and coupled to (1R,2R)-2-phenylcyclopropanecarboxylic acid according to the procedure described for

10 **EXAMPLE 144**, STEPS 2,3,4, providing, after chromatography, N-[(4-fluoro-1-[(1R,2R)-2-phenylcyclopropyl]carbonyl]piperidin-4-yl)methyl]-5-methylpyrimidin-2-amine. M.S.(M+1): 369.

EXAMPLE 148:

15 4-[(4,5-Dimethyl-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



Step 1:

2-Chloro-4,5-dimethylpyrimidine




To 2-chloro-5-methylpyrimidine (**EXAMPLE 144**, STEP 1)(0.50g, 0.00389mol) in diethyl ether (12mL) at -30°C and under nitrogen was added dropwise 1.4M methylolithium (2.90mL, 0.00405mol) and the reaction allowed to stir at -30°C for 30min and at 0°C for 30min. The reaction was quenched with a solution of acetic

acid (0.242mL), water (0.039mL, and THF (0.8mL) and then a solution of DDQ (0.92g, 0.00405mol) in THF was added. The reaction was stirred 5min at rt, recooled to 0°C and 3N sodium hydroxide added. The reaction was allowed to stir at 0°C for 30min after which a thick oily precipitate formed. The organic supernatant was decanted and the residue washed with diethyl ether (2 x 20mL). The organic layers were dried over sodium sulfate, filtered through a pad of silica and the silica pad washed with diethyl ether. The filtrate was concentrated in vacuo to give the title compound as an oil. M.S.(M+1): 143.

$^1\text{H NMR}$ 400 MHz (δ , CDCl_3) δ : 2.22(s, 3H), 2.44(s, 3H), 8.27(s, 1H).

10 Step 2:


4-[(4,5-Dimethyl-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester



The above compound was prepared in a manner similar to that utilized for the preparation of **EXAMPLE 144**, STEP 2 using 2-chloro-4,5-dimethylpyrimidine in place of 2-chloro-5-methylpyrimidine, to give the title compound. M.S.(M+1): 355.

EXAMPLE 149:

20 4-[(4,5-Dimethyl-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid 4-methyl-benzyl ester




The title compound was prepared from 2-chloro-4,5-dimethylpyrimidine (EXAMPLE 148, STEP 1) and INTERMEDIATE 2A as described in a manner similar to that described in EXAMPLE 144, STEP 2 to give the title compound. M.S.(M+1): 369

The following EXAMPLES 150-152 were prepared from 4-((4,5-dimethyl-pyrimidin-2-ylamino)-methyl)-piperidine-1-carboxylic acid benzyl ester (EXAMPLE 148) by hydrogenation of the CBZ group as described in EXAMPLE 145, STEP 1, followed by coupling with the appropriate acid as described in EXAMPLE 145, STEP 2

EXAMPLE 150:


Trans {4-[(4,5-Dimethyl-pyrimidin-2-ylamino)-methyl]-piperidin-1-yl}-(2-phenyl-cyclopropyl)-methanone



M.S.(M+1): 365.

EXAMPLE 151:


{4-[(4,5-Dimethyl-pyrimidin-2-ylamino)-methyl]-piperidin-1-yl}-(2-(2-fluoro-phenyl)-cyclopropyl)-methanone



M.S.(M+1): 383.

EXAMPLE 152:


{4-[(4,5-Dimethyl-pyrimidin-2-ylamino)-methyl]-piperidin-1-yl}-(2-(2,6-difluoro-phenyl)-cyclopropyl)-methanone



M.S.(M+1): 401.


EXAMPLE 153:

5-bromo-N-((1-((1*R*,2*R*)-2-phenylcyclopropyl)carbonyl)piperidin-4-yl)methyl)pyrimidin-2-amine



Step 1:


[1-(2-Phenyl-cyclopropanecarbonyl)-piperidin-4-ylmethyl]-carbamic acid tert-butyl ester



A mixture of *tert*-butyl piperidin-4-ylmethylcarbamate (Epsilon, 0.80g, 3.73mmol), (1*R*,2*R*)-2-phenylcyclopropanecarboxylic acid (T. Riley et al, *J. Med. Chem.*, 15, 1187, 1972) (0.61g, 3.73mmol), EDC (1.07g, 5.60mmol) and HOBT (0.76g, 5.60mmol) in DMF (10mL) was stirred at RT for 2h. The reaction mixture was diluted with ethyl acetate (200mL), washed with aq. sat. NaHCO₃ (50mL), water (5 x 50mL), brine (50mL), dried over Na₂SO₄, filtered and concentrated. The residue was chromatographed on silica gel 60 (90g), eluting with 10:1-10:89-80 CH₂Cl₂:2-propanol:hexane to give the title compound, (1.22 g, 91.0 %). M.S. (M+1): 359.

Step 2:

10 (1-(((1*R*,2*R*)-2-phenylcyclopropyl[carbonyl]piperidin-4-yl)methyl)amine




A solution of *tert*-butyl (1-(((1*R*,2*R*)-2-

phenylcyclopropyl[carbonyl]piperidin-4-yl)methylcarbamate (1.00g, 2.79mmol) in TFA (3mL) and CH₂Cl₂ (3mL) was stirred at R.T. for 0.5h. The reaction mixture was then concentrated to give the title compound as a trifluoroacetate salt.

Step 3:

5-bromo-N-((1-(((1*R*,2*R*)-2-phenylcyclopropyl[carbonyl]piperidin-4-yl)methyl)pyrimidin-2-amine




A mixture of (1-(((1*R*,2*R*)-2-phenylcyclopropyl[carbonyl]piperidin-4-yl)methyl)amine trifluoroacetate salt (1.00g, 2.69mmol), 5-bromo-2-chloro-pyrimidine ((32779-36-5), 0.519g, 2.69mmol) and cesium carbonate (1.75g, 5.37mmol) in DMF (7mL) was heated at 100°C for 1.5h. The reaction mixture was cooled to RT, diluted with ethyl acetate (200mL), washed with water (5 x 20mL), and brine (10mL), then dried over Na₂SO₄, filtered and concentrated. The residue was chromatographed on silica gel 60, eluting with 10:1-20:89-70 CH₂Cl₂:2-propanol:hexane to give the title compound, (0.31 g, 28.1 %). M.S. (M+1): 416.

¹H NMR (400 MHz, CDCl₃): δ 8.26 (s, 2 H, Pyr), 7.28 (t, J = 7.3 Hz, 2 H, Ar), 7.19 (t, J = 7.7 Hz, 1 H, Ar), 7.11 (d, J = 7.6 Hz, 2 H, Ar), 5.21 (s, 1 H, NH), 4.64 (d, J = 11.9 Hz, 1 H, NCH₂), 4.13 (d, J = 12.9 Hz, 1 H, NCH₂), 3.31 (s, 2 H, NCH₂), 3.05 (q, J = 12.6 Hz, 1 H, NCH₂), 2.62 (t, J = 12.3 Hz, 1 H, NCH₂), 2.46 (brs, 1 H, ArCH), 1.98 (m, 1 H, CHCO), 1.87 (m, 1 H, CH₂CHCH₂), 1.80 (d, J = 13.1 Hz, 2 H, CHCH₂CH₂), 1.65 (s, 1 H, CHCH₂CH), 1.25 (s, 1 H, CHCH₂CH), 1.21 (m, 2 H, CHCH₂CH₂).

EXAMPLE 154:

10 N-((1-(((1*R*,2*R*)-2-Phenylcyclopropyl[carbonyl]piperidin-4-yl)methyl)-5-((trimethylsilyl)ethynyl)pyrimidin-2-amine




A mixture of 5-bromo-N-((1-(((1*R*,2*R*)-2-

phenylcyclopropyl[carbonyl]piperidin-4-yl)methyl)pyrimidin-2-amine (EXAMPLE 153) (0.300g, 0.722mmol), trimethylsilylacetylene (0.177g, 1.81mmol), Pd(PPh₃)₄ (0.083g, 0.072mmol), and copper iodide (0.007g, 0.036mmol) in DMSO (1mL) and diethylamine (1mL) was heated in a sealed tube at 100°C for 3h. The reaction mixture was cooled to RT, diluted with ethyl acetate (50mL), washed with water (10mL), and brine (10mL), then dried over Na₂SO₄, filtered and concentrated. The residue was chromatographed on silica gel 60 (35g), eluting with 10:1-10:89-80 CH₂Cl₂:2-propanol:hexane to give the title compound.

EXAMPLE 155:

25 5-Ethynyl-N-((1-(((1*R*,2*R*)-2-phenylcyclopropyl[carbonyl]piperidin-4-yl)methyl)pyrimidin-2-amine



A mixture of N-((1-(((1*R*,2*R*)-2-phenylcyclopropyl[carbonyl]piperidin-4-yl)methyl)-5-


((trimethylsilyl)ethylmethylpyrimidin-2-amine (EXAMPLE 154) (0.200g, 0.462mmol) and potassium carbonate (0.128g, 0.924mmol) in methanol (3mL) was stirred at RT for 0.5h. The reaction mixture was concentrated in vacuo, diluted with ethyl acetate (50mL), washed with water (20mL), and brine (10mL), then dried over Na₂SO₄, filtered and concentrated. The residue was chromatographed on silica gel 60 (35g), eluting with 10:1-20:89-70 CH₂Cl₂:2-propanol:hexane to give the title compound. M.S. (M+1): 361.

¹H NMR (400 MHz, CDCl₃): δ 8.37 (s, 2 H, Pyr), 7.28 (t, J = 7.1 Hz, 2 H, Ar), 7.19 (t, J = 7.0 Hz, 1 H, Ar), 7.11 (d, J = 7.4 Hz, 2 H, Ar), 5.38 (s, 1 H, NH), 4.65 (d, J = 12.6 Hz, 1 H, NCH₂), 4.14 (d, J = 14.1 Hz, 1 H, NCH₂), 3.36 (m, 2 H, NHCH₂), 3.17 (s, 1 H, CCH), 3.05 (q, J = 12.2 Hz, 1 H, NCH₂), 2.62 (t, J = 12.5 Hz, 1 H, NCH₂), 2.46 (brs, 1 H, ArCH), 1.97 (m, 1 H, CHCO), 1.86 (m, 1 H, CH₂CHCH₂), 1.80 (d, J = 12.5 Hz, 2 H, CHCH₂CH₂), 1.65 (m, 1 H, CHCH₂CH), 1.26 (m, 1 H, CHCH₂CH), 1.23 (m, 2 H, CHCH₂CH₂).

15


EXAMPLE 156:

2-(((1-((1R,2R)-2-phenylcyclopropyl)carbonyl)piperidin-4-yl)methyl)amino]pyrimidine-5-carbonitrile



20 Step 1:

4-chloro-2-(methylthio)pyrimidine-5-carbonitrile



A stirred solution of 4-hydroxy-2-(methylthio)pyrimidine-5-

carbonitrile (British patent GB901749) (1.00g, 5.98mmol) in POCl₃ (5mL) was heated to reflux for 2h. The reaction mixture was concentrated in vacuo and the residue quenched with ice (100g). The solution was then basified to pH 8 with sat. aq NaHCO₃ and extracted with ethyl acetate (3 x 50mL). The combined ethyl acetate layers were washed with water (20mL), brine (10mL), dried over Na₂SO₄, filtered and concentrated to give the title compound.

- 168 -

¹H NMR (400 MHz, CD₃OD): δ 8.84 (s, 1 H, Ar), 2.62 (s, 3 H, CH₃)

2-(Methylthio)pyrimidine-5-carbonitrile




To a stirred mixture of 4-chloro-2-(methylthio)pyrimidine-5-

carbonitrile (0.843g, 4.54mmol) and zinc dust (1.48g, 22.71mmol) in ethanol (7.5mL) and water (1.4mL) was slowly added acetic acid (0.29mL, 5.13mmol). The resulting reaction mixture was vigorously stirred for 3h. The solids were removed by filtration and the filtrate concentrated in vacuo. The residue was chromatographed on silica gel 60 (35g), eluting with 10:1-20:89-70 CH₂Cl₂:2-propanol:hexane to give the title compound. M.S. (M+1): 152.

¹H NMR (400 MHz, CDCl₃): δ 8.72 (s, 2 H, Ar), 2.61 (s, 3 H, CH₃).

Step 3:

2-(((1-((1R,2R)-2-Phenylcyclopropyl)carbonyl)piperidin-4-yl)methyl)amino]pyrimidine-5-carbonitrile



A mixture of (1-(((1R,2R)-2-phenylcyclopropyl)carbonyl)piperidin-4-yl)methylamine (EXAMPLE 153, STEP 2) (0.100g, 0.387mmol), 2-

(methylthio)pyrimidine-5-carbonitrile (0.059g, 0.387mmol) and cesium carbonate (0.252g, 0.774mmol) in DMF (1mL) was heated at 70°C for 1h. The reaction mixture was cooled to RT, diluted with ethyl acetate (50mL), washed with water (5 x 10mL), and brine (10mL), then dried over Na₂SO₄, filtered and concentrated. The residue was chromatographed on a reversed phase column, running 5 - 95% 0.1%TFA in CH₃CN/0.1%TFA in water to give the title compound as a TFA salt. M.S. (M+1): 362.


¹H NMR (400 MHz, CDCl₃): δ 8.52 (s, 1 H, Pyr), 8.45 (s, 1 H, Pyr), 7.28 (t, J = 7.9 Hz, 2 H, Ar), 7.20 (t, J = 6.6 Hz, 1 H, Ar), 7.11 (d, J = 7.3 Hz, 2 H, Ar), 5.78 (s, 1 H, NH), 4.66 (d, J = 12.2 Hz, 1 H, NCH₂), 4.15 (d, J = 13.1 Hz, 1 H, NCH₂), 3.41 (m, 2 H, NHCH₂), 3.06 (q, J = 12.4 Hz, 1 H, NCH₂), 2.62 (t, J = 12.7 Hz, 1 H, NCH₂), 2.47 (brs, 1 H, ArCH), 1.96 (m, 1 H, CHCO), 1.91 (m, 1 H,

- 169 -

CH_3CHCH_2 , 1.80 (d, $J = 13.2$ Hz, 2 H, CHCH_2CH_3), 1.65 (s, 1 H, CHCH_2CH_3), 1.27 (s, 1 H, CHCH_2CH_3), 1.25 (m, 2 H, CHCH_2CH_3).

EXAMPLE 157:

5 5-Ethyl-N-((1-(((1*R*,2*R*)-2-phenylcyclopropyl)carbonyl)piperidin-4-yl)methyl)pyrimidin-2-amine



The stirred reaction mixture of (1-(((1*R*,2*R*)-2-


phenylcyclopropyl)carbonyl)piperidin-4-yl)methylamine (EXAMPLE 154, STEP 2) (0.100g, 0.387mmol), 2-chloro-5-ethyl-pyrimidine ([111196-81-7], 0.055g,

0.387mmol) and cesium carbonate (0.252g, 0.774mmol) in DMF (5mL) was heated at 150°C for 7h. The reaction mixture was cooled to RT, diluted with ethyl acetate (100mL), washed with water (5 x 20mL), and brine (10mL), then dried over Na_2SO_4 , filtered and concentrated. The residue was chromatographed on silica gel 60, eluting with 10:1-20:89-70 CH_2Cl_2 :2-propanol:hexane to give the title compound. M.S. (M+1): 365.

^1H NMR (400 MHz, CDCl_3): δ 8.13 (s, 2 H, Pyr), 7.28 (t, $J = 6.6$ Hz, 2 H, Ar), 7.19 (t, $J = 7.2$ Hz, 1 H, Ar), 7.11 (d, $J = 7.3$ Hz, 2 H, Ar), 5.05 (s, 1 H, NH), 4.64 (d, $J = 12.9$ Hz, 1 H, NCH_2), 4.14 (d, $J = 12.8$ Hz, 1 H, NCH_2), 3.32 (s, 2 H, NHCH_2), 3.05 (q, $J = 12.5$ Hz, 1 H, NCH_2), 2.62 (t, $J = 12.6$ Hz, 1 H, NCH_2), 2.46 (q, $J = 7.5$ Hz, 2 H, CH_2CH_3), 2.43 (brs, 1 H, ArCH), 1.97 (m, 1 H, CHCO), 1.86 (m, 1 H, CH_2CHCH_2), 1.82 (d, $J = 13.5$ Hz, 2 H, CHCH_2CH_3), 1.64 (m, 1 H, CHCH_2CH_3), 1.26 (m, 1 H, CHCH_2CH_3), 1.22 (m, 2 H, CHCH_2CH_3), 1.19 (t, $J = 7.6$ Hz, 3 H, CH_3CH_3).

EXAMPLE 158:

5-(Cyclopropylethynyl)-N-((1-(((1*R*,2*R*)-2-phenylcyclopropyl)carbonyl)piperidin-4-yl)methyl)pyrimidin-2-amine



A mixture of 5-bromo-N-((1-(((1*R*,2*R*)-2-phenylcyclopropyl)carbonyl)piperidin-4-yl)methyl)pyrimidin-2-amine (EXAMPLE 153) (0.050g, 0.120mmol), ethynylcyclopropane (0.020g, 0.301mmol), $\text{Pd}(\text{PPh}_3)_4$ (0.014g, 0.012mmol), copper iodide (0.001g, 0.006mmol) in DMSO (1mL) and diethylamine (1mL) was heated in a sealed tube at 100°C for 3h. The reaction mixture


was cooled to RT, diluted with ethyl acetate (50mL), washed with water (10mL), and brine (10mL), then dried over Na_2SO_4 , filtered and concentrated. The residue was chromatographed on a reverse phase column, running 5 - 95% 0.1%TFA in $\text{CH}_3\text{CN}/0.1\%$ TFA in water to give the title compound as a TFA salt. M.S. (M+1):

10 401.

^1H NMR (400 MHz, CDCl_3): δ 8.56 (s, 1 H, Pyr), 8.02 (s, 1 H, Pyr), 7.28 (t, $J = 7.4$ Hz, 2 H, Ar), 7.20 (t, $J = 7.2$ Hz, 1 H, Ar), 7.11 (d, $J = 7.5$ Hz, 2 H, Ar), 4.64 (d, $J = 13.0$ Hz, 1 H, NCH_2), 4.15 (d, $J = 11.7$ Hz, 1 H, NCH_2), 3.50 (s, 1 H, NHCH_2), 3.41 (s, 1 H, NHCH_2), 3.07 (q, $J = 12.8$ Hz, 1 H, NCH_2), 2.64 (t, $J = 12.7$ Hz, 1 H, NCH_2), 2.47 (brs, 1 H, ArCH), 1.98 (m, 1 H, CHCO), 1.92 (m, 1 H, CH_2CHCH_2), 1.79 (d, $J = 13.6$ Hz, 2 H, CHCH_2CH_3), 1.64 (s, 1 H, CHCH_2CH_3), 1.44 (m, 1 H, CHCH_2CH_3), 1.28 (m, 1 H, CHCH_2CH_3), 1.26 (m, 2 H, CHCH_2CH_3), 0.83 (m, 4 H, CH_3).

EXAMPLE 159:

N-((1-(((1*R*,2*R*)-2-Phenylcyclopropyl)carbonyl)piperidin-4-yl)methyl)-5-(phenylethynyl)pyrimidin-2-amine



A mixture of 5-bromo-N-((1-(((1*R*,2*R*)-2-

phenylcyclopropyl)carbonyl)piperidin-4-yl)methyl)pyrimidin-2-amine (EXAMPLE 153) (0.200g, 0.482mmol), ethynylbenzene (0.123g, 0.132mmol), $\text{Pd}(\text{PPh}_3)_4$ (0.056g, 0.048mmol), and copper iodide (0.005g, 0.024mmol) in DMSO (1mL) and diethylamine (1mL) was heated in a sealed tube at 100°C for 3h. The reaction mixture was cooled to RT, diluted with ethyl acetate (50mL), washed with water (10mL), and brine (10mL), then dried over Na_2SO_4 , filtered and concentrated. The residue was chromatographed on a reversed phase column, running 5 - 95% 0.1%TFA in

CH₃CN/0.1%TFA in water to give the title compound as a TFA salt. M.S. (M+1): 437.







¹H NMR (400 MHz, CDCl₃): 8.8.50 (brs, 2 H, Pyr), 7.50 (m, 2 H, Ar), 7.38 (m, 3 H, Ar), 7.28 (t, J = 7.9 Hz, 2 H, Ar), 7.20 (t, J = 7.3 Hz, 1 H, Ar), 7.11 (d, J = 7.5 Hz, 2 H, Ar), 4.66 (d, J = 11.7 Hz, 1 H, NCH₂), 4.16 (d, J = 13.2 Hz, 1 H, NCH₂), 3.46 (m, 2 H, NHCH₂), 3.10 (m, 1 H, NCH₂), 2.64 (t, J = 11.9 Hz, 1 H, NCH₂), 2.47 (brs, 1 H, ArCH), 1.97 (m, 1 H, CHCO), 1.93 (m, 1 H, CH₂CHCH₃), 1.81 (d, J = 11.5 Hz, 2 H, CHCH₂CH₂), 1.65 (brs, 1 H, CHCH₂CH), 1.28 (m, 1 H, CHCH₂CH), 1.25 (m, 2 H, CHCH₂CH₃).







EXAMPLE 160 TO EXAMPLE 180


The following examples were prepared by coupling the appropriate amine (EXAMPLE 143, STEP 1, EXAMPLE 145, STEP 1, or piperidin-4-ylmethyl-pyrimidin-2-yl-amine which was prepared in a manner similar to that described for EXAMPLE 143, STEP 1, replacing 2-chloro-5-methylpyrimidine with 2-chloropyrimidine in STEP 1) with the appropriately substituted trans phenylcyclopropanecarboxylic acid (prepared in a similar manner to that described in (EXAMPLE 136).

EX.	Structure	Name	M.S. (M+1)
160		[4-(Pyrimidin-2-ylaminomethyl)-piperidin-1-yl]-(2-p-tolyl-cyclopropyl)-methanone	351.2
161		[4-(Pyrimidin-2-ylaminomethyl)-piperidin-1-yl]-(2-o-tolyl-cyclopropyl)-methanone	351.4

EX.	Structure	Name	M.S. (M+1)
162		[4-(Pyrimidin-2-ylaminomethyl)-piperidin-1-yl]-(2-m-tolyl-cyclopropyl)-methanone	351.4
163		[2-(4-Fluoro-phenyl)-cyclopropyl]-[4-(pyrimidin-2-ylaminomethyl)-piperidin-1-yl]-methanone	355.2
164		[2-(4-Chloro-phenyl)-cyclopropyl]-[4-(pyrimidin-2-ylaminomethyl)-piperidin-1-yl]-methanone	371.1
165		[2-(3-Chloro-phenyl)-cyclopropyl]-[4-(pyrimidin-2-ylaminomethyl)-piperidin-1-yl]-methanone	371.3
166		[2-(3-Fluoro-phenyl)-cyclopropyl]-[4-(pyrimidin-2-ylaminomethyl)-piperidin-1-yl]-methanone	355.3
167		[2-(2-Methoxy-phenyl)-cyclopropyl]-[4-(pyrimidin-2-ylaminomethyl)-piperidin-1-yl]-methanone	367.3


EX.	Structure	Name	M.S. (M+1)
168		[2-(3-Methoxy-phenyl)-cyclopropyl]-[4-(pyrimidin-2-ylamino-methyl)-piperidin-1-yl]-methanone	367.3
169		[2-(2,6-Difluoro-phenyl)-cyclopropyl]-[4-(pyrimidin-2-ylamino-methyl)-piperidin-1-yl]-methanone	373.3
170		[2-(2,4-Difluoro-phenyl)-cyclopropyl]-[4-(pyrimidin-2-ylamino-methyl)-piperidin-1-yl]-methanone	373.4
171		(2-Phenyl-cyclopropyl)-[4-(pyrimidin-2-ylamino-methyl)-piperidin-1-yl]-methanone	337.2
172		[2-(2,3-Difluoro-phenyl)-cyclopropyl]-[4-(pyrimidin-2-ylamino-methyl)-piperidin-1-yl]-methanone	373.3
173		[4-(5-Methyl-pyrimidin-2-ylamino)-methyl]-piperidin-1-yl-(2-m-tolyl-cyclopropyl)-methanone	365.4

EX.	Structure	Name	M.S. (M+1)
174		[4-(5-Methyl-pyrimidin-2-ylamino)-methyl]-piperidin-1-yl-(2-o-tolyl-cyclopropyl)-methanone	365.3
175		[2-(2-Fluoro-phenyl)-cyclopropyl]-[4-(5-methyl-pyrimidin-2-ylamino)-methyl]-piperidin-1-yl]-methanone	369.3
176		[2-(2,3-Difluoro-phenyl)-cyclopropyl]-[4-(5-methyl-pyrimidin-2-ylamino)-methyl]-piperidin-1-yl]-methanone	387.3
177		[2-(2,6-Difluoro-phenyl)-cyclopropyl]-[4-(5-methyl-pyrimidin-2-ylamino)-methyl]-piperidin-1-yl]-methanone	387.3
178		[4-(5-Fluoro-pyrimidin-2-ylamino)-methyl]-piperidin-1-yl-(2-pentafluorophenyl-cyclopropyl)-methanone	445.3
179		[4-(5-Fluoro-pyrimidin-2-ylamino)-methyl]-piperidin-1-yl-(2-o-tolyl-cyclopropyl)-methanone	369.3

EX.	Structure	Name	M.S. (M+1)
180		(4-((5-Fluoro-pyrimidin-2-ylamino)-methyl)-piperidin-1-yl)-(2-m-tolyl-cyclopropyl)-methanone	369.4

EXAMPLE 181:

4-((3-Fluoro-pyridin-2-ylamino)-methyl)-piperidine-1-carboxylic acid benzyl ester




A solution of 2-chloro-3-fluoropyridine (prepared in a manner similar to that described by W.J. Link, R.F. Borne and F.L. Setliff, *J. Heterocyclic Chem.* 4, 641-3, 1967) (131mg, 1mmol), benzyl 4-(aminomethyl)piperidine-1-carboxylate (EXAMPLE 13, STEP 1) (248mg, 1mmol) and diisopropylethylamine (129mg, 1mmol) were heated to reflux in 2-methoxyethanol for 2 days under nitrogen. The reaction mixture was concentrated, partitioned between ethyl acetate and water, the organic layer washed with brine, dried over anhydrous sodium sulfate and solvent evaporated to give the crude product purified by chromatography on silica.

M.S. (M+1): 344.3

EXAMPLE 182:


(4-((3-Fluoro-pyridin-2-ylamino)-methyl)-piperidin-1-yl)-(2-phenyl-cyclopropyl)-methanone



The title compound was prepared from 4-((3-fluoro-pyridin-2-ylamino)-methyl)-piperidine-1-carboxylic acid benzyl ester (EXAMPLE 181) in a similar manner to that described in EXAMPLE 145). M.S. (M+1): 354.3

EXAMPLE 183:

4-((3-Fluoro-pyridin-2-ylamino)-methyl)-piperidine-1-carboxylic acid benzyl ester

**Step 1:****2-Chloro-3-fluoropyridine**

Prepared in a manner similar to that described by W.J. Link, R.F. Borne and F.L. Setliff, *J. Heterocyclic Chem.* 4, 641-3, (1967).

Step 2:

A mixture of 2mmol of 4-aminomethyl-piperidine-1-carboxylic acid benzyl ester, 1mmol of 2-chloro-3-fluoropyridine, and 1mmol of tributylamine were heated to reflux in 2mL of cyclohexanol for 3 days (or 2-methoxyethanol for 14 days) under nitrogen. Preparative TLC eluting with 75:25 ether:hexane gave 4-((3-fluoro-pyridin-2-ylamino)-methyl)-piperidine-1-carboxylic acid benzyl ester:

¹H NMR (CDCl₃) δ: 7.85 (1H, d), 7.4-7.35 (5H, m), 7.1 (1H, dd), 6.5 (1H, m), 5.15 (2H, s), 4.65 (1H, br m), 4.2 (2H, br s), 3.4 (2H, br m), 2.8 (2H, br m), 1.8 (3H, m), 1.2 (2H, m). Mass spec.: 344.32 (M+1).

A lower band gave 4-((2-chloro-pyridin-3-ylamino)-methyl)-piperidine-1-carboxylic acid benzyl ester:


¹H NMR (CDCl₃) δ: 7.7 (1H, d), 7.4-7.35 (5H, m), 7.1 (1H, dd), 6.82 (1H, d), 5.15 (2H, s), 4.4 (1H, br m), 4.2 (2H, br s), 3.05 (2H, m), 2.8 (2H, br m), 1.8 (3H, m), 1.2 (2H, m). Mass spec.: 360.29 (M+1).

Alternatively, the use of 2,3-difluoropyridine [Finger, G. C.; Starr, L. D.; Roe, A.; Link, W. J., *J. Organic Chem.* 27, 3965-68, 1962.] in place of 2-chloro-3-fluoropyridine in refluxing 2-butanol gave higher yields of product without the 4-

[(2-chloro-pyridin-3-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester by-product.

EXAMPLE 184:

5 [R,R] 4-[(3-Fluoro-pyridin-2-ylamino)-methyl]-piperidin-1-yl)-(2-phenyl-cyclopropyl)-methanone




Prepared from 4-[(3-fluoro-pyridin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester by hydrogenolysis of the benzyloxycarbonyl group followed by EDC, HOBt coupling with [R,R] trans-2-phenyl-1-cyclopropanecarboxylic acid in DMF in the usual manner such as described previously in EXAMPLE 143 above:

15 ¹H NMR (CDCl₃) 8.75 (1H, d), 7.35 (2H, m), 7.2 (1H, dd), 7.1 (3H, m), 6.5 (1H, m), 4.65 (2H, br m), 4.18 (1H, br d), 3.4 (2H, br m), 3.1 (1H, complex m), 2.6 (1H, m), 2.45 (1H, m), 2.0-1.8 (4H, m), 1.62 (1H, m), 1.2 (3H, m). Mass spec.: 354.35 (M+1).

EXAMPLE 185:

20 4-[(4-Methyl-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester




A mixture of 1.6mmol of 4-aminomethyl-piperidine-1-carboxylic acid benzyl ester, 2.4mmol of 2-methanesulfonyl-4-methylpyrimidine, and 3mmol of N,N-diethylethylamine were heated to reflux in either 5mL of 2-butanol for 24h under nitrogen. Preparative TLC eluting with ethyl acetate gave 460mg of 4-[(3-fluoro-pyridin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester.

¹H NMR (CDCl₃) 8.8.1 (1H, d), 7.4-7.35 (5H, m), 6.4 (1H, d), 5.15 (2H, s), 4.2 (2H, br s), 3.35 (2H, m), 2.8 (2H, br m), 2.3 (3H, s), 1.8 (4H, m), 1.2 (2H, m). Mass spec.: 341.4 (M+1).

EXAMPLE 186:


5 [R,R] 4-[(4-Methyl-pyrimidin-2-ylamino)-methyl]-piperidin-1-yl)-(2-phenyl-cyclopropyl)-methanone



Prepared from 4-[(4-methyl-pyrimidin-2-ylamino)-methyl]-piperidine-1-carboxylic acid benzyl ester by hydrogenolysis of the benzyloxycarbonyl group followed by EDC, HOBt coupling with [R,R] trans-2-phenyl-1-cyclopropanecarboxylic acid in DMF as described above in EXAMPLE 143.

Preparative TLC using 90:10 ethyl acetate: methanol gave the product:
15 ¹H NMR (CDCl₃) 8.8.1 (1H, d), 7.35 (2H, m), 7.2 (1H, dd), 7.1 (2H, m), 6.4 (1H, d), 5.3 (1H, br m), 4.6 (1H, br d), 4.15 (1H, br d), 3.35 (2H, m), 3.05 (1H, dd), 2.6 (1H, t), 2.45 (1H, m), 2.3 (3H, s), 2.0 (1H, m), 1.8 (4H, m), 1.6 (1H, s), 1.2 (4H, m). Mass spec.: 351.4 (M+1).

EXAMPLE 187:



20 (+)-N-((8-((trans-2-phenylcyclopropyl)carboxyl)-8-azabicyclo[3.2.1]oct-3-eno-yl)methyl)pyrimidin-2-amine was prepared similarly as described previously above.

CCR3

2. The compound according to Claim 1, or pharmaceutically acceptable salts thereof, wherein
NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom; and
B is $\text{aryl}(\text{CH}_2)_{0-3}-\text{O}-\text{C}(\text{O})-$, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C_1 -4alkyl, C_3 -6cycloalkyl, C_1 -4alkoxy, trifluoromethyl, bromo, fluoro, or chloro.

3. The compound according to Claim 2, or pharmaceutically acceptable salts thereof, wherein
HetAr is a 6 membered heteroaromatic ring containing 1 nitrogen ring atom;

HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C_1 -4alkyl, C_1 -4alkoxy, C_2 -4alkynyl, trifluoromethyl, hydroxy, hydroxy C_1 -4alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, $\text{N}(\text{C}_0$ -4alkyl)(C_0 -4alkyl), nitro, $(\text{C}_1$ -2alkyl)(C_1 -2alkyl) NCH_2- , $(\text{C}_1$ -2alkyl) HNCH_2- , $\text{Si}(\text{CH}_3)_3-\text{C}-$, or $\text{NH}_2\text{C}(\text{O})-$.

4. The compound according to Claim 2, or pharmaceutically acceptable salts thereof, wherein

HetAr is an isoxazolyl optionally substituted with 1 or 2 substituents, each substituent independently is C_1 -4alkyl, C_1 -4alkoxy, C_2 -4alkynyl, trifluoromethyl, hydroxy, hydroxy C_1 -4alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, $\text{N}(\text{C}_0$ -4alkyl)(C_0 -4alkyl), nitro, $(\text{C}_1$ -2alkyl)(C_1 -2alkyl) NCH_2- , $(\text{C}_1$ -2alkyl) HNCH_2- , $\text{Si}(\text{CH}_3)_3-\text{C}-$, or $\text{NH}_2\text{C}(\text{O})-$.

5. The compound according to Claim 2, or pharmaceutically acceptable salts thereof, wherein

HetAr is a thiadiazolyl optionally substituted with 1 or 2 substituents, each substituent independently is C_1 -4alkyl, C_1 -4alkoxy, C_2 -4alkynyl, trifluoromethyl, hydroxy, hydroxy C_1 -4alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, $\text{N}(\text{C}_0$ -4alkyl)(C_0 -4alkyl), nitro, $(\text{C}_1$ -2alkyl)(C_1 -2alkyl) NCH_2- , $(\text{C}_1$ -2alkyl) HNCH_2- , $\text{Si}(\text{CH}_3)_3-\text{C}-$, or $\text{NH}_2\text{C}(\text{O})-$.

4alkyl)(C_0 -4alkyl), nitro, $(\text{C}_1$ -2alkyl)(C_1 -2alkyl) NCH_2- , $(\text{C}_1$ -2alkyl) HNCH_2- , $\text{Si}(\text{CH}_3)_3-\text{C}-$, or $\text{NH}_2\text{C}(\text{O})-$.

6. The compound according to Claim 2, or pharmaceutically acceptable salts thereof, wherein
HetAr is a 5 membered heteroaromatic ring containing 2 nitrogen ring atoms;

HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C_1 -4alkyl, C_1 -4alkoxy, C_2 -4alkynyl, trifluoromethyl, hydroxy, hydroxy C_1 -4alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, $\text{N}(\text{C}_0$ -4alkyl)(C_0 -4alkyl), nitro, $(\text{C}_1$ -2alkyl)(C_1 -2alkyl) NCH_2- , $(\text{C}_1$ -2alkyl) HNCH_2- , $\text{Si}(\text{CH}_3)_3-\text{C}-$, or $\text{NH}_2\text{C}(\text{O})-$.

7. The compound according to Claim 2, or pharmaceutically acceptable salts thereof, wherein
HetAr is quinolinyl optionally substituted with 1 or 2 substituents, each substituent independently is C_1 -4alkyl, C_1 -4alkoxy, C_2 -4alkynyl, trifluoromethyl, hydroxy, hydroxy C_1 -4alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, $\text{N}(\text{C}_0$ -4alkyl)(C_0 -4alkyl), nitro, $(\text{C}_1$ -2alkyl)(C_1 -2alkyl) NCH_2- , $(\text{C}_1$ -2alkyl) HNCH_2- , $\text{Si}(\text{CH}_3)_3-\text{C}-$, or $\text{NH}_2\text{C}(\text{O})-$.

8. The compound according to Claim 2, or pharmaceutically acceptable salts thereof, wherein
HetAr is purinyl optionally substituted with 1 or 2 substituents, each substituent independently is C_1 -4alkyl, C_1 -4alkoxy, C_2 -4alkynyl, trifluoromethyl, hydroxy, hydroxy C_1 -4alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, $\text{N}(\text{C}_0$ -4alkyl)(C_0 -4alkyl), nitro, $(\text{C}_1$ -2alkyl)(C_1 -2alkyl) NCH_2- , $(\text{C}_1$ -2alkyl) HNCH_2- , $\text{Si}(\text{CH}_3)_3-\text{C}-$, or $\text{NH}_2\text{C}(\text{O})-$.

9. The compound according to Claim 2, or pharmaceutically acceptable salts thereof, wherein

HeAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring

atoms;

HeAr is optionally substituted with 1 or 2 substituents, each

substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₄alkyl)(CO₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

10 The compound according to Claim 2, or pharmaceutically acceptable salts thereof, wherein

HeAr is thiazolyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₄alkyl)(CO₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

11 The compound according to Claim 2, or pharmaceutically acceptable salts thereof, wherein

HeAr is pteridinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₄alkyl)(CO₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

12 The compound according to Claim 2, or pharmaceutically acceptable salts thereof, wherein

HeAr is pyrrolopyrimidinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₄alkyl)(CO₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

13 The compound according to Claim 2, or pharmaceutically acceptable salts thereof, wherein

HeAr is a imidazopyridinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₄alkyl)(CO₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

14 The compound according to Claim 2, or pharmaceutically acceptable salts thereof, wherein

HeAr is benzimidazolyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₄alkyl)(CO₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

15 The compound according to Claim 1, or pharmaceutically acceptable salts thereof, wherein

NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom; and

B is aryl(CH₂)₁₋₃-SO₂-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro.

16 The compound according to Claim 15, or pharmaceutically acceptable salts thereof, wherein

HeAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring atoms;

HeAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyc₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₄alkyl)(CO₄alkyl),

nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

17. The compound according to Claim 15, or pharmaceutically

acceptable salts thereof, wherein

HetAr is quinazolinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkenyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

18. The compound according to Claim 15, or pharmaceutically

acceptable salts thereof, wherein

HetAr is purinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkenyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

19. The compound according to Claim 15, or pharmaceutically

acceptable salts thereof, wherein

HetAr is imidazopyridinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkenyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

20. The compound according to Claim 15, or pharmaceutically

acceptable salts thereof, wherein

HetAr is a 6 membered heteroaromatic ring containing 1 nitrogen ring atom; and

HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkenyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

21. The compound according to Claim 1, or pharmaceutically

acceptable salts thereof, wherein

NonAr is a nonaromatic 5 membered ring containing 1 nitrogen ring atom; and

B is aryl(CH₂)₀₋₃-O-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro.

22. The compound according to Claim 21, or pharmaceutically

acceptable salts thereof, wherein

HetAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring atoms;

HetAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkenyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

23. The compound according to Claim 21, or pharmaceutically

acceptable salts thereof, wherein

HetAr is pteridinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkenyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfanyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)NCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

24. The compound according to Claim 21, or pharmaceutically acceptable salts thereof, wherein

HeAr is purinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₄alkyl)(CO₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

25. The compound according to Claim 21, or pharmaceutically acceptable salts thereof, wherein

HeAr is benzimidazolyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₄alkyl)(CO₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

26. The compound according to Claim 1, or pharmaceutically acceptable salts thereof, wherein

NonAr is an aza bicyclo octane ring; and
B is aryl(CH₂)₃-O-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro.

27. The compound according to Claim 26, or pharmaceutically acceptable salts thereof, wherein

HeAr is a 6 membered heteroaromatic ring containing 1 nitrogen ring atom; and

HeAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₄alkyl)(CO₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

28. The compound according to Claim 26, or pharmaceutically acceptable salts thereof, wherein

HeAr is purinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₄alkyl)(CO₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

29. The compound according to Claim 26, or pharmaceutically acceptable salts thereof, wherein

HeAr is a 6 membered heteroaromatic ring containing 2 nitrogen ring atom; and

HeAr is optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO₄alkyl)(CO₄alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

30. The compound according to Claim 1, or pharmaceutically acceptable salts thereof, wherein

NonAr is an aza bicyclo octane ring; and

B is aryl(CH₂)₃-SO₂-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro.



31. The compound according to Claim 1, or pharmaceutically acceptable salts thereof, wherein



NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom; and


B is heteroaryl(CH₂)₃-C(O)-, wherein the heteroaryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro.

32. The compound according to Claim 1, or pharmaceutically acceptable salts thereof, wherein
NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom; and
B is aryl(CH₂)₁₋₃-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro.
33. The compound according to Claim 1, or pharmaceutically acceptable salts thereof, wherein
NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom; and
B is aryl(cyclopropyl)-C(O)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro.
34. The compound according to Claim 33, or pharmaceutically acceptable salts thereof, wherein
HeAr is pyridyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.
35. The compound according to Claim 33, or pharmaceutically acceptable salts thereof, wherein
HeAr is pyrazinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.

36. The compound according to Claim 33, or pharmaceutically acceptable salts thereof, wherein
HeAr is pyridazinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.
37. The compound according to Claim 33, or pharmaceutically acceptable salts thereof, wherein
HeAr is pyrimidinyl optionally substituted with 1 or 2 substituents, each substituent independently is C₁₋₄alkyl, C₁₋₄alkoxy, C₂₋₄alkynyl, trifluoromethyl, hydroxy, hydroxyC₁₋₄alkyl, fluoro, chloro, bromo, iodo, cyano, methylsulfonyl, cyclopropylethynyl-, phenylethynyl-, heteroarylethynyl-, N(CO-4alkyl)(CO-4alkyl), nitro, (C₁₋₂alkyl)(C₁₋₂alkyl)HNCH₂-, (C₁₋₂alkyl)HNCH₂-, Si(CH₃)₃-C-, or NH₂C(O)-.
38. The compound according to Claim 1, or pharmaceutically acceptable salts thereof, wherein
NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom; and
B is heteroaryl(CH₂)₁₋₃-O-C(O)-, wherein the heteroaryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro.
39. The compound according to Claim 1, or pharmaceutically acceptable salts thereof, wherein
NonAr is a nonaromatic 6 membered ring containing 1 nitrogen ring atom; and
B is aryl(CH₂)₁₋₃-NH-C(NCN)-, wherein the aryl is optionally substituted by 1-5 substituents, each substituent independently is C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, trifluoromethyl, bromo, fluoro, or chloro.
40. The compound according to Claim 1, wherein said compound is









or a pharmaceutically acceptable salt thereof.

41. The compound according to Claim 1, wherein said compound is




or a pharmaceutically acceptable salt thereof.

42. The compound according to Claim 1, wherein said compound is




or a pharmaceutically acceptable salt thereof.


43. The compound according to Claim 1, wherein said compound is



or a pharmaceutically acceptable salt thereof.



44. The compound according to Claim 1, wherein said compound is






or a pharmaceutically acceptable salt thereof.

45. The compound according to Claim 1, wherein said compound is




or a pharmaceutically acceptable salt thereof.

46. The compound according to Claim 1, wherein said compound is



or a pharmaceutically acceptable salt thereof.

47. The compound according to Claim 1, wherein said compound is



or a pharmaceutically acceptable salt thereof.

48. A pharmaceutical composition comprising an inert carrier and an effective amount of a compound according to claim 1.
49. The pharmaceutical composition according to claim 48 useful for the treatment of pain.
50. The pharmaceutical composition according to claim 48 useful for the treatment of migraine, depression, anxiety, schizophrenia, Parkinson's disease, or stroke.
51. A method of treating pain comprising a step of administering to one in need of such treatment an effective amount of a compound according to claim 1.
52. A method of treating migraine, depression, anxiety, schizophrenia, Parkinson's disease, or stroke comprising a step of administering to one in need of such treatment an effective amount of a compound according to claim 1.

- 204 -

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 02/05226	
CLASSIFICATION OF SUBJECT MATTER IPC 7 C07D401/12 C07D471/08 C07D487/04 A61K31/44 A61P29/02	
According to International Patent Classification (IPC) or to both national classification and IPC	
B. FIELD OF SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D	
Documentation searched other than minimum documentation to the extent that such documents are included in the data searched	
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, BEILSTEIN Data, CHEM ABS Data, MPI Data, PAJ	
C. DOCUMENTS CONSIDERED TO BE RELEVANT	
Category *	Relevant to claim No.
X	US 3 184 462 A (SCARBOROUGH HOMER C ET AL) 18 May 1965 (1965-05-18) equations 1 and 4 examples II, XV, XXXIII, XXXVII
A	US 6 124 323 A (BIGGEE CHRISTOPHER F ET AL) 26 September 2000 (2000-09-26) claims 1, 2; table 2 -/-
Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.	
* Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance "B" document published after the international filing date but prior to the priority date and not in conflict with the application but which may be considered the principle or theory underlying the invention "C" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to be an improvement over the prior art when the document is taken alone "D" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one of more other such documents in the art "E" document published prior to the international filing date but later than the priority date claimed "F" document member of the same patent family	
Date of the actual completion of the international search 4 June 2002	
Date of mailing of the international search report 19/06/2002	
Name and mailing address of the ISA European Patent Office, P.O. Box 1 NL - 2200 HV Rijswijk Tel. (+31-70) 340-3040, Telex 31 631 epo NL Fax (+31-70) 340-3010	Authorized officer SeeImann, I

Form PCT/ISA/210 (second sheet) July 1992

INTERNATIONAL SEARCH REPORT

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT	
Category*	Citation of document, with indication, where appropriate, of the relevant passages
A	<p>Z.-L. ZHOU, ET AL.: "4-Hydroxy-1'-2-(4-hydroxyphenoxy)ethyl)-4 -(4-methylbenzyl)piperidine: A novel, potent, and selective NR1/2B nmda receptor antagonist" J. MED. CHEM., vol. 42, 1999, pages 2993-3000, XP002201030 cited in the application Results and Discussion abstract</p>
	1-52

from PCT/US02/05226 (publication of second sheet) (July 1997)

INTERNATIONAL SEARCH REPORT

Information on patent family members	
Patent document cited in search report	Publication date
US 3184462	A 18-05-1965
	CH 421114 A 30-09-1966
	FR 1262 M 09-02-1968
	FR 1512306 A 22-04-1964
	GB 956255 A 22-04-1964
	GB 956254 A 22-04-1964
	NL 265777 A
US 6124323	A 26-09-2000
	AU 717185 B2 16-03-2000
	AU 1689997 A 17-07-1997
	BG 63380 B1 31-12-2001
	BG 102562 A 30-04-1999
	BR 9612686 A 20-07-1999
	CA 2240275 A1 03-07-1997
	CZ 9801802 A3 13-01-1999
	EP 0869792 A2 14-10-1998
	HU 9901296 A2 30-08-1999
	NO 982870 A 24-08-1998
	NZ 330624 A 27-03-2000
	PL 327493 A1 21-12-1998
	SK 82498 A3 11-02-1999
	WO 9723216 A1 03-07-1997
	ZA 9610745 A 24-06-1997

from PCT/US02/05226 (patent family sheet) (July 1997)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.